

Variation in Sensitivity Among Some Isolates of *Macrophomina phaseolina* Isolated from Cotton Roots to Flutolanil Fungicide

Yehia A.-G. Mahmoud^{1*}, A.A. Aly², M.R. Omar² and Abdel-Wahab A. Ismail²

¹Botany Department, Mycology Research Lab., Tanta 31527, Egypt

²Plant Pathology Research Institute, Agriculture Research Center, Giza, Egypt.

(Received March 2, 2006)

Toxicity of the fungicide Flutolanil was *in vitro* tested against 20 isolates of *Macrophomina phaseolina* and cotton seedlings of ten commercial cotton cultivars. The isolates were recovered from roots of cotton plants obtained from different cotton-growing areas in Egypt. Most of the tested isolates were sensitive to Flutolanil; however, they varied in sensitivity. Twenty-five percent of the isolates were highly sensitive where IC_{50} ranged from <1 to $5.1 \mu\text{g/ml}$, 20% of the isolates were sensitive where IC_{50} ranged from 15 to $30 \mu\text{g/ml}$, 45% of the isolates were moderately sensitive where IC_{50} ranged from 46 to $58.5 \mu\text{g/ml}$, and 10% of the isolates were not much sensitive (tolerant) where IC_{50} was $>100 \mu\text{g/ml}$. Flutolanil was very safe on both shoots and roots of the tested cultivars ($IC_{50} >100 \mu\text{g/ml}$). Treating cotton seeds with Flutolanil resulted in highly significant ($P < 0.01$) reductions in pathogenicity of 18 isolates and a significant reduction ($P < 0.05$) in pathogenicity of isolate M_2 . M_1 was the only isolate, which was insensitive to the application of Flutolanil. *In vivo* toxicity to Flutolanil was not correlated with its *in vitro* toxicity. However, a highly significant correlation ($r = 0.60$, $P < 0.01$) was observed between pathogenicity of isolates and the *in vivo* toxicity of the fungicide.

KEYWORDS: Cotton, Fungicide, *Macrophomina phaseolina*, Root rot

Macrophomina phaseolina (Tassi) Goid., the causal agent of charcoal rot of cotton, is of widespread distribution in the Egyptian soil, and it is easily and frequently isolated from cotton roots particularly during the late period of the growing seasons (Aly *et al.*, 2000).

M. phaseolina is a non-specialized fungus, attacking more than 500 host species (Sinclair, 1982); therefore, rotation of cotton with other crops is a questionable practice for controlling this pathogen.

Resistance to *M. phaseolina* has not been found in cotton (Hollida and Punithalingam, 1970). Sources of resistance were considered completely lacking in Pakistan (Akhtar, 1977). Aly *et al.* (1998) and Omar (2005) reported variation in susceptibility of Egyptian cottons to *M. phaseolina*; however, none of the tested cultivars showed satisfactory level of resistance.

Thus, fungicides have become indispensable for controlling *M. phaseolina* on cotton. For instance, Dwivedi and Ghaube (1985) studied effect of fungicides on the emergence and infection of cotton seedlings by *M. phaseolina* in pot experiments. They found that emergence was maximized after treatment with a TMTD soil drench and the least seedling infection occurred in soils treated with benlate, TMTD, and PCNB (Quintozene).

Chauhan (1986a) reported that control of seedling disease of cotton due to *M. phaseolina* was best and germination maximized when seeds were treated with carben-

dazim, followed in effectiveness by quintozone.

Chauhan (1986b) evaluated seed treatments of 7 fungicides and selected pair-wise combinations. The percentage of root rot of cotton at harvest ranged from 14 to 22% in treated plots compared with 28% in the control. Benzimidazole-derived fungicides differed in their effects on isolates of *M. phaseolina*. Benlate and carbendazim were most inhibitory but a sesame isolate was less sensitive to all fungicides.

Chauhan (1988) obtained good control of *M. phaseolina* on cotton by seed treatment with carbendazim followed by quintozone, while TMTD and catafol were less effective. A pre-sowing soil drench with quintozone and carbendazim gave very effective control.

Aly *et al.* (2001) evaluated the efficiency of monceren, Pencycuron, carboxin 200, Tolclofos-methyl, and maximum AP for controlling *M. phaseolina* on cotton, under greenhouse conditions. Monceren, Pencycuron, and Tolclofos-methyl were equally effective in increasing the percentage of surviving seedlings and the plant height; however, monceren was the only fungicide, which significantly increased dry weight of seedlings. Omar (2005) found that monceren and Tolclofos-methyl were the best performing fungicides in controlling *M. phaseolina* on cotton, under greenhouse conditions. This superiority was attributed to the following reasons: First, they were effective in increasing the percentage of surviving seedlings by 203.09 and 170.10%, respectively, regardless of the tested isolate. Second, they were also effective in in-

*Corresponding author <E-mail: Yehiam2001@yahoo.com>

creasing the height of surviving seedlings by 93.21 and 62.57%, respectively. Third, monceren and Tolclofos-methyl were effective in increasing dry weight of the surviving seedlings in 50 and 40% of the tested isolates, respectively.

The objective of this investigation was to evaluate variation in sensitivity to the fungicide Flutolanil among isolates of *M. phaseolina* isolated from cotton roots.

Materials and Methods

In vitro study. A laboratory study was carried out to evaluate sensitivity of *M. phaseolina* isolates to five concentrations of the fungicide flutolanil (Flutolanil, 25% WP). Stock solution of flutolanil (500 $\mu\text{g/ml}$) was prepared by dissolving an appropriate amount in sterile distilled water. Then the tested concentrations of 1, 15, 25, 50 and 100 $\mu\text{g/ml}$ were obtained by adding the appropriate amounts of stock suspension of the fungicide to autoclaved PDA medium cooled to about 45°C. Fungicide was not added to control treatments. Fungicide-amended and non amended PDA for each isolate was poured into 9 cm diameter plates. There were five replicates (plates) for each treatment. After solidification of PDA, each plate was centrally inoculated with 5 mm-diameter disc taken from margin of actively growing fungal colony on PDA. Plates were incubated at $30 \pm 3^\circ\text{C}$, and colony diameters were measured when the fungal growth completely covered plates of the control treatments. Percentage toxicity was calculated according to the following formula (Topps and Wain, 1957): percentage of inhibition of mycelial growth (I% toxicity) = $[(A - B)/A] \times 100$ where A = mean diameter of growth in the control treatment and B = mean diameter of growth of a given isolates grown on the pesticide amended PDA. Linear regression equations were fitted to logarithmic probability data of fungicide concentration and percentage of toxicity of Flutolanil-treated isolates. Slope values and IC_{50} values were interplotted (Finney, 1970). IC_{50} is the dose required to cause fungi toxic inhibition of the tested isolates by 50% in comparison with the control treatment.

Phytotoxicity test. The test was carried out according to El-Nawawy *et al.* (1972). Cotton (*Gossypium barbadense* L.) seeds of 10 commercially grown cultivars (Table 2) were dipped in water for 3 to 4 hrs. then incubate in wet cotton cloth for 24 hrs. The selected germinating seeds with their rootless were immersed slightly in the surface of agar in a test tube (1.5 \times 20 cm) containing IC_{50} of Flutolanil. Percentage inhibition of both roots and shoots were measured according to the following formula: $\text{I}\% = [(A - B)/A] \times 100$ where A = length of shoot or root in the control and B = length of shoot or root in a given treatment.

In vivo toxicity under greenhouse conditions. Substrate for growth of each isolate of *M. phaseolina* was prepared in 500-ml glass bottles; each bottle contained 50 g of sorghum grains and 40 ml of tap water. Contents of each bottle were autoclaved for 30 minutes. Isolate inoculum, taken from one-week-old culture on PDA, was aseptically introduced into the bottle and allowed to colonized sorghum for three weeks. The sorghum-fungus mixture of each isolate was used to infest autoclaved clay loam soil at a rate of 50 g/kg soil, and infested soil was dispensed in 10-cm-diameter clay pots. Flutolanil was added to slightly moist seeds (cultivar Giza 89) at a rate of 2 g/kg seeds. The seeds were shaken thoroughly in plastic bag for five minutes and allowed to dry before being planted in the infested pots of each isolate. Untreated seeds were planted in the infested pots of each isolate as a control. There were three replicates (pots) for each treatment. The pots were randomly distributed on a greenhouse bench under temperature regime ranged from $22 \pm 3^\circ\text{C}$ to $42.5 \pm 3.5^\circ\text{C}$. Percentages of dead seedlings were recorded 45 days after planting.

A randomized complete block design with three replicates was used in the greenhouse study. Percentage data were transformed into arc sine angles before carrying out analysis of variance (ANOVA) to produce approximately constant variance. Least significant difference (LSD) was applied for comparing between untreated and treated seeds of each isolate. ANOVA and correlation analysis were performed with a computerized program.

Results and Discussion

List of fungal strains of *M. phaseolina* and its geographic origin have shown in Table 1. It is quite clear that the isolates under the study nearly obtained from all over the country (Egypt). Most of the tested isolates were sensitive to Flutolanil; however, they varied in sensitivity (Table 2). In general, it was possible to classify these isolates into four distinct categories: highly sensitive, sensitive, moderately sensitive, and poorly sensitive (tolerant). The highly sensitive category included M_4 , M_5 , M_{13} , M_{18} , and M_{25} (25% of the isolates) where IC_{50} ranged from < 1 to 5.1 $\mu\text{g/ml}$. The sensitive category included M_{15} , M_{19} , M_{20} and M_{29} (20% of the isolates where IC_{50} ranged from 15 to 30 $\mu\text{g/ml}$).

The moderately sensitive category included M_1 , M_2 , M_3 , M_8 , M_{10} , M_{11} , M_{14} , M_{17} and M_{28} (45% of the isolates) where IC_{50} ranged from 46 to 58.5 $\mu\text{g/ml}$. The poorly sensitive (tolerant) category included M_7 and M_{12} (10% of the isolates) where IC_{50} was $> 100 \mu\text{g/ml}$. This high level of variability in sensitivity to Flutolanil among *M. phaseolina* isolates is in conformity with the results of Omar (2005) who found *in vivo* interaction between isolates of *M. phaseolina* and seed-dressing fungicides-that is, the

Table 1. List of fungal isolates of *Macrophomina phaseolina* used in this study

Isolate number	Geographic origin
M ₁	Minufiya-Sirs El-Laian
M ₂	Assiut El-Kosiya
M ₃	Giza-Giza
M ₄	Gharbiya-Tanta
M ₅	Minufiya-Shoubra Blol
M ₇	Gharbiya-Tanta
M ₈	Beheira-Damanhour
M ₁₀	Damietta-Kafr Saad
M ₁₁	Daqahliya-Minyet El-Nasr
M ₁₂	Minufiya-Berkit El-Saba
M ₁₃	Kafr El-Sheikh El-Riyad
M ₁₄	Assiut-Manfalout
M ₁₅	Giza-Giza
M ₁₇	Sharqiya-Zagazig
M ₁₈	Daqahliya-Aga
M ₁₉	Sharqiya-Diyarb Nigm
M ₂₀	Beheira-Kom Hamada
M ₂₅	El-Minya-Bani Mazar
M ₂₈	Sohag-Tahta
M ₂₉	Sohag-Tema

Table 2. IC₅₀ values and slopes of Flutolanil fungicide, dosage - linear growth response curves for *Macrophomina phaseolina* isolates in fungicide amended with PDA after 72 hrs of growth at 28°C

Isolate no.	IC ₅₀ (µg/ml)*	Slope
M ₁	58.5	0.511
M ₂	53	0.512
M ₃	52	0.320
M ₄	1	0.045
M ₅	5.1	0.229
M ₇	>100	0.305
M ₈	51	0.500
M ₁₀	50.5	0.440
M ₁₁	50	0.864
M ₁₂	>100	0.460
M ₁₃	<1	0.193
M ₁₄	46	0.290
M ₁₅	15	0.376
M ₁₇	50	0.299
M ₁₈	<1	0.120
M ₁₉	18	0.179
M ₂₀	30	0.374
M ₂₅	<1	0.144
M ₂₈	48	0.360
M ₂₉	17	0.241

*IC₅₀ (µg/ml) and slopes were determined by extrapolation from log - dosage probit.

isolates responded differently to the application of fungicides.

Treating cotton seeds with Flutolanil resulted in highly significant ($P < 0.01$) reductions in pathogenicity of 18

Table 3. Effect of the seed-dressing Flutolanil on pathogenicity of 20 isolates of *M. phaseolina* on cotton seedlings (cultivar Giza 89) under greenhouse condition

Isolate no.	Percentage of dead seedlings (Pathogenicity (%)) ^a		Reduction in pathogenicity ^b	<i>In vivo</i> toxicity ^c
	Untreated seed (T ₁)	Flutolanil-treated seed (T ₂)		
	M ₁	53.33		
M ₂	76.67	23.33	53.34**	69.58
M ₃	63.33	20.00	43.33**	68.42
M ₄	93.33	43.33	50.00**	53.57
M ₅	96.67	40.00	56.67**	58.62
M ₇	93.33	33.33	60.00*	64.29
M ₈	80.00	20.00	60.00**	75.00
M ₁₀	53.33	30.00	23.33**	43.75
M ₁₁	66.67	13.33	53.34**	80.01
M ₁₂	86.67	30.00	56.67**	65.39
M ₁₃	100.00	16.67	83.33**	83.33
M ₁₄	100.00	36.67	63.33**	63.33
M ₁₅	86.67	23.33	63.37**	73.08
M ₁₇	86.67	16.67	70.00**	80.77
M ₁₈	80.00	30.00	50.00**	62.50
M ₁₉	100.00	23.33	76.67**	76.67
M ₂₀	100.00	10.00	90.00**	90.00
M ₂₅	66.67	26.67	40.00**	60.00
M ₂₈	80.00	13.33	66.67**	83.34
M ₂₉	56.67	36.67	20.00*	35.29

^aPathogenicity is the percentage of dead seedlings after 45 days from planting date. Percentage data were transformed into arc sine angles before carrying ANOVA to produce approximately constant variance.

^bReduction in pathogenicity was non significant (N.S), significant at $P < 0.05$ (*), or highly significant $P < 0.01$ (**) according to LSD test. LSD was 17.11 ($P < 0.05$) or 22.98 ($P < 0.01$).

^cToxicity was calculated according to the following formula: $[(T_1 - T_2 / T_1) \times 100]$, where T₁ is the number of untreated seeds and T₂ is the number of Flutolanil-treated seeds.

isolates and a significant reduction ($P < 0.05$) in pathogenicity of M₂₉. *In vivo* toxicity of flutolanil against *M. phaseolina* isolates ranged from 31.24 to 90%. M₁ was the only isolate, which was insensitive to the application of Flutolanil (Table 3). Data of Table 4 showed that Flutolanil was very safe on both shoots and roots of the tested cultivars (IC₅₀ > 100 µg/ml). These results are almost in agreement with the results of (Ismail and Aly 1997; Zein, et al., 1999; Ismail, 2004).

Although *In vivo* toxicity of flutolanil against *M. phaseolina* isolates ranged from 31.24 to 90%; however, *in vivo* toxicity of Flutolanil was not correlated with its *in vitro* toxicity (Table 5). This lack of correlation may be attributed to the fact that *in vitro* efficiency of a fungicide is an outcome of a direct interaction between the chemical composition of the fungicide and the genotype of the pathogen. Under greenhouse conditions, other factors may interfere modifying the outcome of the *in vitro* interac-

Table 4. IC₅₀ and slopes values of the fungicide Flutolanil on root and shoot systems of cotton seedling on different cultivars

Cultivar	Root system		Shoot system	
	IC ₅₀ (µg/ml) ^a	Slope	IC ₅₀ (µg/ml)	Slope
Giza86	>100	0.77	>100	0.73
Giza88	>100	0.70	>100	0.66
Giza80	>100	0.68	>100	0.79
Giza70	>100	0.65	>100	0.72
Giza90	>100	0.80	>100	0.53
Giza89	>100	0.75	>100	0.59
Giza83	>100	0.63	>100	0.68
Giza45	>100	0.87	>100	0.69
Giza91	>100	0.55	>100	0.60
Giza85	>100	0.79	>100	0.52

^aThe concentration of Flutolanil (dose) that is required to cause phytotoxic inhibition of the cultivar by 50% in comparison with control treatment.

Table 5. Correlation coefficients among pathogenicity of *M. phaseolina* isolates, *in vitro* toxicity; and *in vivo* toxicity of Flutolanil against these isolates


Variable	1	2	3
1. Pathogenicity ^a	–	–0.145	0.596** ^d
2. <i>In vitro</i> toxicity ^b	–0.145	–	0.006
3. <i>In vivo</i> toxicity ^c	0.596** ^d	0.006	–

^aPercentage of dead seedlings when untreated seeds were planted in *M. phaseolina*-infested soil.

^b*In vitro* toxicity was expressed as IC₅₀ (µg/ml) under pure culture conditions.

^c*In vivo* toxicity is the magnitude of reduction in pathogenicity under greenhouse condition when Flutolanil - treated seeds were planted in the infested soil.

^dCorrelation coefficient is significant at $p < 0.01$ (**).

**Fig. 1.** Correlation coefficient between isolates pathogenicity and the *in vivo* toxicity of the fungicide.

tion. These factors may include (stability of the fungicide in soil (Snel *et al.*, 1970; Buchenauer, 1975; Huppertz *et al.*,

1984), activity of fungicide against ineffective propagules of the isolates (Weinhold and Bowman, 1974; Ktaria and Grover, 1975; Huppertz *et al.*, 1983), the host cultivar (Katoria and Verma, 1990), and the inoculum density (Gaber *et al.*, 1979). This result indicates that *in vitro* performance of Flutolanil is of no practical value in predicting its *in vivo* performance under greenhouse conditions.

A highly significant correlation was observed between pathogenicity of isolates and the *in vivo* toxicity of the fungicide (Table 5 and Fig. 1). This correlation indicates that Flutolanil had the highest inhibitory against the most pathogenic isolates of *M. phaseolina*, while it showed the lowest inhibitory effect against the least pathogenic isolates.

References

- Akhtar, C. M. 1977. Biological control of some plant diseases lacking genetic resistance of the host crops in Pakistan. *Annals of the New York Academy of Science* **287**: 45-56.
- Aly, A. A., El-Shazly; A. M. M., Youssef, R. M. and Omar, M. R. 2001. Chemical and biological control of charcoal rot of cotton caused by *Macrophomina phaseolina*. *J. Agric. Sci. Mansoura Univ.* **26**: 7661-7674.
- _____, Hussein; E. M., Allam, A. D. A., Amein, A. M. and El-Samawaty, A. M. A. 2000. Pathological studies on fungi involve in damping-off of cotton seedlings and root rot of adult plants in Upper Egypt Governorates. *J. Agric. Sci. Mansoura Univ.* **25**: 4015-4034.
- _____, Mostafa, M. A., Salwa, A., El-Rehim, Abd. and Omar, M. R. 1998. Susceptibility of some Egyptian cotton cultivars to *Macrophomina phaseolina* and its deteriorative effects on fiber quality. *Menofiya J. Agric. Res.* **23**: 1157-1167.
- Buchenauer, H. 1975. Difference sunlight stability of some carboxylic acid anilide fungicides in relation to their applicability for seed and foliar treatment. *Pestic. Sci.* **6**: 525-535.
- Chauhan, M. S. 1986a. Comparative efficacy of fungicides for the control of seedling diseases of cotton due to *Rhizoctonia* spp. *Ind. Mycol. Pl. Pathol.* **16**: 335-337.
- Chauhan, M. S. 1986b. Systemic and non-systemic fungicides against root rot of cotton in Haryana. *Ind. Mycol. Pl. Pathol.* **16**: 226-227.
- _____. 1988. Relative efficiency of different methods for the control of seedling disease of cotton by *Rhizoctonia bataticola*. *Ind. Mycol. Pl. Pathol.* **18L**: 25-30.
- Dwivedi, T. S. and Chaube H. S. 1985. Effect of fungicides on the emergence and infection of cotton seedlings by *Macrophomina phaseolina* (Tassi) Goid. *Ind. Mycol. Pl. Pathol.* **15**: 295-296.
- El-Nawawy, A. S., Kadouse, E. A. and Tag El-Din. 1972. Phytocidal properties of several N-aryl uranium and S-alkylisothiouranium arylarsonate. Proc.11th BL.Weed control comp. Pp 1226-1330.
- Finney, D. J. 1952. Probit analysis. Cambridge Univ. Press. p. 318.
- Garber, R. H., DeVay; J. E., Weinhold, A. R. and Matheron, D. 1979. Relationship of pathogen inoculum to cotton seedling disease control with fungicides. *Plant Dis. Repr.* **63**: 264-250.
- Holliday, P. and Pinithalingam, E. 1970. *Macrophomina phaseo-*

- lina*. CMI Description of Pathogenic Fungi and Bacteria. No. 275. Commonwealth Agricultural Bureaux, Farnham Royal, Bucks, UK.
- Huppatz, J. L., Phillips, J. N. and Witrzens, B. 1983. Laboratory and greenhouse studies on the activity of carboxamide derivatives against *Rhizoctonia solani* in cotton. *Plant Dis.* **67**: 45-47.
- _____, _____ and _____. 1984. Structure-activity relationship in a series of fungicidal pyrazole carboxanilides. *Agric. Biol. Chem.* **48**: 45-50.
- Ismail, A. A. and Aly, A. A. 1997. Sensitivity of some isolates of *Rhizoctonia solani* isolated from cotton seedlings to the insecticide Gauche in combination with seed-dressing fungicides used for controlling seedling disease. *J. Agric. Science, Mansoura Univ.* **22**: 753-755.
- Kataria, H. R. and Verma, P. R. 1990. Efficacy of fungicidal seed treatments against pre-emergence damping-off and post-emergence seedling root rot of growth-chamber grown canola caused by *Rhizoctonia solani* AG-2-1 and AG-4. *Can. J. Plant Pathol.* **12**: 409-416.
- Kataria, H. R. and Grover, R. K. 1975. Inhibition of infection structures of *Rhizoctonia solani* by fungi toxicants. *Z. Pflkrank. Pflschutz.* **82**: 226-232.
- Omar, M. R. 2005. Pathological and biochemical studies on *Macrophomina phaseolina* pathogenic on cotton. Ph.D. Thesis. Suez Canal Univ., Ismailia, p 182.
- Sinclair, J. B. 1982. Compendium of soybean diseases. The American Phytopathological Society St. Paul., Minnesota, p 104.
- Snel, M., Von Schmelting, B. and Edginton, L. V. 1970. Fungi toxicity and structure-activity relationships of some oxathin and thiazole derivatives. *Phytopathology* **60**: 1164-1169.
- Topps, J. H. and Wain, R. L. 1957. Investigation of fungicides. III. The fungitoxicity crop.sci. 12: 847-850 of 3 and 5 alkyl salicylanilide and para chloro anilines. *Ann. Appl. Biol.* **45**: 506-511.
- Weinhold, A. R. and Bowman, T. 1974. Repression of virulence in *Rhizoctonia solani* by glucose and 3-0-methyl glucose. *Phytopathology* **64**: 985-990.
- Zein, A. A., Shady, M. F. and Ismail, A. A. 1999. Phytotoxicity of certain pesticides, plant extracts and their mixtures on cotton soybean and wheat seedlings. *J. Agric. Science, Mansoura Uni.* **24**: 2009-2019.