

Identification of differentially expressed cDNAs in *Acanthamoeba culbertsoni* after mouse brain passage

Kyu-Lee HAN, Jongweon LEE, Don-Soo KIM, Soon-Jung PARK, Kyung-il IM and
Tai-Soon YONG*

Department of Parasitology and Institute of Tropical Medicine, Yonsei University College of Medicine, Seoul 120-752, Korea

Abstract: Free-living amoebae of the genus *Acanthamoeba* are causative agents of granulomatous amebic encephalitis and amebic keratitis. Because the virulence of *Acanthamoeba culbertsoni* cultured in the laboratory is restored by consecutive brain passages, we examined the genes induced in mouse brain-passaged *A. culbertsoni* by differential display reverse transcriptase polymerase chain reaction (DDRT-PCR). Enhanced *A. culbertsoni* virulence was observed during the second mouse brain passage, i.e., infected mouse mortality increased from 5% to 70%. Ten cDNAs induced during mouse brain passage were identified by DDRT-PCR and this was confirmed by northern blot analysis. BlastX searches of these cDNAs indicated the upregulations of genes encoding predictive NADH-dehydrogenase, proteasomal ATPase, and GDP-mannose pyrophosphorylase B, which have previously been reported to be associated with *A. culbertsoni* virulence factors.

Key words: *Acanthamoeba culbertsoni*, free-living ameba, mouse, DDRT-PCR, virulence

INTRODUCTION

Free-living amoebae of the genus *Acanthamoeba* occur in various environments, and some cause granulomatous amebic encephalitis and amebic keratitis in humans (Gardner et al., 1991; Tan et al., 1993; Cohen et al., 1987). Moreover, amebic virulence is attenuated by the long-term cultivation of amebas in vitro and can be recovered by serial mouse brain passages via intranasal infection (Lee et al., 1983; Marzur et al., 1994). Peroxidase and proteinase activities have also been reported to increase in *Acanthamoeba* in parallel with the recovery of virulence by mouse brain passage (Marzur et al., 1994). On recovering virulence, protease activities and some secretory products were

found to be increased in pathogenic *Acanthamoeba* (Hadas et al., 1993). We speculated that virulence restoration also occurs in parallel with the upregulation of virulence-related gene expressions, and that the gene products of these genes may play roles in the pathogenesis of *Acanthamoeba*. Accordingly, using DDRT-PCR, we identified upregulations in the transcription of some *A. culbertsoni* genes during mouse brain passages.

MATERIALS AND METHODS

Cultivation of amebas and mouse brain passages

Acanthamoeba culbertsoni (ATCC No. 30171) was axenically cultured at 37°C in Casitone-Glucose-Vitamin (CGV) medium (Willaert, 1971). *A. culbertsoni* trophozoites (1×10^6) were inoculated intranasally

• Received 29 August 2005, accepted after revision 20 January 2006.

*Corresponding author (e-mail: tsyong212@yumc.yonsei.ac.kr)

into 8-week-old male ICR mice under anesthesia. Infected ICR mouse mortality was observed over 25 days post-infection. Amebas recovered from infected mice were also cultured in CGV medium for 1 week and were used to re-infect mice. In order to recover amebas, autopsies were performed immediately after sacrificing animals and portions of brain tissues were placed in CGV medium.

Preparation of *A. culbertsoni* DNA and RNA

The genomic DNA of *A. culbertsoni* was prepared by lysing 1×10^7 trophozoites in 1% SDS, and following this with phenol/chloroform extraction and ethanol precipitation. Total RNA was isolated from *A. culbertsoni* (in vitro cultivated, and from animals recovered from the 1st, 2nd and 3rd passages) using TRIZOL reagent (Gibco/BRL, Gaithersburg, Maryland, U.S.A.), according to the manufacturer's instructions.

DDRT-PCR

DDRT-PCR was performed as described previously (Liang et al., 1992) using a No.1 RNA image kit (GenHunter, Nashville, TN). DNA-free RNAs were obtained by treating total RNA with RNase-free DNase I. Total RNA from long-term in vitro cultured and mouse brain passaged amebas were converted to cDNA using 1 of the 3 oligo-dTs provided in the kit (T₁₁A, T₁₁G, or T₁₁C). Reactions were initiated at 65°C for 5 min and 37°C for 10 min, 100 units of Superscript II reverse transcriptase (Gibco/BRL) was then added, and reaction mixes were incubated at 37°C for 50 min and then at 75°C for 5 min. Using a 20 ng portion of cDNA as a template, a pool of cDNA was generated by PCR using 1 of the arbitrary primers and an appropriate oligo-dT. Parameters for the PCR reaction were as follows; pre-incubation for 2 min at 94°C, 40 amplification cycles (30 sec at 94°C, 2 min at 40°C, 30 sec at 72°C), and a final extension for 10 min at 72°C, cDNA bands were visualized by including a trace amount of ³²P (1 µCi/20 µl reaction) in the PCR reaction. The cDNAs generated were then separated on 6% SDS-urea polyacrylamide gels. Differentially displayed cDNAs were re-amplified using the same PCR conditions, and the re-amplified cDNAs were cloned into

pGEM-T easy vector (Promega, Madison, Wisconsin, U.S.A.) for sequencing and Northern blot analysis.

Northern blot analysis

cDNA probes were labeled with ³²P using a Random labeling kit (Takara, Japan), and used to examine expression patterns after 3 cycles of infection and in long-term in vitro cultured amebas. For Northern hybridization, samples of total RNA (10 µg) were fractionated in 1% formaldehyde agarose gels, blotted on Hybond-N membranes by capillary transfer in 20× SSPE (3 M NaCl, 0.2 M NaH₂PO₄, 0.02 M EDTA) and immobilized on membranes using a UV-crosslinker. Blots were prehybridized in 5× SSPE, 50% formamide, 5× Denhardt's solution, 0.5% SDS and 100 µg/ml salmon sperm DNA for 2 hr at 42°C, and hybridization was continued overnight at 42°C in the presence of a ³²P labeled probe. Membranes were washed twice in 2× SSPE/0.1% SDS at room temperature for 15 min, and then twice at 60°C for 30 min in 0.2× SSPE/0.5% SDS. Blots were hybridized with a labeled beta-actin DNA fragment, as an internal RNA control. mRNAs levels were measured using a Phosphorimager BAS 2500 (Fuji-Film Co. Ltd., Japan) and analyzed using TINA 2.0 software.

Sequence analysis

cDNA clones showing an increased hybridization pattern after mouse passages versus in vitro cultivated amebas were sequenced using the dideoxy nucleotide chain termination method. The reactions were run and analyzed on a capillary automated sequencing machine (ABI 377 XL90; Applied Biosystems). For data analysis, the nucleotide sequences of the selected cDNA clones were compared with entries in the EMBL and GenBank databases, and all 6 reading frames of the selected cDNA clones were also searched for proteins and proteins in other organisms homologous to the determined *A. culbertsoni* sequences using BlastX.

RESULTS

Mouse-passaged amebas showed increased viru-




Fig. 1. Survival rate of mice intranasally infected with *A. culbertsoni*; from 5% of mortality in the first infection to 70% in the third infection observed at 20 days of infection.




Fig. 2. Differential display using A, G or C-base anchored oligo-dT primers. PCR was carried out using single-strand cDNA produced from the total RNA from non-mouse brain passaged (0) and mouse brain passaged (1st, 2nd and 3rd) *A. culbertsoni*. Lane zero (non-mouse brain passage), 1, 2 and 3 (1st, 2nd and 3rd mouse brain passage) were compared by differential display using dT11A (or G, C) primer in combination with 8 arbitrary primers. Arrows indicate bands showing increased intensity after mouse brain passage.

lence; from 5% of mortality in the first infection to 70% in the third infection observed at 25 days of infection (Fig. 1). DDRT-PCR was performed upon the mRNAs of brain-passaged amebas and upon the RNA of long-term in vitro cultured amebas using 24 combinations of 8 arbitrary primers and 3 oligo-dT₁₁ primers. Fifteen amebic cDNAs were found to show upregulat-




Fig. 3. Northern blot analysis of *A. culbertsoni* RNA obtained before- mouse brain passage (0) or after mouse brain passage (1st, 2nd, and 3rd mouse brain passages). The same membrane was stripped and re-hybridized with a non-regulated probe encoding *A. culbertsoni* actin as an RNA loading control. Signals were quantified using a Bio-Imaging Analyser System 2500 (Fuji-Film).

ed expression patterns following mouse brain passages, and were identified as shown in Fig. 2. To confirm the elevated expressions of these selected cDNA clones in *A. culbertsoni* we performed Northern blot hybridization. Ten of the 15 isolated cDNAs showed increased expressions in mouse brain-passaged amebas, ranging from 1.5 to 25-fold in 4G-1 and 3A-1 (Fig. 3). DNA sequencing of 10 clones with a higher hybridization pattern in mouse brain-passaged amebas showed that 4 of these clones (3A-1, 3A-2, 3A-3 and 3G) had almost the same sequence. BlastX searches for 5 (3A-1, 3A-2, 3A-3, 3G and 4G-1) of these 10 clones failed to find any protein sequences in public databases. The sequences of the remaining 5 amplicons provided useful information with respect to their putative functions (Table 1). One clone 2A showed homology to genes encoding p47 tumor suppressor protein of human and mouse. Interestingly their mRNAs rapidly increased at the first mouse brain passage and decreased marginally after the second and third passages. 7C showed homology to a mitochondrial metabolic enzyme – NADH-dehydrogenase

Table 1. Identification of clones selected by DDRT-PCR

Clone ^{a)}	cDNA size	Northern blot analysis ^{b)}	Putative identification	E-value	Accession number ^{c)}
8G	138 bp (partial)	3	GDP-mannose pyrophosphorylase B	0.41	AY161267
7G-1	447 bp (partial)	3	Proteasomal ATPase	4e-47	AY161265
7G-2	252 bp (partial)	2	ATP(GTP)-binding fet5 mRNA	2e-07	AY161266
7C	357 bp (partial)	10	NADH-dehydrogenase (<i>Acanthamoeba castellanii</i>)	1e-37	AY161264
2A	231 bp (partial)	2	P47 tumor suppressor protein	2e-09	AY161262
4G-1	229 bp (partial)	1.5	Novel gene		AY161268
3G	195 bp (partial)	25	Novel gene		AY161263
3A-1	191 bp (partial)	25	Novel gene		AY161263
3A-2	185 bp (partial)	10	Novel gene		AY161263
3A-3	191 bp (partial)	25	Novel gene		AY161263

^{a)}Arabic numbers represent arbitrary primers. Alphabet (A, G or C) represents the one-nucleotide base anchored to oligo-dT primers.

^{b)}Numbers indicate highest fold of the hybridization intensity on northern blot analysis among 1st, 2nd, or 3rd mouse brain passage compared with non-passaged amebas.

^{c)}Nucleotide sequences submitted to EMBL and GenBank™ databases by this study.

subunit I. 7G-1 was found to be homologous to the proteasomal ATPase gene. 7G-2 matched ATP/GTP-binding fet5 mRNA, which is a known eukaryotic transcription factor of unknown function. 8G showed homology to GDP-mannose pyrophosphorylase B, a known virulence factor.

DISCUSSION

It has been postulated that the expressions of *A. culbertsoni* genes that are directly or indirectly responsible for virulence, may be upregulated by consecutive brain passages, since the virulence of long-term cultured *A. culbertsoni* in the laboratory is recovered by consecutive brain passages. DDRT-PCR has been applied to analyze tissues and developmental stages by comparing reverse transcribed cDNA from purified mRNA. For example, cancer cell-specific mRNA, human brain tissue-specific mRNA (Sokolov et al., 1994), and developmental stage-specific mRNA in *Eimeria bovis* of the coccidian parasite (Abrahamsen et al., 1995) and in *Fasciola hepatica* (Reed et al., 1998) have been successfully identified using this technique. In the present study, DDRT-PCR produced 3 kinds of cDNA pools from mRNA templates when different oligo-dT₁₁-A, G or C primers were used during the

reverse transcription reaction. Each cDNA pool was analyzed by PCR using the same oligo-dT₁₁-A, G, or C primer used for cDNA synthesis, and an arbitrary primer.

In this study, we examined differentially expressed genes in *A. culbertsoni* after mouse brain passages and compared these with genes expressed after long-term in vitro culture, by DDRT-PCR. Ten differentially expressed clones in mouse brain-passaged amebas were identified by DDRT-PCR, and confirmed by Northern blot analysis. Based on sequencing analysis results, clone 2A showed homology to the human p47/p33 tumor suppressor protein. And, p47 tumor suppressor protein was reported to play an important role in cellular stress response to ultraviolet irradiation, during DNA repair and apoptosis (Chen et al., 2003). Other amplicons were matched with NADH-dehydrogenase, proteasomal ATPase, ATP/GTP-binding fet5 mRNA, and GDP-mannose pyrophosphorylase B. The *Salmonella* serotype Gallinarum containing mutated NADH dehydrogenase was reported to be less invasive than wild *Salmonella* serotype Gallinarum, and showed no evidence of multiplication in the liver or spleen (Li et al., 1998). In particular, proteasomal ATPase is known to function in the main degradation pathway of most proteins in the cyto-

plasm and nuclei of eukaryotic cells (Hershko et al., 1998; DeMartino et al., 1994). The proteasomal ATPase subunit was reported to recognize the polyubiquitin degradation signal (Lam et al., 2002; Aaron, 1994), and to be required for nitric oxide resistance (Darwin et al., 2003). These findings suggest that the proteasomal degradation pathway of cellular protein and the nitric oxide protection mechanism are turned on during *A. culbertsoni* infection in mice. In addition, ATP/GTP-binding fet5 mRNA is regulated by iron and upregulated in cells grown in iron-limited media, which suggests that Fet5 protein normally plays a role in the transport of iron (Spizzo et al, 1997), an important factor in virulence change. In fungi, GDP-mannose pyrophosphorylase B gene deletion was found to be lethal (Varki, 1999), and *Leishmania mexicana* containing a mutated form of this gene, lost its ability to infect murine macrophages (Attila et al., 2001). Moreover, GDP-mannose pyrophosphorylase B was suggested to be a virulence factor for other pathogens, and to play a significant role in host invasion (Attila et al., 2001). Although it is difficult to conclude that any direct relationship exists between these induced genes and increased *A. culbertsoni* virulence caused by mouse brain passage, proteasomal ATPase and GDP-mannose pyrophosphorylase B may be expected to participate in the virulence of amebic infection in the mouse brain.

The information gathered from the present study provides resources for further investigations into the molecular mechanisms of *Acanthamoeba* virulence. The availability of the cDNA sequence coding for *Acanthamoeba* should provide the opportunity to design experiments for recombinant protein and immunization of selected genes.

REFERENCES

- Aaron C (1994) The ubiquitin-proteasome proteolytic pathway. *Cell* **79**: 13-21.
- Abrahamsen MS, Johnson RR, Hathway M, White MW (1995) Identification of *Eimeria bovis* merozoite cDNAs using differential mRNA display. *Mol Biochem Parasitol* **71**: 183-191.
- Arentsen JJ, Laibson PR (1985) Diagnosis and management of *Acanthamoeba keratitis*. *Am J Ophthalmol* **100**: 389-395.
- Attila G, Thomas L (2001) Disruption of mannose activation in *Leishmania mexicana* GDP-mannose pyrophosphorylase is required for virulence, but not for viability. *EMBO J* **20**: 3657-3666.
- Chen B, Campos EI, Crawford R, Martinka M, Li G (2003) Analysis of the tumour suppressor ING1 expression and gene mutation in human basal cell carcinoma. *Int J Oncol* **22**: 927-931.
- Cohen EJ, Parlato CJ, Arentsen JJ, Genvert GI, Eagle RC Jr Wieland MR, Laibson PR (1987) Medical and surgical treatment of *Acanthamoeba*. *J Protozool* **32**: 604-607.
- Darwin KH, Ehrst S, Gutierrez-Ramos JC, Weich N, Nathan CF (2003) The proteasome of *Mycobacterium tuberculosis* is required for resistance to nitric oxide. *Science* **302**: 1963-1966.
- DeMartino GN, Moomaro CR, Zagnitko OP, Proske RJ, Chu-Ping M, Afendis SJ, Swatfield JC, Slaughter CA (1994) PA700, an ATP-dependent activator of the 20S proteasome, and ATPase is containing multiple members of nucleotide-binding protein family. *J Biol Chem* **269**: 20878-20884.
- Gardner HAR, Martnez AJ, Visvesvara GS, Sotrel A (1991) Granulomatous amoebic encephalitis in an AIDS patient. *Neurology* **41**: 1993-1995.
- Hadas E, Mazur T (1993) Proteolytic enzymes of pathogenic and non-pathogenic strains of *Acanthamoeba* spp. *Trop Med Parasitol* **44**: 197-200.
- Hershko A, Iechanover A (1998) The ubiquitin system. *Ann Rev Biochem* **67**: 425-479.
- Lam YA, Lawson TG, Velayutham M, Zweier JL, Pickart CM (2002) A proteasomal ATPase subunit recognizes the polyubiquitin degradation signal. *Nature* **18**: 763-767.
- Lee DK, Lee KT, Im KI (1983) Changes in the pathogenicity of *Naegleria fowleri* by several brain passaged in mice. *Korean J Parasitol* **21**: 234-240.
- Li ZB, Turner AK, Dougan G, Barrow PA (1998) Protection of chickens against experimental fowl typhoid using a nuoG mutant of *Salmonella* serotype Gallinarum. *Vaccine* **16**: 899-903.
- Liang P, Averboukh L, Keyomarsi K, Sager R, Pardee AB (1992) Differential display and cloning of mRNAs from human breast cancer versus mammary epithelial cells. *Cancer Res* **52**: 6966-6968.
- Marzur T, Hadas E (1994) The effect of the passages of *Acanthamoeba* strains through mice tissues on their virulence and its biochemical markers. *Parasitol Res* **80**: 431-

434.

- Reed MB, Spothill TW, Strugnell RA, Panaccio M (1998) *Fasciola hepatica*; Stage-specific expression of novel gene sequences as identified by differential display. *Exp Parasitol* **89**: 169-179.
- Sokolov BP, Prockop DJ (1994) A rapid and simple PCR-based method for isolation of cDNA from differentially expressed genes. *Nucleic Acids Res* **22**: 4009-4015.
- Spizzo T, Byersdorfer C, Dueterhoeft S, Eide D (1997) The yeast FET5 gene encodes a FET3-related multicopper oxidase implicated in iron transport. *Mol Gen Genet* **256**: 547-556.
- Tan B, Weldon-Linne M, Rhone DP, Penning CL, Visvesvara GS (1993) *Acanthamoeba* infection presenting as skin lesions in patients with the acquired immunodeficiency syndrome. *Archpathol Lab Med* **117**: 1043-1046.
- Varki A (1999) Exploring the biological roles of glycans, *Essentials of Glycobiology* p57-68, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY.
- Willaert E (1971) Isolation and *in vitro* culture of the amoeba of the genus *Naegleria*. *Ann Soc Belg Med Trop* **51**: 701-708.