

The Anti-allergic Effect of Vinegar Treated Small Black Soybean (*Glycine max* Merr.) is Mediated Through Inhibition of Histamine Release and Inflammatory Cytokine Secretion

Tae-Yong Shin*, Hye-Young Shin¹, Jong-Pil Lim, Hoon Jeon, Dae Keun Kim, Byeong-Suk Chae, Young-Ee Kwon, Sin-Myoung Kang, Ik-Jun Chang, Jeong-Suk Park², Jae-Hyeok Lee³

College of Pharmacy, Woosuk University, Jeonbuk 565-701, Korea

¹Department of Infection Biology, Zoonosis Research Center, School of Medicine, Wonkwang University, Jeonbuk 570-749, Korea

²Department of Oriental Alternative Medicine, Nambu University, Gwangju 506-706, Korea

³Department of Oriental Pharmaceutical Development, Nambu University, Gwangju, 506-706, Korea

Abstract – In this report, we investigated the effect of aqueous extract of vinegar treated small black soybean (*Glycine max* Merr.) (Leguminosae) (VSBS) on mast cell-mediated allergic reaction and pro-inflammatory cytokine secretion. VSBS inhibited compound 48/80-induced systemic reactions. VSBS attenuated immunoglobulin (Ig) E-mediated passive cutaneous anaphylaxis. In addition, VSBS decreased the phorbol 12-myristate 13-acetate plus calcium ionophore A23187-stimulated secretion of tumor necrosis factor (TNF)- α , interleukin (IL)-6 and interleukin (IL)-8 in human mast cells. Our findings provide evidence that VSBS inhibits mast cell-derived allergic reactions.

Keywords – *Glycine max* Merr., allergic reaction, mast cells, tumor necrosis factor (TNF)- α , interleukin (IL)-6, interleukin (IL)-8

Introduction

Mast cells are important producers of mediators of inflammatory responses such as allergy and immediate-type allergic reaction. Immediate-type allergic reaction is mediated by histamine released in response to the antigen cross-linking of immunoglobulin E (IgE) bound to Fc ϵ RI on mast cells. After activation via the Fc ϵ RI, the mast cells start the process of degranulation which results in the releasing of mediators, such as products of arachidonic acid metabolism and an array of inflammatory cytokines (Metacalf, *et al.*, 1981; Church, *et al.*, 1997; Miyajima, *et al.*, 1997). Among the inflammatory substances released from mast cells, histamine remains the best-characterized and most potent vasoactive mediator implicated in the acute phase of immediate hypersensitivity (Cook, *et al.*, 2002). Mast cell degranulation also can be elicited by the synthetic compound 48/80, and it has been used as a direct and convenient reagent to study the mechanism of anaphylaxis (Ennis, *et al.*, 1980). Activated mast cells can

produce histamine, as well as a wide variety of other inflammatory mediators such as eicosanoids, proteoglycans, proteases, and several pro-inflammatory and chemotactic cytokines such as TNF- α , IL-6, IL-4, IL-8, IL-13, and transforming growth factor- β (Burd, *et al.*, 1989; Plaut, *et al.*, 1989; Galli, *et al.*, 1991; Bradding, *et al.*, 1993). Therefore, modulation of secretion of these cytokines from mast cells can provide a useful therapeutic strategy for allergic inflammatory disease. Small black soybean (*Glycine max* Merr.) (Leguminosae) has been used as a health food in Korea, and it also has medicinal uses due to its physiological functions. It has been reported that small black soybean has physiological activities, including antioxidant activity (Sa, *et al.*, 2003, hypoglycemic effect (Lee, *et al.*, 2004) and anti-allergic effect (Shin, *et al.*, 2007). In this study, we evaluated the effect of vinegar treated small black soybean (VSBS) on systemic and local allergic reaction. In addition, the effect of VSBS on phorbol 12-myristate 13-acetate (PMA) plus calcium ionophore A23187 (A23187)-induced TNF- α , IL-6 and IL-8 secretion in a human mast cell line (HMC-1) was also investigated.

*Author for correspondence

Fax: +82-63-290-1812; E-mail: tyshin@woosuk.ac.kr

Experimental

Culture of HMC-1 Cells – HMC-1 cells, a human mast cell line were grown in Iscove's media supplemented with 10% FCS and 2 mM glutamine.

Animals – The original stock of male ICR mice were purchased from Dae-Han Biolink Co. Ltd. (Daejeon, Korea). The animals were maintained in the College of Pharmacy, Woosuk University. The animals were housed five per cage in a laminar air flow room maintained at a temperature of 22 ± 2 °C, with a relative humidity of $55 \pm 5\%$ throughout the study. The care and treatment of the mice were in accordance with the guidelines established by the Public Health Service Policy on the Humane Care and Use of Laboratory Animals and were approved by the Institutional Animal Care and Use Committee.

Reagents – Compound 48/80, anti-DNP IgE, DNP-human serum albumin (HSA), phorbol 12-myristate 13-acetate, calcium ionophore A23187 were purchased from Sigma Chemical Co. (St Louis, MO). rTNF- α , rIL-6 and rIL-8 were purchased from R & D Systems Inc. (Minneapolis, MN).

Preparation of VSBS – Small black soybean of *G max* (the 2003 product) were purchased at Imsil, Jeonbuk, Korea. The soybeans were kept in un-milled rice vinegar in two weeks, and air-dried. A voucher specimen was deposited at the Herbarium of the College of Pharmacy, Woosuk University. The plant sample was extracted with purified water at 70 °C for 5 h. The extract was filtered, and lyophilized. The dried extract was dissolved in saline or Tyrode buffer A (10 mM HEPES, 130 mM NaCl, 5 mM KCl, 1.4 mM CaCl₂, 1 mM MgCl₂, 5.6 mM glucose, 0.1% bovine serum albumin) before use.

Compound 48/80-induced systemic anaphylaxis – Mice were given an intraperitoneal injection of 0.008 g/kg body weight (BW) of the mast cell degranulator, compound 48/80. VSBS was dissolved in saline and administered intraperitoneally ranging from 0.05 to 1 g/kg BW 1 h before the injection of compound 48/80 (n = 10/group). In the time dependent experiment, VSBS (1 g/kg) was administered intraperitoneally at 5, 10, 15, and 20 min after injection of compound 48/80 (n = 10/group). Mortality was monitored for 1 h after induction of anaphylactic shock.

Passive cutaneous anaphylaxis (PCA) – The mice were injected intradermally with 0.5 mg of anti-DNP IgE into each of 2 dorsal skin sites that had been shaved 48 h earlier. The sites were outlined with a water-insoluble red marker. After 48 h, each mouse was received an injection of 1 mg of DNP-HSA in PBS containing 4% Evans blue (1 : 4) via the tail vein. VSBS (0.001 to 1 g/kg BW) was

intraperitoneally administered 1 h before the challenge. Thirty minutes after the challenge, the mice were killed and the dorsal skin was removed for measurement of the pigment area. The amount of dye was determined colorimetrically after extraction with 1 mL of 1 M KOH and 9 mL of a mixture of acetone and phosphoric acid (5 : 13) based on the previous report (Katayama, *et al.*, 1978). The absorbent intensity of the extraction was measured at 620 nm by using a spectrophotometer, and the amount of dye was calculated with Evans blue measuring-line.

Assay of TNF- α , IL-6 and IL-8 secretion – TNF- α , IL-6 and IL-8 secretion was measured by modification of an enzyme-linked immunosorbent assay (ELISA) as described previously (Scuderi, *et al.*, 1986). HMC-1 cells were cultured with α -MEM plus 10% FBS and resuspended in Tyrode buffer A. The cells were sensitized with PMA (20 nM) plus A23187 (1 μ M) for 8 h in the absence or presence of VSBS. The ELISA was performed by coating 96-well plates with 6.25 ng/well of monoclonal antibody with specificity for TNF- α , IL-6 and IL-8 respectively. Before use and between subsequent steps in the assay, the coated plates were washed twice with PBS containing 0.05% Tween-20 and twice with PBS alone. For the standard curve, rTNF- α , rIL-6 and rIL-8 were added to serum previously determined to be negative to endogenous TNF- α , IL-6 and IL-8. After exposure to the medium, the assay plates were exposed sequentially to biotinylated anti-human TNF- α , IL-6 and IL-8, 2,2'-azino-bis (3-ethylbenzthiazoline-6-sulfonic acid) (ABTS) tablets substrates. Optical density readings were made within 10 min of the addition of the substrate with a 405 nm filter.

Statistical analysis – Statistical analyses were performed using SAS statistical software (SAS Institute, Cary, NC). Treatment effects were analyzed using one-way ANOVA, followed by Duncan's multiple range tests. $P < 0.05$ was used to indicate significance.

Results

Effect of VSBS on compound 48/80-induced systemic reaction – To determine the effect of VSBS on allergic reaction, an *in vivo* model of a systemic reaction was used. Compound 48/80 (0.008 g/kg) was used as a model of induction of systemic fatal allergic reaction. After the intraperitoneal injection of compound 48/80, the mice were monitored for 1 h, after which the mortality rate was determined. Compound 48/80 induced fatal shock in 100% of animals. When VSBS was administered

Table 1. Effect of VSBS on compound 48/80-induced systemic reaction




VSBS treatment (g/kg BW)	Compound 48/80 (0.008 g/kg BW)	Mortality (%)
none (saline)	+	100
0.05	+	100
0.1	+	70
0.5	+	50
1	+	20
1	-	0

Groups of mice ($n = 10/\text{group}$) were intraperitoneally pretreated with 200 μl of saline or VSBS at various doses 1 h before the intraperitoneal injection of compound 48/80. Mortality (%) within 1 h following compound 48/80 injection is represented as the number of dead mice $\times 100/\text{total number of experimental mice}$.

Table 2. Time-dependent effect of VSBS on compound 48/80-induced systemic reaction

VSBS treatment (g/kg, BW)	Time (min)	Compound 48/80 (0.008 g/kg BW)	Mortality (%)
none (saline)	0	+	100
1	5	+	30
	10	+	60
	15	+	60
	20	+	100

Groups of mice ($n = 10/\text{group}$) were intraperitoneally pretreated with 200 μl of saline or VSBS. VSBS (1 g/kg) was given at 5, 10, 15, and 20 min after the intraperitoneal injection of compound 48/80. Mortality (%) within 1 h following compound 48/80 injection is represented as the number of dead mice $\times 100/\text{total number of experimental mice}$.

**Fig. 1.** Effect of VSBS on the PCA reaction. VSBS was intraperitoneally administered 1 h prior to the challenge with antigen. Each date represents the mean \pm SEM of three independent experiments. * $p < 0.05$; significantly different from the saline value.**Fig. 2.** Effect of VSBS on the TNF- α secretion. PMA plus A23187-stimulated HMC-1 cells were incubated for 8 h in the absence or presence of VSBS. TNF- α secreted into the medium are presented as the mean \pm SEM of three independent experiments. * $p < 0.05$; significantly different from the PMA + A23187 value.**Fig. 3.** Effect of VSBS on the IL-6 secretion. PMA plus A23187-stimulated HMC-1 cells were incubated for 8 h in the absence or presence of VSBS. IL-6 secreted into the medium are presented as the mean \pm SEM of three independent experiments. * $p < 0.05$; significantly different from the PMA + A23187 value.

intraperitoneally at a concentrations ranging from 0.05 to 1 g/kg BW for 1 h, the mortality with compound 48/80 was dose-dependently reduced (Table 1). In addition, the mortality of mice administered with VSBS (1 g/kg) 5, 10, 15, and 20 min after compound 48/80 injection increased time-dependently (Table 2).

Effect of VSBS on IgE-mediated PCA reaction – To confirm the anti-allergic effects of VSBS in *in vivo* model, we used PCA model. A local extravasation was induced by a local injection of anti-DNP IgE followed by an intravenous antigenic challenge. VSBS was




Fig. 4. Effect of VSBS on the IL-8 secretion. PMA plus A23187-stimulated HMC-1 cells were incubated for 8 h in the absence or presence of VSBS. IL-8 secreted into the medium are presented as the mean \pm SEM of three independent experiments. * $p < 0.05$; significantly different from the PMA + A23187 value.

administered intraperitoneally 1 h prior to the challenge with antigen. The administration of VSBS showed a dose-dependent inhibition in the PCA reaction (Fig. 1).

Effect of VSBS on the secretion of pro-inflammatory cytokines from HMC-1 cells – We examined whether VSBS could regulate pro-inflammatory cytokines such as TNF- α , IL-6 and IL-8 in HMC-1 cells. TNF- α , IL-6 and IL-8 are important inflammatory cytokines released from mast cells. The HMC-1 cell line is a useful cell for studying the cytokine activation pathway (Kim, *et al.*, 2006). Stimulation of HMC-1 cells with PMA plus A23187 induced the secretion of TNF- α , IL-6 and IL-8. However, pretreatment with VSBS decreased PMA plus A23187-induced TNF- α , IL-6 and IL-8 secretion (Fig. 2, Fig. 3, Fig. 4).

Discussion

The results of this study demonstrated that VSBS has anti-allergic properties. VSBS inhibited compound 48/80-induced systemic allergic reaction and anti-DNP IgE-mediated local allergic reaction. These results indicate that mast cell-mediated immediate-type allergic reactions are inhibited by VSBS. Numerous reports established that stimulation of mast cells with compound 48/80 or IgE initiates the activation of signal-transduction pathway, which leads to histamine release. Several recent studies have shown that compound 48/80 and other polybasic compounds are able, apparently directly, to activate G-proteins (Mousli, *et al.*, 1990a; Mousli, *et al.*, 1990b;

Chahdi, *et al.*, 2000). Compound 48/80 increases the permeability of the lipid bilayer membrane by causing a perturbation in the membrane. This result indicates that the increase in membrane permeability may be an essential trigger for the release of the mediator from mast cells. In this sense, anti-allergic agents having a membrane-stabilizing action may be desirable. VSBS might stabilize the lipid bilayer membrane, thus preventing the perturbation being induced by compound 48/80. Additionally, the VSBS administered mice are protected from anti-DNP IgE-mediated PCA, one of the most important *in vivo* models of anaphylaxis in local allergic reaction. This finding suggests that VSBS might be useful in the treatment of allergic skin reactions. Pro-inflammatory cytokines, including TNF- α , IL-6 and IL-8, play a major role in triggering and sustaining the allergic inflammatory response in mast cells (Walsh, *et al.*, 1991; Mullarkey, *et al.*, 1994). Mast cells are a principal source of TNF- α in human dermis. TNF- α has an important amplifying effect in asthmatic inflammation and potently stimulates airway epithelial cells to produce cytokines (Galli, *et al.*, 1991). It is also a potent inducer of other inflammatory cytokines, including IL-1, IL-6, IL-8 and GM-CSF. IL-6 is produced from mast cells and its local accumulation is associated with PCA reaction (Mican, *et al.*, 1992). IL-8 from mast cells, acts on surrounding cells such as neutrophils, T lymphocytes, and eosinophils, and plays a role in activation of inflammatory effector cells (Sengupta, *et al.*, 2004). These reports indicate that the reduction of pro-inflammatory cytokines from mast cells is one of the key indicators of reduced allergic symptom. In our present study, VSBS inhibited the secretion of TNF- α , IL-6 and IL-8 in PMA plus A23187-stimulated HMC-1 cells. This result suggests that the anti-allergic effect of VSBS results from its reduction of TNF- α , IL-6 and IL-8 release from mast cells.

The results obtained in the present study shows that VSBS contributes to the prevention or treatment of mast cell-mediated allergic diseases.

Acknowledgements

“This work was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD)” (The Regional Universities Program/Center for Healthcare Technology Development)

References

Bradding, P., Feather, I.H., Wilson, S., Bardin, P.G., Heusser, C.H.,

- Holgate, S.T., and Howarth, P.H., Immunolocalization of cytokines in the nasal mucosa of normal and perennial rhinitic subjects. The mast cell as a source of IL-4, IL-5, and IL-6 in human allergic mucosal inflammation. *J. Immunol.* **151**, 3853-3865 (1993).
- Burd, P.R., Rogers, H.W., Gordon, J.R., Martin, C.A., Jayaraman, S., Wilson, S.D., Dvorak, A.M., Galli, S.J., and Dorf, M.E., Interleukin 3-dependent and -independent mast cells stimulated with IgE and antigen express multiple cytokines. *J. Exp. Med.* **170**, 245-257 (1989).
- Chahdi, A., Fraundorfer, P.F., and Beaven, M.A., Compound 48/80 activates mast cell phospholipase D via heterotrimeric GTP-binding proteins. *J. Pharmacol. Exp. Ther.* **292**, 122-130 (2000).
- Church, M.K., and Levi-Schaffer, F., The human mast cell. *J. Allergy Clin. Immunol.* **99**, 155-160 (1997).
- Cook, E.B., Stahl, J.L., Barney, N.P., and Graziano, F.M., Mechanisms of antihistamines and mast cell stabilizers in ocular allergic inflammation. *Curr. Drug Targets Inflamm. Allergy* **1**, 167-180 (2002).
- Ennis, M., Pearce, F.L., and Weston, P.M., Some studies on the release of histamine from mast cells stimulated with polylysine. *Br. J. Pharmacol.* **70**, 329-334 (1980).
- Galli, S.J., Gordon, J.R., and Wershil, B.K., Cytokine production by mast cells and basophils. *Curr. Opin. Immunol.* **3**, 865-872 (1991).
- Katayama, S., Shionoya, H., and Ohtake, S., A new method for extraction of extravasated dye in the skin and the influence of fasting stress on passive cutaneous anaphylaxis in guinea pigs and rats. *Microbiol. Immunol.* **22**, 89-101 (1978).
- Kim, S.H., Jun, C.D., Suk, K., Choi, B.J., Lim, H., Park, S., Lee, S.H., Shin, H.Y., Kim, D.K., and Shin, T.Y., Gallic acid inhibits histamine release and pro-inflammatory cytokine production in mast cells. *Toxicol. Sci.* **91**, 123-131 (2006).
- Lee, D.H., Kwak, D.H., Kim, S.M., Ju, E.J., Choi, H.G., Kim, O.H., Hwang, J.B., Bae, N.G., Jung, K.Y., Han, J.C., Park, H.D., and Choo, Y. K., Effect of small black soybean powder on blood glucose and insulin sensitivity in streptozotocin-induced diabetic rats. *J. Korean Soc. Food Sci. Nutr.* **33**, 1618-1625 (2004).
- Metcalfe, D.D., Kaliner, M., and Donlon, M.A., The mast cell. *Crit. Rev. Immunol.* **3**, 23-27 (1981).
- Mican, J.A., Arora, N., Burd, P.R., and Metcalfe, D.D., Passive cutaneous anaphylaxis in mouse skin is associated with local accumulation of interleukin-6 mRNA and immunoreactive interleukin-6 protein. *J. Allergy Clin. Immunol.* **90**, 815-824 (1992).
- Miyajima, I., Dombrowicz, D., Martin, T.R., Ravetch, J.V., Kinet, J.P., and Galli, S.J., Systemic anaphylaxis in the mouse can be mediated largely through IgG1 and Fc gammaRIII. Assessment of the cardiopulmonary changes, mast cell degranulation, and death associated with active or IgE- or IgG1-dependent passive anaphylaxis. *J. Clin. Invest.* **99**, 901-914 (1997).
- Mousli, M.C., Bronner, C., Bockaert, J., Rouot, B., and Landry, Y., Interaction of substance P, compound 48/80 and mastoparan with a-subunit C-terminal of G protein. *Immunol. Lett.* **25**, 355-358 (1990a).
- Mousli, M.C., Bronner, C., Landry, Y., Bockaert, J., and Rouot, B., Direct activation of GTP-binding regulatory proteins (G proteins) by substance P and compound 48/80. *FEBS Lett.* **259**, 260-262 (1990b).
- Mullarkey, M.F., Leifeman, K.M., Peters, M.S., Caro, I., Roux, E.R., Hanna, P.K., Rubin, A.S., and Jacobs, C.A., Human cutaneous allergic late-phase response is inhibited by soluble IL-1 receptor. *J. Immunol.* **152**, 2033-2041 (1994).
- Plaut, M., Pierce, J.H., Watson, C.J., Hanley-Hyde, J., Nordan, R.P., and Paul, W.E., Mast cell lines produce lymphokines in response to cross-linkage of Fc epsilon RI or to calcium ionophores. *Nature* **339**, 64-67 (1989).
- Sa, J.H., Shin, I.C., Jeong, K.J., Shim, T.H., Oh, H.S., Kim, Y.J., Cheung, E.H., Kim, G.G., and Choi, D.S., Antioxidative activity and chemical characteristics from different organs of small black soybean (Yak-Kong) grown in the area of Jungsun. *Korean J. Food Sci. Technol.* **35**, 309-315 (2003).
- Scuderi, P., Sterling, K.E., Lam, K.S., Finley, P.R., Ryan, K.J., Ray, C.G., Petersen, E., Slymen, D.J., and Salmon, S.E., Raised serum levels of tumor necrosis factor in parasitic infections. *Lancet* **13**, 1364-1365 (1986).
- Sengupta, B., Banerjee, A., and Sengupta, P.K., Investigations on the binding and antioxidant properties of the plant flavonoid fisetin in model biomembranes. *FEBS Lett.* **570**, 77-81 (2004).
- Shin, T.Y., Shin, H.Y., Lim, J.P., Kim, D.K., Chae, B.S., Kwon, Y.E., Oh, C.H., Cho, M.G., Lee, T.K., Park, J.S., Lee, J.H., and Jeon H., Small black soybean (*Glycine max* Merr.) inhibits mast cell-mediated allergic reaction and inflammatory cytokine secretion. *Nat. Prod. Sci.* **13**, 263-267 (2007).
- Walsh, L.J., Trinchieri, G., Waldorf, H.A., Whitaker, D., and Murphy, G.F., Human dermal mast cells contain and release tumor necrosis factor alpha, which induces endothelial leukocyte adhesion molecule 1. *Proc. Natl. Acad. Sci. USA* **88**, 4220-4224 (1991).

(Accepted November 12, 2007)