

Cloning and Expression Analysis of a Chitinase Gene *Crchi1* from the Mycoparasitic Fungus *Clonostachys rosea* (syn. *Gliocladium roseum*)

Zhongwei Gan[§], Jinkui Yang[§], Nan Tao, Zefen Yu, and Ke-Qin Zhang*

Laboratory for Conservation and Utilization of Bio-resources, and Key Laboratory for Microbial Resources of the Ministry of Education, Yunnan University, Kunming 650091, P. R. China

(Received August 13, 2007 / Accepted October 16, 2007)

Clonostachys rosea (syn. *Gliocladium roseum*) is a well-known biocontrol agent and widely distributed around the world. In this study, an endochitinase gene *Crchi1* was isolated from the mycoparasitic fungus *C. rosea* using the DNA walking strategy. The *Crchi1* ORF is 1,746 bp long and interrupted by three introns. The cloned gene *Crchi1* encodes 426 amino acid residues and shares a high degree of similarity with other chitinases from entomopathogenic and mycoparasitic fungi. Several putative binding sites for transcriptional regulation of *Crchi1* in response to carbon (5'-SYGGRG-3') and nitrogen (5'-GATA-3') were identified in the upstream of *Crchi1*. Expression of *Crchi1* gene in different carbon sources was analyzed using real-time PCR (RT-PCR). We found that the *Crchi1* expression was suppressed by glucose but strongly stimulated by chitin or solubilized components of the cell wall from *Rhizoctonia solani*. Phylogenetic analysis of chitinases from entomopathogenic and mycoparasitic fungi suggests that these chitinases have probably evolved from a common ancestor.

Keywords: carbon starvation, chitinase, *Clonostachys rosea*, expression, phylogenetic analysis, real-time PCR

The filamentous fungus *Clonostachys rosea* (syn. *Gliocladium roseum*: teleomorph *Bionectria ochroleuca*) is a common saprotroph with a worldwide distribution. It is associated with organisms such as fungi, nematodes and plants and is versatile in terms of its ecological niches, frequently found in cultivated soils, in wastes containing phytotoxic phenolic acids, and in aerial plant organs (Sutton *et al.*, 1997; Krauss *et al.*, 2001). Many isolates of *C. rosea* are highly efficient antagonists against several plant pathogenic fungi, as shown in the controls of *Botrytis cinerea* in strawberry, raspberry, and tomato (Sutton *et al.*, 1997), *Rhizoctonia solani* on tobacco (Lahoz *et al.*, 2002), and as an entomopathogenic fungus of insects *Oncometopia tucumana* and *Sonesimia grossa* (Toledo *et al.*, 2006). The extraordinary ecological versatility renders *C. rosea* a good candidate for use in agriculture.

As a mycoparasite, *C. rosea* has been tested successfully as a biological control agent against divergent fungal plant pathogens (Sutton *et al.*, 1997; Xue, 2003). The mycoparasitic activity has been attributed to the secretion of its cell-wall degrading enzymes, including chitinases. Chitinase (EC3.2.1.14) is a type of enzyme capable of degrading chitin. A wide variety of organisms can produce and secrete chitinase. Recent genetic evidence has suggested that a 42 kDa extracellular chitinase present in several *Trichoderma* species plays a critical role in their antagonistic activities against *B. cinerea* and *R. solani* (Lorito *et al.*, 1998; Baek *et al.*, 1999; Kim *et al.*, 2002). In addition, there is evidence

that in *Trichoderma* spp. chitinase *ech42* was inducible by fungal cell walls and by colloidal chitin (Carsolio *et al.*, 1994; Garcia *et al.*, 1994) as well as by carbon starvation (Margolles-Clark *et al.*, 1996). In addition, *ech42* expression appeared to be governed by carbon catabolite repression (Carsolio *et al.*, 1994; Lorito *et al.*, 1996).

In our previous reports, two serine proteases were identified from *C. rosea*, and both were found to play a role in infection against nematodes (Zhao *et al.*, 2005; Li *et al.*, 2006). However, little is known about the encoding genes of chitinase from *C. rosea* and the regulation of its expression. In this study, we described the cloning of an endochitinase gene from *C. rosea* (*Crchi1*) and the regulation of its expression using real-time PCR (RT-PCR) analysis. The phylogenetic analysis of chitinases from different organisms was also discussed.

Materials and Methods

Strains, plasmids, and growth conditions

The isolate of mycoparasitic fungus *C. rosea* used in this study was originally isolated from a field soil sample in Yunnan Province and had been deposited in China General Microbiological Culture Collection Center (CGMCC 0806). The fungal strain was grown on Potato Dextrose Agar (PDA) medium at 28°C. *R. solani* was incubated in Potato Dextrose Broth and the cell walls were prepared as described previously (Schirmböck *et al.*, 1994). *Escherichia coli* DH5 α , used as the recipient strain for recombinant plasmids, and was grown in Luria-Bertani medium. The vector pMD18-T (Takara, Japan) was used for PCR cloning.

§ These authors contributed equally to this work.

* To whom correspondence should be addressed.

(Tel) 86-871-503-4878; (Fax) 86-871-503-4838

(E-mail) kqzhang111@yahoo.com.cn

Primer design

A pair of degenerate primers (Table 1, primer set 1) was synthesized according to a previous report (Bogo *et al.*, 1998) and used to amplify the conserved sequence of the chitinase gene from *C. rosea*. Two additional pairs of nested PCR primers (Table 1, primer set 2 and 3) were designed according to the amplified sequence to amplify the whole gene. A pair of specific primers containing restriction sites (*Eco*RI and *Bam*HI, respectively) (Table 1, primer set 4) were designed to amplify the cDNA sequence and for cloning. For *Crchi1* expression studies, the *Crchi1* primer (Table 1, primer set 5) was synthesized according to the *Crchi1* gene, and the beta-tubulin gene primers (*Tubb*) (Table 1, primer set 6) were synthesized based on the *C. rosea* beta-tubulin gene (AF358169) for endogenous control.

Primer dimer formation and the integrity of amplification reactions were confirmed by agarose gel electrophoresis (1.5% agarose) and melting curve analysis. In order to calculate the PCR efficiencies from *Crchi1* and *tubb* genes, a standard dilution curve was performed in duplicates using serial dilutions of standard cDNA. The efficiencies were calculated using a previously established method (Kenneth and Thomas, 2001).

Amplification of a chitinase gene from strain *C. rosea* CGMCC 0806

C. rosea was cultured on PDA medium for 7 days at 28°C. Mycelium for DNA extraction was collected from the plates and ground to a fine powder in liquid nitrogen. DNA was extracted using the CTAB method described previously (Zhang *et al.*, 1996).

A pair of degenerate primers (Table 1, primer set 1) was used to amplify the conserved sequence of chitinase genes from *C. rosea*. The genomic DNA was used as templates, and PCR conditions followed those described in a previous report (Yang *et al.*, 2005). To obtain the complete chitinase gene from *C. rosea*, two pairs of nested PCR primers were designed targeting the conserved sequence. The 5' and 3' terminal fragments of the chitinase gene were amplified using the DNA Walking Speedup™ Premix Kit (Seegene, Korea),

separately. PCR amplification was performed according to previous report (Gan *et al.*, 2007). All PCR products were purified and sequenced using an ABI PRISM 3730 automated sequencer (Perkin-Elmer, USA) with four fluorescent dyes.

Sequence analysis

Analysis of the DNA sequence was done using the DNAMAN software package (Version 5.2.2, Lynnon BioSoft, Canada). Homology search was performed using the BLAST algorithms against various databases in the GenBank (<http://www.ncbi.nlm.nih.gov/BLAST>). Potential N-glycosylation sites were analyzed by NetNGlyc 1.0 Server (<http://www.cbs.dtu.dk/services/NetNGlyc/>). Molecular weight, theoretical pI and amino acid composition were analyzed by ProtParam tool (<http://us.expasy.org/tools/protparam.html>).

Gene expression analysis

For *Crchi1* expression studies, *C. rosea* was first precultured for 3 days on a rich medium (Potato Dextrose Broth). The mycelium was collected by filtration, washed three times with sterilized water and transferred to liquid synthetic medium (SM) as described previously (Robert *et al.*, 1999). The carbon source in the synthetic medium was glucose, colloidal chitin, cell wall components of *R. solani* or periostracum cicadae. Mycelium was harvested after 48 h of incubation for RNA extraction.

Total RNA was isolated using the RNAPrep Plant Kit (DP402 TianGen, China) according to the manufacturer's protocol. Integrity of RNA was confirmed by running samples in a 1% agarose gel. RT-PCR amplification was performed using the RNA PCR Kit (AMV) Ver. 2.1 (TaKaRa, Japan) according to the manufacturer's protocol.

Expression levels of *Crchi1* were quantified with a real-time RT-PCR assay performed in an ABI Prism 7000 sequence detection system (Applied Biosystems, USA), using the SYBR Green Jump Start Taq Ready Mix (Sigma, USA). Each 20 µl reaction mixture consisted of 10 µl SYBR Green Supermix, 1 µl of each primer at 10 µmol, 6 µl nuclease-free water and 2 µl diluted cDNA. Synthesis of cDNA and

Table 1. List of oligonucleotide primers used in this study

	Primers	Sequence (5'-3')
1	Forward	GC(TC) GT(TC) TA(TC) TT(TC) AC(TC) AA(TC) TGG
	Reverse	GG(AG) TA(TC) TCC CA(AG) AT(AG) TC(CGAT) AT
2	Forward TSP1	TGC CCC TGA AAA ATG GAT GA
	Forward TSP3	CAT GAA TGG TCA CGA TGG ACT AAC
3	Reverse TSP1	CAT TGG TGG CTG GAC TTG GT
	Reverse TSP3	GCT GTT GAG TTC ATG AAG GAT TGG
4	Former	CCGGAATTGATG AAG TCG TTT TTC ACC CG
	Reserve	CGCGGATCCTCA GGA GAG GCT GTT CTT G
5	Crchi1F	CTC AAC TCC ATT CAG CAC CAA G
	Crchi1R	TGT AGT CCC AGA TTC CGC TCT C
6	TubbF	CAT CTT CAG ACC GGT CAG TG
	TubbR	AAG TAG ACG TTC ATG CGC TC

tcatgcc**atggcgg**tgccattcatccaaatccgatctgggtccatgcgtgctc -473
actccatgcttcttcatctgatgtctgggtcatgcagccacatgccc**cccc**caagtaac -420
 atcaacttgcgggaatgtcgg**tggttgg**ctgggttcagggcaacatgacatcatgctcga -360
 tcttgactcttgcgggttgtggaaggttgc**agaggg**cagca**ttggag**atgaaacctaa -300
 gagact**atggcgc****cccc****ttc**gataaatagtatct**gtggc**gtatctcatcctc**TATATA**at -240
 catatcaagttctccctctcagctcctgggttcttttctctcctcagcagcagagc -180
 tctctcctcccaaccattccccagcaacttccaectccttggacatttgtctaacctagcc -120
 tgggtgctggtaacttgcattcaagttctctctctctgagtttatcaactgagagccggttctc -60
 ATGAAGTCGTTTTTCA**CCCGT**CGCTGGCTCTTCTGGCCCTTGGCCAGCTACATTTGG**A** 60
M K S F F T R S L A L L A L A Q S T F G 20
 CGTGC**A**ACTCCTCGCATGGAGGACTTGGCTTCCACTGACCTTTCCACGGCTGCAACTGGA 120
R A T P R M E D L A S T D L S T R A T G 40
 TCCATCAATGCTGTCTACTTTACCAACTGgttagtccatcgtgaccatctcagggctcat 180
S I N A V Y F T N W 50
 tgtacagcttccataataatcatccatttttccagGGGCATCTATGGACGCAACTTCCAGCC 240
G I Y G R N F Q P 59
 TGCGGATCTTCAAGCTTCAAAAATTTTCGCATGTCTCTATTCTTTCATGAACCTCCGAGC 300
A D L Q A S K I S H V L Y S F M N L R A 79
 AGATGGCACGGTgttaagttaccttatgctcttcaagcctttgtgacccagttctggctaat 360
D G T V 83
 attgaac**ag**TTACTCGGGTGACACCTACGCGGACTTGGAGAAACATTACTCTGACGACT 420
Y S G D T Y A D L E K H Y S D D 99
gtaagcatcagcttcccttcttaaac**ag**ctctccggcactttgctaacgggtgacgggat 480

agCGTGGAAATGATATCGGAACAAATGCCTATGGCTGTGTC**AAGCAGCT**CTAC**AAGCTCAA** 540
S W N D I G T N A Y G C V K Q L Y K L K 119
 GAAGGCRAACCGCTCGCTCAAGATCATGTGTCCATTTGGTGGCTGGACTTGGTCGACCAA 600
K A N R S L K I M L S I G G W T W S T N 139
 CTTTCCCGCTGCTGCCTCCACCGAGGCTACCCGCTGTACATTTGCCAAGACCGCTGTGTA 660
F P A A A S T E A T F A K T A V E 159
 GTTCATGAAGGATTGGGGCTTTGACGGCATTGACGTCGACTGGGAGTACCCCGCCAGTGA 720
F M K D W G F D G I D V D W E Y P A S E 179
 GACTGACGCCAATAACATGGTCTCCTTCTTCAGCGAGTTCCGCAGGAGCTCGACTCGTA 780
T D A N N M V L L L Q R V R Q E L D S Y 199
 CTCCGCAACATATGCTAATGGCTATCACTTCCAACCTCTCCATTGCCGCTCCCGCAGGACC 840
S A T Y A N G Y H F Q L S I A A P A G P 219
 TAGCCATTACAACGTTCTTAAAGCTAGCCAGCTCGGTTCCGCTCTCGACAACATCAACCT 900
S H Y N V L K L A Q L G S V L D N I N L 239
 CATGGC**A**TACGACTATGCAGGTTCTGGGACAGTGTACGGCCATCAAGCC**A**CCCTGTA 960
M A Y D Y A G S W D S V S G H Q A N L Y 259
 TCCTAGCACATCAAAACCCAGCTCAACTCCATTCAAGCACC**AAGGCT**GGCGTGCAGGCATA 1020
P S T S N P S S T P F S T K A A V D A Y 279
 CATCGCAGCCGGCGTCCCTGC**AAGCAAGATC**ATTCTGGGTATGCCCATCTACGGCAGAGC 1080
I A A G V P A S K I I L G M P I Y G R A 299
 TTTTGTGGGAACCGACGGACCAGGCAAGCCCTACTCCACTATCGGCGAAGGCAGCTGGGA 1140
F V G T D G P G K P Y S T I G E G S W E 319
 GAGCGGAATCTGGGACTACAAAGTACTTCCC**AAGCCGGCGCT**TACCGTGATTACCGACTC 1200
S G I W D Y K V L P K A G A T V I T D S 339
 CGCGCCGGTGTACTTACAGCTACGACTCCAGCAGCAGGACCATGATCTCATACGATAC 1260
A A G A T Y S Y D S S S R T M I S Y D T 359
 CCCCATATGGTCCGCACAAAGGTCTCATATGCTAAGGGCCCTCGGACTCGGAGGCAGCAT 1320
P D M V R T K V S Y A K G L G L G S M 379
 GTTCGGGAGGCATCGGCCGACAAGACTGGCTCTGACTCGCTTATCGGC**A**CTGCCCTCAG 1380
F W E A S A D K T G S D S L I G T A L S 399
 CAGC**A**TGGGTAGCCTTGATAGCACCCAGAAC**TGCTCAGCT**ACCCCAACTCCAAGTTCGA 1440
S M G S L D S T Q N C L S Y P N S K F D 419
 CAACATCAAGAACAGCCTCTCTGAgcgggacccccgacccctcgtctggcatacttctg 1498
N I K N S L S * 427

Fig. 1. Nucleotide sequence of the *Crchl* from *C. rosea* (GenBank accession no. EU000575). The deduced amino acid sequence is shown in one-letter code under the DNA sequence. Nucleotide coordinates are shown to the right of the sequence. Introns are shown in lower-case letters, and their conserved 5'GT, 3'AG are in bold. General transcription control signals including TATAA regions upstream of the gene are in bold and underlined. Putative transcription control sites within the promoter region for carbon regulation (5'-SYGGRG-3') are in bold italics and underlined, sites for nitrogen regulation (5'-GATA-3') are in bold italics. The possible signal sequence cleavage point is represented by vertical arrows between the A₂₂ and T₂₃ residues. Potential N-glycosylation sites are by NRS and NPS residues and shaded in black. The white boxes (amino acids 130-133 and 166-174) corresponds to the conserved motif from family 18 of the glycosyl hydrolases.




Fig. 2. Alignment of the amino acid sequences of chitinases from different fungi. Areas shaded in black are conserved regions (100% similarity), and unshaded areas are regions of variability between the chitinases. Signal peptide sequences are marked on arrow, *** indicated the conserved substrate binding sites (SXGG), and ▲ indicated the catalytic domains (DXXDXDXE) of glycosyl hydrolase family 18.

RT-PCR was performed in triplicates for each gene. The PCR conditions followed the manufacturer's instructions. RT-PCR reactions for the endogenous control beta-tubulin gene (*Tubb*) and the target gene *Crchi1* were conducted in separate tubes. The expression level of *Crchi1* for 48 h on 1% glucose was used as the calibrator and defined as

one. The expression level calculated by the formula $2^{-\Delta\Delta CT}$ represents the x-fold difference from the calibrator.

Determination of fungal biomass

Fungal biomass was determined using a method described previously (Robert *et al.*, 1999). Values were in milligrams

of extractable protein per milliliter of culture.

Phylogenetic analysis

Amino acid sequences of chitinases from different organisms were aligned with CLUSTAL W (Thompson *et al.*, 1994) and a phylogenetic tree was constructed using PAUP version 4.0b10 (Swofford, 2002). Confidence values for individual branches were assessed from 1000 bootstrap replicates of the original sequence data.

Results

Sequence analysis of the chitinase gene *Crchi1* from *C. rosea* CGMCC 0806

A 579 bp PCR product was amplified by using degenerate primers set 1 (Table 1). The 5' and 3' flanking sequences were amplified using the DNA Walking Speedup™ Premix Kit, separately. The complete chitinase gene was obtained from the assembled DNA sequence with the DNAMAN software package. The sequence was verified as a chitinase gene by homologous analysis with other fungal chitinase genes. The complete nucleotide sequence of *Crchi1* has been submitted to GenBank (accession number EU000575).

The encoding gene of *CrChi1* consisted of one open reading frame (ORF) and three short introns (64, 58, and 62 bp in length, respectively). The splice junctions were determined by comparing the sequence of *Crchi1* with that of its corresponding cDNA sequence. Each intron began with GT and ended with AG, a common feature of fungal introns (Gurr *et al.*, 1987). The elements commonly found within fungal promoters were identified (Fig. 1) and include a TATAA box located at 188 bp upstream of the translation start codon ATG. In addition, putative binding sites for transcriptional regulation of *Crchi1* in response to carbon (binding site 5'-SYGGRG-3') (Sophianopoulou *et al.*, 1993) and nitrogen (binding site 5'-GATA-3') (Kudla *et al.*, 1990) are presented in Fig. 1.

The chitinase gene *Crchi1* encodes a polypeptide of 426 amino acid residues. Homologous analysis indicates that

CrChi1 is synthesized as a preproenzyme, and the first 22 amino acids have many characteristics of a signal peptide and the signal peptidase cleavage site is between A₂₂ and T₂₃. After processing of the 22 N-terminal amino acids, the calculated size of *CrChi1* is 43.8 kDa. Two potential N-glycosylation sites (NRS and NPS) were found by computational prediction (Fig. 1). Comparison of the *CrChi1* amino acid sequence with other chitinases in the GenBank database revealed that they shared a high degree of similarity among fungal chitinases, and the regions of SIGGW and FDGIDVDWE were highly conserved among those chitinases of the glycosyl hydrolase family 18 (Fig. 2). The active site centered on the catalytic residues Asp₁₆₇, Asp₁₇₀, Asp₁₇₂ and Glu₁₇₄ (in *CrChi1*) are conserved in all aligned sequences. Previous studies have indicated that substitution at any of these sites (D or E) could result in greatly reduced activity or a total elimination of activity (Seur *et al.*, 2007).

Moreover, the chitinase gene *Crchi1* shared a high degree of similarity with another endochitinase gene *Crech42* from *C. rosea*. The percent sequence identity was 85.12% at the nucleotide sequence ORF level (GenBank accession no. DQ523687) and 81.69% at the deduced amino acid sequence level (GenBank accession no. ABG02418). *Crchi1* contained three short introns, while *Crech42* contained only two introns.

Gene expression analysis

Amplification using the *Crchi1* primer (Table 1, primer set 5) yielded a 179 bp product, and the *tubb* primers (Table 1, primer set 6) yielded a 137 bp product. When analyzing the melting curve, a single product-specific melting curve was obtained for primers of *Crchi1* and *tubb*, respectively, indicating that primers were designed with optimal efficiency and were effective at targeting and amplifying only the genes of interest (data not shown). A plot of the log cDNA dilution versus ΔC_T ($C_{T, Crchi1} - C_{T, Tubb}$) showed that the absolute value of the slope is close to 0.10, suggesting that the efficiencies of the target gene and the reference gene were similar. We therefore used the $\Delta\Delta C_T$ calculation for the relative quantifications of the target gene.




Fig. 3. RT-PCR analyses of the expression of *Crchi1* in *C. rosea*. Strains were cultivated on SM with different C sources [colloidal chitin (Cch), glucose (G), and *R. solani* cell walls (CWRs or R)] for 48 h. The numbers 0.1, 0.5, 1 and 2 in X axis represent the different concentrations [weight in volume (w/v)] of carbon sources. A housekeeping gene, beta-tubulin (*Tubb*) was used as the endogenous control to help quantify PCR products. The expression level of *Crchi1* grown for 48 h on 1% glucose was used as the calibrator and defined as one. The expression level calculated by the formula $2^{-\Delta\Delta C_T}$. Y axis represents the x-fold difference from the calibrator. Error bars indicate SD.

Quantitative analysis of expression levels of *Crchi1* was achieved by a RT-PCR assay (Fig. 3). Our results showed that both genes (*Crchi1* and *tubb*) were found to be expressed constitutively. The highest level of *Crchi1* expression was observed with 0.5% CWRs (cell walls of *R. solani*) (Fig. 3A), followed by 0.5% CCh (colloidal chitin) (Fig. 2A and B), and 1% G (glucose) (Fig. 3A and C). The induction of *Crchi1* by the different concentrations of glucose and colloidal chitin were also tested (Fig. 3B and C). Glucose induced the highest level of expression at the concentration of 0.1% and the lowest level of expression at 2% (Fig. 3C). Similarly, CCh induced the highest level of expression at the concen-




Fig. 4. Biomass formation (total extractable protein) of *C. rosea* during cultivation on different concentration of glucose (□) and colloidal chitin (△). Values are means of at least three separate experiments carried out with the spores (6×10^6 /ml), error bars indicate standard deviation.

tration of 0.1% and the lowest level of expression at 1% (Fig. 3B). *Crchi1* mRNA showed a relative low level expression on 1% glucose in the presence of the 0.5% CCh (Fig. 3C).

Furthermore, with respect to *C. rosea* growth in SM with different concentrations of glucose and colloidal chitin, fungal biomass was determined (Fig. 4). The biomass results contrasted those of *Crchi1* expression. Specifically, biomass formation was lowest grown on 0.1% glucose and colloidal chitin. These results suggested that carbon starvation induced *Crchi1* expression in *C. rosea*, consistent with previous reports (Margolles-Clark *et al.*, 1996; Robert *et al.*, 1999).

Phylogenetic analysis

The phylogenetic tree of chitinases (Fig. 5) from various organisms was constructed using PAUP version 4.0b10 (Swofford, 2002). One chitinase belonging to glycosyl hydrolase family 19, identified from *Zea diploperennis* (Tiffin, 2004) was used as the outgroup. Other chitinases all belonged to glycosyl hydrolase family 18. In family 18, chitinases from *Neurospora crassa*, *Botryotinia fuckeliana*, *Aspergillus fumigatus* and *Coniothyrium mimitans* were clustered and formed a clade (clade II), while other chitinases from entomopathogenic and mycoparasitic fungi were clustered and formed a monophyletic clade (clade I). The monophyletic lineages provided a support for the hypothesis that these chitinases likely evolved from one common ancestor. Moreover, clade I consisted of three subclades: chitinases from mycoparasitic fungi (*Hypocrea* spp., *Trichoderma* spp.) were clustered and formed the subclade Ia, chitinases from different entomopathogenic fungi (e.g. *Beauveria bassiana*, *Paecilomyces fumosoroseus*, *Metarhizium anisopliae*) were clustered and formed the subclade Ib, and chitinase CrChi1 and CreCh42 from *C. rosea* formed the subclade Ic, and there is a statistical support (54%) for the significant divergence of CrChi1 from other chitinases.

Interestingly, the topology of the phylogenetic tree co-




Fig. 5. Phylogenetic tree showing the relationship between CrChi1 and other chitinases. The tree was constructed using the PAUP version 4.0b10 (Swofford 2002) program packages. The chitinase gene of *Z. diploperennis* (AAT40053) was used as an outgroup to root the tree. Confidence values were assessed from 1,000 bootstrap replicates of the original sequence data.

Table 2. Chitinases from different organisms used in our phylogenetic analysis

Taxonomy	Name	Accession no.
Eukaryota		
Fungi		
Ascomycota		
Eurotiomycetes		
Eurotiales		
Trichocomaceae		
<i>Aspergillus</i>	<i>Aspergillus fumigatus</i>	AAP23218
Leotiomycetes		
Helotiales		
Sclerotiniaceae		
<i>Botryotinia</i>	<i>Botryotinia fuckeliana</i>	AAM94807
Dothideomycetes		
Pleosporales		
Leptosphaeriaceae		
<i>Coniothyrium</i>	<i>Coniothyrium minitans</i>	AAG00504
Sordariomycetes		
Sordariales		
Sordariaceae		
<i>Neurospora</i>	<i>Neurospora crassa</i>	XP_957924
mitosporic Ascomycota		
<i>Aphanocladium</i>	<i>Aphanocladium album</i>	P32470
Sordariomycetes		
Hypocreales		
Bionectriaceae		
<i>Bionectria</i>	<i>Clonostachys rosea</i> (1)	ABG02418
	<i>Clonostachys rosea</i> (2)	EU000575
Clavicipitaceae		
<i>Lecanicillium</i>	<i>Lecanicillium psalliotae</i>	EF203917
<i>Beauveria</i>	<i>Beauveria bassiana</i>	AAN41260
<i>Paecilomyces</i>	<i>Paecilomyces fumosoroseus</i>	AAX19146
	<i>Paecilomyces farinosus</i>	ABD64606
<i>Torrubiella</i>	<i>Torrubiella confragosa</i>	AAV98692
<i>Nomuraea</i>	<i>Nomuraea rileyi</i>	AAP04616
<i>Metarhizium</i>	<i>Metarhizium anisopliae</i>	AAY32603
	<i>Metarhizium flavoviride</i>	CAB44709
mitosporic Hypocreales		
<i>Verticillium</i>	<i>Verticillium fungicola</i>	AAP45631
<i>Trichoderma</i>	<i>Trichoderma hamatum</i>	AAC60385
	<i>Trichoderma longibrachiatum</i>	ABD42921
	<i>Trichoderma asperellum</i>	AAF19623
	<i>Trichoderma aureoviride</i>	AAW31950
	<i>Trichoderma viride</i>	AAF19618
Hypocreaceae		
<i>Hypocrea</i>	<i>Hypocrea lixii</i> (1)	CAA55928
	<i>Hypocrea lixii</i> (2)	P48827
	<i>Hypocrea koningii</i>	AAF19612
	<i>Hypocrea rufa</i>	AAF19616
	<i>Hypocrea vinosa</i>	AAF19625

incided with the taxonomic status of these fungi, except two species *Aphanocladium album* (P32470) and *Verticillium fungicola* (AAP45631) (Table 2). All fungi analyzed here were from the same class Ascomycetes. Fungi in the first subgroup (clade I) belong to the order Hypocreales and com-

prises several families (Bionectriaceae, Clavicipitaceae, Hypocreaceae) that correspond to their additional groupings based on the chitinase sequences. The second subgroup (clade II) contained fungi from families Sordariales of the order Sordariomycetes (*N. crassa*), and from Orders

Leotiomycetes (*B. fuckeliana*), Eurotiomycetes (*A. fumigatus*) and Dothideomycetes (*C. minitans*).

Discussion

Clonostachys rosea displayed strong antagonistic activities against many soil and root-dwelling fungi. The success of *C. rosea* as a biocontrol agent is believed to involve many factors and diverse modes of action. The hydrolytic enzymes (e.g. protease and chitinase) produced by *C. rosea* likely play a key role in its ability to penetrate and kill a host. In this study, we cloned a chitinase gene (*Crchi1*) from *C. rosea* using the DNA walking technology. Comparison of the 5'-flanking DNA sequences of co-regulated genes may help identify conserved regulatory sequences (Gwynne *et al.*, 1987). Such regions may include binding sites for either general or special transcription regulatory proteins. Thus, this strategy was used to identify promoter elements that might be involved in the coordinated expression of *Crchi1*. The *Crchi1* promoter displayed eight consensus sites (5'-SY GGRG-3') for binding of the carbon catabolite repressor CreA (Sophianopoulou *et al.*, 1993). These sites are located at the upstream of the TATA box. One GATA site is present in the *Crchi1* promoter. The GATA sequences are recognized by the major positive regulatory elements AreA (in *Aspergillus nidulans*) and Gln3 (in *Saccharomyces cerevisiae*) (Kudla *et al.*, 1990; Minehart and Magasanik, 1991). Both AreA and Gln3 mediate global nitrogen repression and derepression. The role of these sequences in the regulation of *Crchi1* is unknown and requires further investigation.

A high level *Crchi1* expression was observed in medium with 0.5% CWRs (cell walls of *R. solani*) (Fig. 3A). This result suggests that a soluble component from the cell walls may act as an inducer for *Crchi1* expression. *C. rosea* has been reported as an efficient antagonist against *R. solani* on tobacco (Lahoz *et al.*, 2002). Therefore, chitinase CrChi1 may play a role during the interaction between *C. rosea* and the phytopathogen *R. solani*. In addition, 0.5% CCh (colloidal chitin) also induced a high level expression of *Crchi1*. Because purified colloidal chitin used in our assays is significantly different from the naturally occurring polymer in fungal cell walls, therefore, their induction potentials could also be very different. However, the expression of *Crchi1* could not be detected when the strain was grown in 0.5% periostracum cicadae, even though *C. rosea* was reported as an entomopathogenic fungus of *Oncometopia tucumana* and *Sonesimia grossa* (Toledo *et al.*, 2006). These results suggest other hydrolytic enzymes (e.g. protease) must be playing a significant role during infections against these hosts. In contrast, cultures grown in 1% glucose had a low level *Crchi1* mRNA. This was likely due to the lack of an inducer or the presence of a suppressor, since the promoter region of *Crchi1* contains putative binding sites for the regulatory protein CreA. The expression of *Crchi1* decreased with increasing glucose concentrations (Fig. 3B). In addition, our results showed that *Crchi1* mRNAs accumulated in fungi grown on 1% glucose in the presence of the 0.5% CCh (colloidal chitin), but reaching a low expression level. The low level expression suggests that glucose have a suppressive effect on the expression of *Crchi1*. Similarly, the expression

of *Crchi1* also decreased following the increase of colloidal chitin (Fig. 3C).

Several chitinases have been identified from the fungal species *Trichoderma* sp. (Seidl *et al.*, 2005). However, only the 42 kDa (Lorito *et al.*, 1998) and the 33 kDa proteins (Limon *et al.*, 1999) were found inducible to higher biological activity levels. In this study, the 42 kDa chitinases derived from various organisms were selected to investigate the evolutionary relationships between CrChi1 and other chitinases. Interestingly, except two species *A. album* and *V. fungicola*, the topology of the chitinase phylogenetic tree was very similar to the taxonomic relationships among these fungi inferred based on their structural and reproductive features. The incongruence for two of the species is understandable because some fungal species are highly pleomorphic and have anamorphic life cycles with widely divergent types of propagations (De Hoog *et al.*, 1994). Species in some of the genera cannot be identified using morphological characters alone. Additional physiological tests and sequence analyses may be needed. Recently, Schickler *et al.* (1998) developed an electrophoretic method using chitinases (Ech42) as a marker for identifying species in the genus *Trichoderma*. Therefore, the 42 kDa chitinases derived from ascomycetes may serve as a reference marker in future phylogenetic analyses. Moreover, the redundancy of the chitinase genes within the same species may reflect their functional differences between related proteins (Orikoshi *et al.*, 2003) and the adaptive evolution of the chitinases.

Acknowledgements

We thank Dr. Jianping Xu (McMaster University), Chenggang Zou and Yueshui Zhao for their help and advice in our studies. This work was funded by National Basic Research Program of China (2007CB411600), by the National Natural Science Foundation of China (approved nos. 30570059, 30630003 and 30660107), and by the Department of Science and Technology of Yunnan Province (approval Nos.2004C0001Z and 2005NG05).

References

- Baek, J.M., C.R Howell, and C.M. Kenerley. 1999. The role of an extracellular chitinase from *Trichoderma virens* Gv29-8 in the biocontrol of *Rhizoctonia solani*. *Curr. Genet.* 35, 41-50.
- Bogo, M.R., C.A. Rota, H.J. Pinto, M. Ocampos, C.T. Correa, M.H. Vainstein, and A. Schrank. 1998. A chitinase encoding gene (chit1 Gene) from the entomopathogen *Metarhizium anisopliae*: isolation and characterization of genomic and full-length cDNA. *Curr. Microbiol.* 37, 221-225.
- Carsolio, C., A. Gutierrez, B. Jimenez, M. Montagu, and A.E. Herrera. 1994. Characterization of ech-42, a *Trichoderma harzianum* endochitinase gene expressed during mycoparasitism. *Proc. Natl. Acad. Sci. USA* 91, 10903-10907.
- De Hoog, G.S., K. Takeo, S. Yoshida, E. Göttlich, K. Nishimura, and M. Miyaji. 1994. Pleoanamorphic life cycle of *Exophiala (Wangiella) dermatitidis*. *Antonie Van Leeuwenhoek* 65, 143-153.
- Gan, Z.W., J.K. Yang, N. Tao, L.M. Liang, Q.L. Mi, J. Li, and K.Q. Zhang. 2007. Cloning of the gene *Lecanicillium psalliotae* chitinase *Lpch1* and identification of its potential role in the biocontrol of root-knot nematode *Meloidogyne incognita*. *Appl. Microbiol. Biotechnol.* 76, 1309-1317.

- Garcia, I., J.M. Lora, J. Cruz, T. Benitez, A. Llobell, and J.A.T. Pintor. 1994. Cloning and characterization of a chitinase (chit42) cDNA from the mycoparasitic fungus *Trichoderma harzianum*. *Curr. Genet.* 27, 83-89.
- Gurr, S.J., S.E. Unkles, and J.R. Kinghorn. 1987. The structure and organisation of nuclear genes of filamentous fungi, p. 93-139. In J.L. Kinghorn (ed.), *Gene Structure in Lower Eukaryotes*. Oxford, IRL Press, UK.
- Gwynne, D.I., F.P. Buxton, S. Sibley, R.W. Davies, R.A. Lockington, C. Scazzocchio, and H.M. Sealy-Lewis. 1987. Comparison of the cis-acting control regions of two coordinately controlled genes involved in ethanol utilization in *Aspergillus nidulans*. *Gene* 51, 205-216.
- Kenneth, J.L. and D.S. Thomas. 2001. Analysis of relative gene expression data using Real-Time quantitative PCR and the $\Delta\Delta C_T$ method. *Methods* 25, 402-408.
- Kim, D.J., J.M. Baek, P. Uribe, C.M. Kenerley, and D.R. Cook. 2002. Cloning and characterization of multiple glycosyl hydrolase genes from *Trichoderma virens*. *Curr. Genet.* 40, 374-384.
- Krauss, U., P. Matthews, R. Bidwell, M. Hocart, and F. Anthony. 2001. Strain discrimination by fungal antagonists of *Colletotrichum musae*: implications for biocontrol of crown rot of banana. *Mycol. Res.* 105, 67-76.
- Kudla, B., M.X. Caddick, T. Langdon, N.M. Martinez-Rossi, C.F. Bennett, S. Sibley, R.W. Davies, and H.N. Arst. 1990. The regulatory gene are mediating nitrogen metabolite repression in *Aspergillus nidulans*. Mutations affecting specificity of gene activation alter a loop residue of a putative zinc finger. *EMBO J.* 9, 1355-1364.
- Lahoz, E., R. Nicoletti, F. Porrone, F. Raimo, L. Covarelli, and R. Contillo. 2002. Selection of fungal isolates with antagonistic effect against *Rhizoctonia solani* (AG 4 and AG 2-1 Nt) and growth promotion on tobacco. In *Proceedings of CORESTA Congress*, New Orleans, USA, 22-27 September.
- Li, J., J.K. Yang, X.W. Huang, and K.Q. Zhang. 2006. Purification and characterization of an extracellular serine protease from *Clonostachys rosea* and its potential as a pathogenic factor. *Process Biochem.* 41, 925-929.
- Limon, M.C., J.A. Pintor-Toro, and T. Benitez. 1999. Increased antifungal activity of *Trichoderma harzianum* transformants that overexpress a 33-KDa chitinase. *Phytopathology* 89, 254-261.
- Lorito, M., R.L. Mach, P. Sposato, J. Strauss, C.K. Peterbauer, and C.P. Kubicek. 1996. Mycoparasitic interaction relieves binding of Cre1 carbon catabolite repressor protein to promoter sequence of ech-42 (endochitinase encoding) gene of *Trichoderma harzianum*. *Proc. Natl. Acad. Sci. USA* 93, 14868-14872.
- Lorito, M., S.L. Woo, I. Garcia, G. Colucci, G.E. Harman, J.A. Pintor-Toro, E. Filippone, S. Muccifora, C.B. Lawrence, A. Zoina, S. Tuzun, F. Scala, and I.G. Fernandez. 1998. Genes from mycoparasitic fungi as a source for improving plant resistance to fungal pathogens. *Proc. Natl. Acad. Sci. USA* 95, 7860-7865.
- Margolles-Clark, E., G.E. Harman, and M. Penttila. 1996. Improved production of *Trichoderma harzianum* endochitinase by expression on *Trichoderma reesei*. *Appl. Environ. Microbiol.* 62, 2152-2155.
- Minehart, P.L. and B. Magasanik. 1991. Sequence and expression of GLN3, a positive nitrogen regulatory gene of *Saccharomyces cerevisiae* encoding a protein with a putative zinc finger DNA-binding domain. *Mol. Cell. Biol.* 11, 6216-6228.
- Orikoshi, H., N. Baba, S. Nakayama, H. Kashu, K. Miyamoto, M. Yasuda, Y. Inamori, and H. Tsujibo. 2003. Molecular analysis of the gene encoding a novel cold-adapted chitinase (ChiB) from a marine bacterium, *Alteromonas* sp. strain O-7. *J. Bacteriol.* 185, 1153-1160.
- Robert, L.M., K.P. Clemens, P. Kathrin, J. Sylvia, L.W. Sheridan, Z. Susanne, M.K. Cornelia, L. Matteo, and P.K. Christian. 1999. Expression of two major chitinase genes of *Trichoderma atroviride* (*T. harzianum* P1) is triggered by different regulatory signals. *Appl. Environ. Microbiol.* 5, 1858-1863.
- Schickler, H., B.C. Danin-Gehali, S. Haran, and I. Chet. 1998. Electrophoretic characterization of chitinases as a tool for the identification of *Trichoderma harzianum* strains. *Mycol. Res.* 102, 373-377.
- Schirmböck, M., M. Lorito, Y.L. Wang, C.K. Hayes, I. Arisan-Atac, F. Scala, G.E. Harman, and C.P. Kubicek. 1994. Parallel formation and synergism of hydrolytic enzymes and peptaibol antibiotics, molecular mechanisms involved in the antagonistic action of *Trichoderma harzianum* against phytopathogenic fungi. *Appl. Environ. Microbiol.* 60, 4364-4370.
- Seidl, V., B. Huemer, and C.P. Kubicek. 2005. A complete survey of *Trichoderma* chitinases reveals three distinct subgroups of family 18 chitinases. *FEBS J.* 272, 5923-5939.
- Seur, K.P., W.K. Chi, K. Hoon, S.J. Jae, and G.E. Harman. 2007. Cloning and high-level production of a chitinase from *Chromobacterium* sp. and the role of conserved or nonconserved residues on its catalytic activity. *Appl. Microbiol. Biotechnol.* 74, 791-804.
- Sophianopoulou, V., T. Suárez, G. Diallinas, and C. Scazzocchio. 1993. Operator derepressed mutations in the proline utilisation gene cluster of *Aspergillus nidulans*. *Mol. Gen. Genet.* 236, 209-213.
- Sutton, J.C., D.W. Li, P. Gang, Y. Hai, and P. Zhang. 1997. *Gliocladium roseum* a versatile adversary of a *Botrytis cinerea* in crops. *Plant Dis.* 81, 316-328.
- Swofford, D.L. 2002. PAUP*: Phylogenetic analysis using parsimony (*and other methods). Version 4.0b10. Sunderland, Massachusetts: Sinauer Associates, USA.
- Thompson, J.D., D.G. Higgins, and T.J. Gibson. 1994. CLUSTALW: improving the sensitivity of progressive multiple sequence alignment through sequence weighting, position-specific gap penalties and weight matrix choice. *Nucleic Acids Res.* 22, 4673-4680.
- Tiffin, P. 2004. Comparative evolutionary histories of chitinase genes in the genus *zea* and family poaceae. *Genetics* 167, 1331-1340.
- Toledo, A.V., E. Virla, R.A. Humber, S.L. Paradell, and L.C.C. López. 2006. First record of *Clonostachys rosea* (Ascomycota: Hypocreales) as an entomopathogenic fungus of *Oncometopia tucumana* and *Sonesimia grossa* (Hemiptera: Cicadellidae) in Argentina. *J. Invertebr. Pathol.* 92, 7-10.
- Xue, A.G. 2003. Biological control of pathogens causing root rot complex in weld pea using *Clonostachys rosea* strain ACM941. *Phytopathology* 93, 329-335.
- Yang, J.K., X.W. Huang, B.Y. Tian, H. Sun, J.X. Duan, W.P. Wu, and K.Q. Zhang. 2005. Characterization of an extracellular serine protease gene from the nematophagous fungus *Lecanicillium psalliotae*. *Biotechnol. Lett.* 27, 1329-1334.
- Zhang, D., Y. Yang, L.A. Castlebury, and C.E. Cerniglia. 1996. A method for the large scale isolation of high transformation efficiency fungal genomic DNA. *FEMS Microbiol. Lett.* 145, 216-265.
- Zhao, M.L., J.S. Huang, M.H. Mo, and K.Q. Zhang. 2005. A potential virulence factor involved in fungal pathogenicity serine-like protease activity of nematophagous fungus *Clonostachys rosea*. *Fungal Divers.* 19, 217-234.