

## MONOTONE ITERATION SCHEME FOR IMPULSIVE THREE-POINT NONLINEAR BOUNDARY VALUE PROBLEMS WITH QUADRATIC CONVERGENCE

BASHIR AHMAD, AHMED ALSAEDI, AND DOA'A GAROUT

ABSTRACT. In this paper, we consider an impulsive nonlinear second order ordinary differential equation with nonlinear three-point boundary conditions and develop a monotone iteration scheme by relaxing the convexity assumption on the function involved in the differential equation and the concavity assumption on nonlinearities in the boundary conditions. In fact, we obtain monotone sequences of iterates (approximate solutions) converging quadratically to the unique solution of the impulsive three-point boundary value problem.

### 1. Introduction

In recent years, a number of research papers have dealt with dynamical systems with impulse effect as a class of general hybrid systems. Impulsive hybrid systems are composed of some continuous variable dynamic systems along with certain reset maps that define impulsive switching among them. It is the switching that resets the modes and changes the continuous state of the system. There are three classes of impulsive hybrid systems, namely, impulsive differential systems [33, 41], sampled data or digital control system [30, 44] and impulsive switched system [19, 24]. Applications of such systems include air traffic management [43], automotive control [5, 9], real-time software verification [6], transportation systems [37, 45], manufacturing [39], mobile robotics [10], process industry [25], etc. In fact, hybrid systems have a central role in embedded control systems that interact with the physical world. Using hybrid models, one may represent time and event-based behaviors more accurately so as to meet challenging design requirements in the design of control systems for problems such as cut-off control and idle speed control of the engine. For more details, see [7] and the references therein.

Impulsive differential equations, which provide a natural description of observed evolution processes, are regarded as important mathematical tools for

---

Received December 14, 2006.

2000 *Mathematics Subject Classification.* 34A37, 34B10, 34B15.

*Key words and phrases.* quasilinearization, three-point boundary value problems, quadratic convergence.

the better understanding of several real world problems in biology, physics, engineering, etc. Thus, the theory of impulsive differential equations is much richer than the corresponding theory of ordinary differential equations without impulse effects since a simple impulsive differential equation may exhibit several new phenomena such as rhythmical beating, merging of solutions and noncontinuability of solutions. For the theoretical aspect of impulsive differential equations, we refer the reader to the references [3, 4, 8, 13, 14, 16, 21, 33, 36, 40, 41, 47] whereas the applications of impulsive differential equations can be found in [15, 18, 26, 42].

Multi-point nonlinear boundary value problems which take into account the boundary data at intermediate points of the interval under consideration, have been addressed by many authors, for example, see [22, 27-29]. In this paper, we develop an extended method of quasilinearization (the generalized quasilinearization) for a nonlinear impulsive three-point boundary value problem given by

$$(1.1) \quad x''(t) = f(t, x(t), x'(t)), \quad t_k < t < t_{k+1}, \quad k = 0, 1, 2, \dots, m,$$

$$(1.2) \quad px(0) - qx'(0) = g_1(x(\frac{1}{2})), \quad px(1) + qx'(1) = g_2(x(\frac{1}{2})),$$

and for  $k = 1, 2, \dots, m$ ,

$$\Delta x(t_k) = u_k,$$

$$(1.3) \quad \Delta x'(t_k) = v_k(x(t_k), x'(t_k)),$$

where  $f : [0, 1] \times \mathbb{R}^2 \rightarrow \mathbb{R}$  is continuous,  $p, q$  are chosen appropriately with  $p > 1$  and  $g_{1,2} : \mathbb{R} \rightarrow \mathbb{R}$  are continuous,  $u_k \in \mathbb{R}$  and  $v_k : \mathbb{R}^2 \rightarrow \mathbb{R}$  are continuous for  $k = 1, 2, \dots, m$ . This work is motivated by Doddaballapur, Eloë, and Zhang [17] who discussed the method of quasilinearization for two-point boundary value problems with impulse. The origin of the quasilinearization lies in the theory of dynamic programming [11, 12]. This method applies to semilinear equations with convex or concave nonlinearities and generates a monotone iteration scheme whose iterates converge quadratically to a unique solution of the given problem. The nineties brought new dimensions to this technique when Lakshmikantham [31, 32] generalized the method of quasilinearization by relaxing the concavity/convexity assumption. This development proved to be of immense value and the method was extensively developed and applied to a wide range of initial and boundary value problems for different types of differential equations, see [1, 22, 23, 34, 35] and references therein. Some real-world applications of the quasilinearization technique can be found in [2, 38, 46]. In Section 2, we present some basic results which play an important role to prove the main result of the paper. In Section 3, we apply the method of generalized quasilinearization to obtain monotone sequences of upper and lower solutions that converge quadratically to the unique solution of the problem (1.1)-(1.3).

### 2. Preliminary results

Let  $PC[0, 1]$  denote the class of piecewise continuous functions on  $[0, 1]$  and  $PC^1[0, 1]$  denote the class of functions  $x$  such that  $x \in PC[0, 1]$  and  $x' \in PC[0, 1]$ . Define an appropriate Banach space  $B$  by

$$B = \{x \in PC^1[0, 1] : x^{(i)}|_{[t_k, t_{k+1}]} \in C^i[t_k, t_{k+1}], k = 0, 1, \dots, m, i = 0, 1\},$$

with

$$\|x\|_B = \max_{k=0,1,\dots,m} \|x\|_k \text{ and } \|x\|_k = \max_{i=0,1} \sup_{t_k \leq t \leq t_{k+1}} |x^{(i)}(t)|.$$

For the fixed impulsive points  $t_k$  satisfying  $0 = t_0 < t_1 < \dots < t_m < t_{m+1} = 1$ , we define the impulse by  $\Delta x(t_k) = x(t_k^+) - x(t_k^-)$  with the convention  $x(t_k) = x(t_k^-)$ ,  $k = 1, \dots, m$ .

We say that  $\alpha \in B$  is a lower solution of the BVP with impulse (1.1)-(1.3) if

$$\begin{aligned} \alpha''(t) &\geq f(t, \alpha(t), \alpha'(t)), \quad t_k < t < t_{k+1}, \quad k = 0, 1, \dots, m, \\ p\alpha(0) - q\alpha'(0) &\leq g_1(\alpha(\frac{1}{2})), \quad p\alpha(1) + q\alpha'(1) \leq g_2(\alpha(\frac{1}{2})), \end{aligned}$$

and for  $k = 1, 2, \dots, m$ ,

$$\Delta\alpha(t_k) = u_k,$$

$$\Delta\alpha'(t_k) \geq v_k(\alpha(t_k), \alpha'(t_k)).$$

Analogously,  $\beta \in B$  is an upper solution of the BVP with impulse (1.1)-(1.3) if

$$\begin{aligned} \beta''(t) &\leq f(t, (\beta(t), \beta'(t)), \quad t_k < t < t_{k+1}, \quad k = 0, 1, \dots, m, \\ p\beta(0) - q\beta'(0) &\geq g_1(\beta(\frac{1}{2})), \quad p\beta(1) + q\beta'(1) \geq g_2(\beta(\frac{1}{2})), \end{aligned}$$

and for  $k = 1, 2, \dots, m$ ,

$$\Delta\beta(t_k) = u_k,$$

$$\Delta\beta'(t_k) \leq v_k(\beta(t_k), \beta'(t_k)).$$

Finally, we define a partial order on  $B$  as

$$\alpha|_{[t_k, t_{k+1}]}(t) \leq \beta|_{[t_k, t_{k+1}]}(t), \quad t_k < t < t_{k+1}, \quad k = 0, 1, \dots, m.$$

As argued in [17],  $x$  is a solution of BVP with impulse (1.1)-(1.3) if and only if  $x \in B$  and  $Tx = x$ , where  $T$  is a fixed point operator defined by

$$(2.1) \quad Tx(t) = p(t) + I(t, x) + \int_0^1 G(t, s)f(s, x(s), x'(s))ds,$$

where

$$p(t) = g_1(x(\frac{1}{2}))(\frac{-t}{p+2q} + \frac{p+q}{p^2+2pq}) + g_2(x(\frac{1}{2}))(\frac{t}{p+2q} + \frac{q}{p^2+2pq}),$$

$$G(t, s) = \frac{1}{(p^2+2pq)} \begin{cases} (pt+q)(p(s-1)-q), & \text{if } 0 \leq t \leq s \leq 1, \\ (p(t-1)-q)(ps+q), & \text{if } 0 \leq s \leq t \leq 1, \end{cases}$$

and  $I(t, x) = \sum_{k=1}^m I_k(t, x)$  with

$$I_k(t, x) = \frac{1}{p^2 + 2pq} \begin{cases} (pt + q)[-u_k p + (p(t_k - 1) - q)v_k(x(t_k), x'(t_k))], & 0 \leq t \leq t_k, \\ (p(t - 1) - q)[-u_k p + (pt_k + q)v_k(x(t_k), x'(t_k))], & t_k \leq t \leq 1. \end{cases}$$

Now, we are in a position to present some basic results. For the sake of completeness, we provide the proof of these theorems although the method of proof is similar to the one employed in [17].

**Theorem 2.1.** *Assume that*

- (i)  $\alpha, \beta \in B$  are lower and upper solutions of the BVP with impulse (1.1)-(1.3) respectively.
- (ii)  $f, f_x \in C([0, 1] \times \mathbb{R}^2)$  such that  $f_x(t, x, y) > 0$  and  $v_k \in C^1(\mathbb{R}^2)$  with  $v_{kx}(x, y) > 0, v_{ky}(x, y) > 0, (x, y) \in \mathbb{R}^2$  for  $k = 1, 2, \dots, m$ .
- (iii)  $g_i$  are continuous on  $\mathbb{R}$  and satisfy one-sided Lipschitz condition:

$$g_i(x) - g_i(y) \leq L_i(x - y), \quad 0 \leq L_i < 1, \quad i = 1, 2.$$

Then  $\alpha(t) \leq \beta(t)$ .

*Proof.* Let us set  $w(t) = \alpha(t) - \beta(t)$ . For the sake of contradiction, we assume that  $w$  is positive on  $[0, 1]$ , that is,  $\alpha(t) > \beta(t)$ . By definition of upper and lower solutions and using (iii), we find that

$$\begin{aligned} pw(0) - qw'(0) &\leq g_1(\alpha(\frac{1}{2})) - g_1(\beta(\frac{1}{2})) \\ &\leq L_1(\alpha(\frac{1}{2}) - \beta(\frac{1}{2})) \\ &< \alpha(\frac{1}{2}) - \beta(\frac{1}{2}) = w(\frac{1}{2}). \end{aligned}$$

From this inequality, it follows that  $w(t)$  does not have a local positive maximum at  $\tau = 0$  as  $w'(0) = 0$  implies that  $w(0) < w(\frac{1}{2})$  for  $p > 1$ . Similarly, it can be shown that  $w(t)$  cannot have a local positive maximum at  $\tau = 1$ . Thus,  $w$  has a positive maximum at some  $\tau \in (0, 1)$ . Let us assume that  $\tau \in \bigcup_{k=0}^m (t_k, t_{k+1})$ . Then  $w''(\tau) \leq 0$  and  $\alpha'(\tau) = \beta'(\tau)$ . Since  $\alpha$  and  $\beta$  are lower and upper solutions of the BVP with impulse (1.1)-(1.3) and  $f_x > 0$ , it follows that

$$w''(\tau) = \alpha''(\tau) - \beta''(\tau) \geq f(\tau, \alpha(\tau), \alpha'(\tau)) - f(\tau, \beta(\tau), \beta'(\tau)) > 0.$$

This provides a contradiction and so,  $\tau \notin \bigcup_{k=0}^m (t_k, t_{k+1})$ .

Now, we assume that  $\tau = t_k$  for some  $k \in \{1, 2, \dots, m\}$ . Then, by Taylor's theorem,  $w'(t_k^-) \geq 0$  and  $w'(t_k^+) \leq 0$ , or  $\Delta w'(t_k) \leq 0$  and

$$\alpha'(t_k^-) = \alpha'(t_k) \geq \beta'(t_k) = \beta'(t_k^-).$$

In view of the fact that  $v_{kx}(x, y) > 0$ ,  $v_{ky}(x, y) > 0$ ,  $(x, y) \in \mathbb{R}^2$ , it follows that

$$\Delta w'(t_k) = \Delta \alpha'(t_k) - \Delta \beta'(t_k) \geq v_k(\alpha(t_k), \alpha'(t_k)) - v_k(\beta(t_k), \beta'(t_k)) > 0,$$

which is a contradiction and thus  $\tau \notin \{t_1, \dots, t_m\}$ . Hence, we conclude that  $\alpha(t) \leq \beta(t)$ ,  $t \in [0, 1]$ . This completes the proof.  $\square$

**Theorem 2.2.** Assume that  $f \in C([0, 1] \times \mathbb{R}^2)$ ,  $z_k \in C(\mathbb{R}^2)$ ,  $k = 1, \dots, m$  and each  $z_k(x, y)$  is monotone increasing in  $y$  for fixed  $x$ . Further, each solution of  $x''(t) = f(t, x(t), x'(t))$  extends to  $[0, 1]$  or becomes unbounded on its maximal interval of convergence. Let  $\alpha$ ,  $\beta$  be lower and upper solutions respectively of the BVP

$$(2.2) \quad \begin{aligned} x''(t) &= f(t, x(t), x'(t)), \quad t_k < t < t_{k+1}, \\ \Delta x(t_k) &= u_k, \end{aligned}$$

$$(2.3) \quad \Delta x'(t_k) = z_k(x(t_k), x'(t_k)), \quad k = 1, 2, \dots, m,$$

with boundary conditions (1.2) such that  $\alpha \leq \beta$ . Further,  $g_i$  ( $i = 1, 2$ ) in (1.2) are continuous on  $\mathbb{R}$  and satisfy one-sided Lipschitz condition. Then, there exists a solution  $x$  of the BVP with impulse (2.2)-(2.3), (1.2) satisfying  $\alpha \leq x \leq \beta$ .

*Proof.* We define

$$\hat{f}(t, x, y) = \begin{cases} f(t, \beta(t), y) + \frac{x - \beta(t)}{1 + (x - \beta(t))}, & \text{if } x(t) > \beta(t), \\ f(t, x, y), & \text{if } \alpha(t) \leq x(t) \leq \beta(t), \\ f(t, \alpha(t), y) + \frac{x - \alpha(t)}{1 + |x - \alpha(t)|}, & \text{if } x(t) < \alpha(t), \end{cases}$$

$$\hat{v}_k(x, y) = \begin{cases} z_k(\beta(t_k), y) + \frac{x - \beta(t_k)}{1 + (x - \beta(t_k))}, & \text{if } x > \beta(t_k), \\ z_k(x, y), & \text{if } \alpha(t_k) \leq x \leq \beta(t_k), \\ z_k(\alpha(t_k), y) + \frac{x - \alpha(t_k)}{1 + |x - \alpha(t_k)|}, & \text{if } x < \alpha(t_k) \end{cases}$$

for  $k = 1, 2, \dots, m$  and

$$G_i(x) = \begin{cases} g_i(\beta(\frac{1}{2})), & \text{if } x > \beta(\frac{1}{2}), \\ g_i(x), & \text{if } \alpha(\frac{1}{2}) \leq x \leq \beta(\frac{1}{2}), \\ g_i(\alpha(\frac{1}{2})), & \text{if } x < \alpha(\frac{1}{2}) \end{cases}$$

for  $i = 1, 2$ . Let  $N > 0$  be such that  $|\alpha'(t)| \leq N$ ,  $|\beta'(t)| \leq N$ ,  $t \in [t_k, t_{k+1}]$ ,  $k = 0, 1, \dots, m$ .

For each positive integer  $l$ , define

$$f_l(t, x, y) = \begin{cases} \hat{f}(t, x, N + l), & \text{if } y > N + l, \\ \hat{f}(t, x, y), & \text{if } |y| \leq N + l, \\ \hat{f}(t, x, -(N + l)), & \text{if } y < -(N + l), \end{cases}$$

and

$$v_{k,l}(x, y) = \begin{cases} \hat{v}_k(x, N + l), & \text{if } y > N + l, \\ \hat{v}_k(x, y), & \text{if } |y| \leq N + l, \\ \hat{v}_k(x, -(N + l)), & \text{if } y < -(N + l). \end{cases}$$

Notice that  $f_l$ ,  $G_i$  and  $v_{kl}$  are bounded and continuous on their respective domains. With a standard application of the Schauder fixed point theorem to the operator  $T$ , defined by (2.1), we can obtain a solution,  $x_l \in B$ , to the BVP with impulse (2.2)-(2.3), (1.2) with  $f = f_l$ , each  $G_i = g_i$  and each  $v_k = v_{kl}$ .

Now, we want to show that each solution  $x_l$  satisfies  $\alpha \leq x_l \leq \beta$ . We set  $r(t) = x_l(t) - \beta(t)$  and assume that  $r(t)$  has a positive maximum at  $\tau \in (0, 1)$  as in the proof of Theorem 2.1. If  $\tau \in \bigcup_{k=0}^m (t_k, t_{k+1})$ , then  $r''(\tau) \leq 0$ , that is,  $x_l''(\tau) \leq \beta''(\tau)$ , and  $|x_l'(\tau)| = |\beta'(\tau)| \leq N < N + l$ . Thus

$$\begin{aligned} r''(\tau) &= x_l''(\tau) - \beta''(\tau) \\ &\geq f(\tau, \beta(\tau), x_l'(\tau)) + \frac{x_l(\tau) - \beta(\tau)}{1 + (x_l(\tau) - \beta(\tau))} - f(\tau, \beta(\tau), \beta'(\tau)) \\ &= \frac{x_l(\tau) - \beta(\tau)}{1 + (x_l(\tau) - \beta(\tau))} > 0, \end{aligned}$$

which is a contradiction. Now, suppose that  $\Delta r'(\tau) \leq 0$  for  $\tau = t_k$ ,  $k = 1, 2, \dots, m$ . Since each  $z_k(x, y)$  is monotone increasing in  $y$  for fixed  $x$ , it follows that each  $v_{kl}(x, y)$  is monotone increasing in  $y$  for fixed  $x$ . Moreover, we note that each  $v_{kl}(\beta(t_k), \beta'(t_k)) = z_k(\beta(t_k), \beta'(t_k))$ . Thus,

$$\begin{aligned} \Delta r'(t_k) &\geq v_{kl}(\beta(t_k), x_l'(t_k)) - v_{kl}(\beta(t_k), \beta'(t_k)) + \frac{x_l(t_k) - \beta(t_k)}{1 + (x_l(t_k) - \beta(t_k))} \\ &\geq \frac{x_l(t_k) - \beta(t_k)}{1 + (x_l(t_k) - \beta(t_k))} > 0, \end{aligned}$$

which is also a contradiction. Therefore,  $r(t) \leq 0$  or  $x_l \leq \beta$ . On the same pattern, we can prove that  $\alpha(t) \leq x_l(t)$ .

For each  $l$ , there exists  $t_l \in [0, t_1]$  such that

$$t_1 |x'_{kl}(t_l)| = |x_{kl}(t_1) - x_{kl}(0)| \leq \max\{|\beta(0) - \alpha(t_1)|, |\beta(t_1) - \alpha(0)|\}.$$

Thus, the sequences  $\{x_{kl}(t_l)\}$  and  $\{x'_{kl}(t_l)\}$  are bounded. Now, applying the Kamke convergence theorem for solutions of initial value problems, we obtain a subsequence of  $\{x_{kl}\}$  which converges to a solution of  $x''(t) = \hat{f}(t, x(t), x'(t))$  on a maximal subinterval of  $[0, t_1]$ . Clearly,  $\alpha(t) \leq x(t) \leq \beta(t)$  and the solutions of  $x''(t) = f(t, x(t), x'(t))$  extend to all of  $[0, 1]$  or become unbounded implying that  $x''(t) = \hat{f}(t, x(t), x'(t))$  on  $[0, t_1]$ .

Now, we apply the impulse  $\Delta x'(t_k) = z_k(x(t_k), x'(t_k))$  at  $t_1$ . Applying the Kamke theorem on the subsequence obtained earlier, we can employ  $t_1 = t_l$  for each  $l$ . Thus, we can obtain a further subsequence which converges to a solution  $x$  of  $x''(t) = \hat{f}(t, x(t), x'(t))$  on  $(0, t_1) \cup (t_1, t_2)$  such that  $x$  satisfies the impulse  $\Delta x'(t_k) = z_k(x(t_k), x'(t_k))$  at  $t_1$ . This procedure can be continued inductively by first applying the impulse at each  $t_j$  and then applying the Kamke convergence theorem on the subinterval  $(t_j, t_{j+1})$ . Hence  $\alpha(t) \leq x(t) \leq \beta(t)$  and  $\hat{f}(t, x(t), x'(t)) = f(t, x(t), x'(t))$ . This completes the proof.  $\square$

*Remark.* The simplified version of the condition that each solution of  $x''(t) = f(t, x(t), x'(t))$  extends to  $[0, 1]$  or becomes unbounded on its maximal interval of convergence is that  $f$  satisfies a Nagumo condition, that is, for each  $M > 0$ , there exists a positive continuous function  $h_M$  on  $[0, \infty)$  such that  $|f(t, x, x')| \leq h_M(|x'|)$  for all  $(t, x, x') \in [0, 1] \times [-M, M] \times \mathbb{R}$  and

$$\int_0^\infty (s/h_M(s))ds = \infty.$$

### 3. Iteration scheme with quadratic convergence

**Theorem 3.1.** *Assume that*

- (A<sub>0</sub>)  $\alpha_0, \beta_0 \in B$  are lower and upper solutions of the BVP with impulse (1.1)-(1.3) respectively.
- (A<sub>1</sub>)  $f(t, x, y) \in C([0, 1] \times \mathbb{R}^2)$  be such that  $f_x(t, x, y) > 0$ ,  $\frac{\partial^2}{\partial x^2}(f(t, x, y) + \phi_1(t, x, y)) \geq 0$ , where  $\frac{\partial^2}{\partial x^2}\phi_1(t, x, y) \geq 0$  for some continuous function  $\phi_1(t, x, y)$ . Moreover,  $f$  satisfies a Nagumo condition in  $x'$ .
- (A<sub>2</sub>) For  $i = 1, 2$ ,  $g_i(x), g'_i(x)$  are continuous on  $\mathbb{R}$  with  $0 \leq g'_i \leq 1$  and  $g''_i(x) + \psi''_i(x) \leq 0$  with  $\psi''_i \leq 0$  for some continuous functions  $\psi_i(x)$  on  $\mathbb{R}$ .
- (A<sub>3</sub>)  $v_k \in C^1(\mathbb{R}^2)$  such that  $v_{kx}(x, y) > 0$ ,  $v_{ky}(x, y) > 0$ ,  $(x, y) \in \mathbb{R}^2$  and  $v''_k \in C(\mathbb{R}^2)$  for  $k = 1, 2, \dots, m$ .

Then there exists monotone sequences  $\{\alpha_n\}$  and  $\{\beta_n\}$ , which converge in  $B$  to a unique solution  $x$  of the BVP with impulse (1.1)-(1.3) and the convergence is quadratic.

*Proof.* Let  $F(t, x) : [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$  be such that  $F, F_x$  and  $F_{xx}$  are continuous on  $[0, 1] \times \mathbb{R}$  and it follows from (A<sub>1</sub>) that  $F_{xx} \geq 0$  on  $[0, 1] \times \mathbb{R}$ , where we have set [23]

$$f(t, x_1, x_2) = F(t, x_1) - \phi_1(t, x_1, x_2).$$

Applying the mean value theorem on  $F(t, x_1)$  gives

$$F(t, x_1) \geq F(t, y_1) + F_x(t, y_1)(x_1 - y_1),$$

which together with the definition of  $F(t, x_1)$  takes the form

$$(3.1) \quad f(t, x_1, x_2) \geq f(t, y_1, y_2) + F_x(t, y_1)(x_1 - y_1) - \phi_1(t, x_1, x_2) + \phi_1(t, y_1, y_2)$$

for  $x_1, y_1, x_2, y_2 \in \mathbb{R}$ .

We define  $G_i : \mathbb{R} \rightarrow \mathbb{R}$  by  $G_i(x) = g_i(x) + \psi_i(x)$ ,  $i = 1, 2$ . From (A<sub>2</sub>), it follows that

$$(3.2) \quad g_i(x) \leq g_i(y) + G'_i(y)(x - y) + \psi_i(y) - \psi_i(x), i = 1, 2, x, y \in \mathbb{R}.$$

For each  $k = 1, \dots, m$ , let  $V_k(x) : \mathbb{R} \rightarrow \mathbb{R}$  be such that  $V_k, V'_k, V''_k$  are continuous on  $\mathbb{R}$  with  $V''_k(x) \geq 0$ ,  $x \in \mathbb{R}$ . Let us set  $\phi_k(x_1, x_2) = V'_k(x_1) - v_k(x_1, x_2)$

on  $\mathbb{R}^2$ . By the mean value theorem, we have  $V_k(x_1) \geq V_k(y_1) + V'_k(y_1)(x_1 - y_1)$  for  $x_1, y_1 \in \mathbb{R}$ , which can alternatively be written as

$$(3.3) \quad v_k(x_1, x_2) \geq v_k(y_1, y_2) + V'_k(y_1)(x_1 - y_1) - \phi_k(x_1, x_2) + \phi_k(y_1, y_2)$$

for  $x_1, x_2, y_1, y_2 \in \mathbb{R}$ . Now, we define

$$\begin{aligned} Q(t, x_1, x_2; \alpha_0, \beta_0, \alpha'_0) &= f(t, \alpha_0(t), \alpha'_0(t)) + F_x(t, \beta_0(t))(x_1 - \alpha_0(t)) \\ &\quad - \phi_1(t, x_1, x_2) + \phi_1(t, \alpha_0(t), \alpha'_0(t)), \\ J(t, x_1, x_2; \beta_0, \beta'_0) &= f(t, \beta_0(t), \beta'_0(t)) + F_x(t, \beta_0(t))(x_1 - \beta_0(t)) \\ &\quad - \phi_1(t, x_1, x_2) + \phi_1(t, \beta_0(t), \beta'_0(t)), \\ h_k(x_1, x_2; \alpha_0, \beta_0, \alpha'_0) &= v_k(\alpha_0(t_k), \alpha'_0(t_k)) + V'_k(\beta_0(t_k))(x_1 - \alpha_0(t_k)) \\ &\quad - \phi_k(x_1, x_2) + \phi_k(\alpha_0(t_k), \alpha'_0(t_k)), \\ H_k(x_1, x_2; \beta_0, \beta'_0) &= v_k(\beta_0(t_k), \beta'_0(t_k)) + V'_k(\beta_0(t_k))(x_1 - \beta_0(t_k)) \\ &\quad - \phi_k(x_1, x_2) + \phi_k(\beta_0(t_k), \beta'_0(t_k)), \\ r_i(x(\frac{1}{2}); \alpha_0, \beta_0) &= g_i(\alpha_0(\frac{1}{2})) + G'_i(\beta_0(\frac{1}{2}))(x(\frac{1}{2}) - \alpha_0(\frac{1}{2})) \\ &\quad + \psi_i(\alpha_0(\frac{1}{2})) - \psi_i(x(\frac{1}{2})), i = 1, 2, \\ R_i(x(\frac{1}{2}); \beta_0) &= g_i(\beta_0(\frac{1}{2})) + G'_i(\beta_0(\frac{1}{2}))(x(\frac{1}{2}) - \beta_0(\frac{1}{2})) \\ &\quad + \psi_i(\beta_0(\frac{1}{2})) - \psi_i(x(\frac{1}{2})), i = 1, 2. \end{aligned}$$

Observe that

$$(3.4) \quad \begin{aligned} Q(t, \alpha_0(t), \alpha'_0(t); \alpha_0, \beta_0, \alpha'_0) &= f(t, \alpha_0(t), \alpha'_0(t)), \\ f(t, x_1, x_2) &\geq Q(t, x_1, x_2; \alpha_0, \beta_0, \alpha'_0). \end{aligned}$$

$$(3.5) \quad r_i(\alpha_0(\frac{1}{2}); \alpha_0, \beta_0) = g_i(\alpha_0(\frac{1}{2})), \quad g_i(x) \leq r_i(x(\frac{1}{2}); \alpha_0, \beta_0) \quad i = 1, 2.$$

$$(3.6) \quad \begin{aligned} h_k(\alpha_0(t_k), \alpha'_0(t_k); \alpha_0, \beta_0, \alpha'_0) &= v_k(\alpha_0(t_k), \alpha'_0(t_k)), \\ v_k(x_1(t), x_2(t)) &\geq h_k(x_1, x_2; \alpha_0, \beta_0, \alpha'_0). \end{aligned}$$

Also

$$(3.7) \quad J(t, \beta_0(t), \beta'_0(t); \beta_0, \beta'_0) = f(t, \beta_0(t), \beta'_0(t)), \quad f(t, x_1, x_2) \leq J(t, x_1, x_2; \beta_0, \beta'_0).$$

$$(3.8) \quad R_i(\beta_0(\frac{1}{2}); \beta_0) = g_i(\beta_0(\frac{1}{2})), \quad g_i(x) \geq R_i(x(\frac{1}{2}); \beta_0), \quad i = 1, 2.$$

$$(3.9) \quad \begin{aligned} H_k(\beta_0(t_k), \beta'_0(t_k); \beta_0, \beta'_0) &= v_k(\beta_0(t_k), \beta'_0(t_k)), \\ v_k(x_1(t), x_2(t)) &\leq H_k(x_1, x_2; \beta_0, \beta'_0). \end{aligned}$$



Consider the BVP with impulse

$$(3.10) \quad x''(t) = Q(t, x(t), x'(t); \alpha_0, \beta_0, \alpha'_0), \quad t_k < t < t_{k+1}, \quad k = 0, \dots, m,$$

$$(3.11) \quad px(0) - qx'(0) = r_1(x(\frac{1}{2}); \alpha_0, \beta_0), \quad px(1) + qx'(1) = r_2(x(\frac{1}{2}); \alpha_0, \beta_0),$$

and for  $k = 1, \dots, m$ ,

$$\Delta x(t_k) = u_k,$$

$$(3.12) \quad \Delta x'(t_k) = h_k(x(t_k), x'(t_k); \alpha_0, \beta_0, \alpha'_0).$$

We show that  $\alpha_0$  and  $\beta_0$  are respectively lower and upper solutions of the BVP with impulse (3.10)-(3.12).

Using (3.4)-(3.6) together with the fact that  $\alpha_0$  is a lower solution of (1.1)-(1.3), we obtain

$$\begin{aligned} \alpha''_0(t) &\geq f(t, \alpha_0(t), \alpha'_0(t)) \\ &= Q(t, \alpha_0(t), \alpha'_0(t); \alpha_0, \beta_0, \alpha'_0), \quad t_k < t < t_{k+1}, \quad k = 0, 1, \dots, m, \\ p\alpha_0(0) - q\alpha'_0(0) &\leq g_1(\alpha_0(\frac{1}{2})) = r_1(\alpha_0(\frac{1}{2}); \alpha_0, \beta_0), \\ p\alpha_0(1) + q\alpha'_0(1) &\leq g_2(\alpha_0(\frac{1}{2})) = r_2(\alpha_0(\frac{1}{2}); \alpha_0, \beta_0), \end{aligned}$$

and for  $k = 1, \dots, m$ ,

$$\Delta \alpha_0(t_k) = u_k,$$

$$\Delta \alpha'_0(t_k) \geq v_k(\alpha_0(t_k), \alpha'_0(t_k)) = h_k(\alpha_0(t_k), \alpha'_0(t_k); \alpha_0, \beta_0, \alpha'_0),$$

which implies that  $\alpha_0$  is a lower solution of (3.10)-(3.12).

Using (3.1) and the fact that  $\beta_0$  is an upper solution of (1.1)-(1.3), we get

$$\begin{aligned} \beta''_0(t) &\leq f(t, \beta_0(t), \beta'_0(t)) \\ &\leq f(t, \alpha_0(t), \alpha'_0(t)) - F_x(t, \beta_0(t))(\alpha_0(t) - \beta_0(t)) \\ &\quad + \phi_1(t, \alpha_0(t), \alpha'_0(t)) - \phi_1(t, \beta_0(t), \beta'_0(t)) \\ &= Q(t, \beta_0(t), \beta'_0(t); \alpha_0, \beta_0, \alpha'_0), \quad t_k < t < t_{k+1}, \quad k = 0, 1, \dots, m. \end{aligned}$$

Using the mean value theorem and the non-increasing property of  $G'_1$ , we have

$$\begin{aligned} &g_1(\beta_0(\frac{1}{2})) - r_1(\beta_0(\frac{1}{2}); \alpha_0, \beta_0) \\ &= g_1(\beta_0(\frac{1}{2})) - g_1(\alpha_0(\frac{1}{2})) - G'_1(\beta_0(\frac{1}{2}))(\beta_0(\frac{1}{2}) - \alpha_0(\frac{1}{2})) \\ &\quad - \psi_1(\alpha_0(\frac{1}{2})) + \psi_1(\beta_0(\frac{1}{2})) \\ &= G_1(\beta_0(\frac{1}{2})) - G_1(\alpha_0(\frac{1}{2})) - G'_1(\beta_0(\frac{1}{2}))(\beta_0(\frac{1}{2}) - \alpha_0(\frac{1}{2})) \\ &= (G'_1(c_0) - G'_1(\beta_0(\frac{1}{2}))) (\beta_0(\frac{1}{2}) - \alpha_0(\frac{1}{2})) \geq 0, \end{aligned}$$

$(\alpha_0(\frac{1}{2}) \leq c_0 \leq \beta_0(\frac{1}{2}))$  which yields

$$p\beta_0(0) - q\beta'_0(0) \geq g_1(\beta_0(\frac{1}{2})) \geq r_1(\beta_0(\frac{1}{2}); \alpha_0, \beta_0).$$

Similarly, it can be shown that

$$p\beta_0(1) + q\beta'_0(1) \geq r_2(\beta_0(\frac{1}{2}); \alpha_0, \beta_0).$$

In view of (3.3), we get

$$\begin{aligned} \Delta\beta'_0(t_k) &\leq v_k(\beta_0(t_k), \beta'_0(t_k)) \\ &\leq v_k(\alpha_0(t_k), \alpha'_0(t_k)) - V'_k(\beta_0(t_k))(\alpha_0(t_k) - \beta_0(t_k)) \\ &\quad + \phi_k(\alpha_0(t_k), \alpha'_0(t_k)) - \phi_k(\beta_0(t_k), \beta'_0(t_k)) \\ &= h_k(\beta_0(t_k), \beta'_0(t_k); \alpha_0, \beta_0, \alpha'_0), \quad k = 1, \dots, m. \end{aligned}$$

Thus, we conclude that  $\beta_0$  is an upper solution of the BVP with impulse (3.10)-(3.12). Moreover, each  $h_k$  satisfies the hypotheses of Theorem 2.2 and a limit comparison implies that  $Q$  satisfies a Nagumo condition in  $x'$ . Hence, by Theorem 2.2, there exists a solution  $\alpha_1$  of the BVP with impulse (3.10)-(3.12) such that

$$(3.13) \quad \alpha_0(t) \leq \alpha_1(t) \leq \beta_0(t).$$

Now, we consider the BVP with impulse

$$(3.14) \quad x''(t) = J(t, x(t), x'(t); \beta_0, \beta'_0), \quad t_k < t < t_{k+1}, \quad k = 0, \dots, m,$$

$$(3.15) \quad px(0) - qx'(0) = R_1(x(\frac{1}{2}); \beta_0), \quad px(1) + qx'(1) = R_2(x(\frac{1}{2}); \beta_0),$$

and for  $k = 1, \dots, m$ ,

$$\Delta x(t_k) = u_k,$$

$$(3.16) \quad \Delta x'(t_k) = H_k(x(t_k), x'(t_k); \alpha_0, \beta_0, \beta'_0).$$

Since  $J$  and each  $H_k$  satisfy the hypotheses of Theorem 2.2, we can repeat the earlier arguments to conclude that  $\alpha_0$  and  $\beta_0$  are respectively lower and upper solutions of the BVP with impulse (3.14)-(3.16). Thus, by Theorem 2.2, there exists a solution  $\beta_1$  of the BVP with impulse (3.14)-(3.16) such that

$$(3.17) \quad \alpha_0 \leq \beta_1 \leq \beta_0.$$

Next, we show that  $\alpha_1$  and  $\beta_1$  are respectively lower and upper solutions of the BVP with impulse (1.1)-(1.3). Since  $\alpha_1$  is a solution of (3.10)-(3.12) such that

$\alpha_0 \leq \alpha_1 \leq \beta_0$ , it follows from  $(A_1)$  and (3.1) that

$$\begin{aligned} \alpha_1''(t) &= Q(t, \alpha_1(t), \alpha_1'(t); \alpha_0, \beta_0, \alpha_0') \\ &= f(t, \alpha_0(t), \alpha_0'(t)) + F_x(t, \beta_0(t))(\alpha_1(t) - \alpha_0(t)) \\ &\quad - \phi_1(t, \alpha_1(t), \alpha_1'(t)) + \phi_1(t, \alpha_0(t), \alpha_0'(t)) \\ &\geq f(t, \alpha_1(t), \alpha_1'(t)) + F_x(t, \alpha_1(t))(\alpha_0(t) - \alpha_1(t)) \\ &\quad + \phi_1(t, \alpha_1(t), \alpha_1'(t)) - \phi_1(t, \alpha_0(t), \alpha_0'(t)) \\ &\quad + F_x(t, \beta_0(t))(\alpha_1(t) - \alpha_0(t)) - \phi_1(t, \alpha_1(t), \alpha_1'(t)) + \phi_1(t, \alpha_0(t), \alpha_0'(t)) \\ &= f(t, \alpha_1(t), \alpha_1'(t)) + (F_x(t, \beta_0(t)) - F_x(t, \alpha_1(t)))(\alpha_1(t) - \alpha_0(t)) \\ &\geq f(t, \alpha_1(t), \alpha_1'(t)), \quad t \in \bigcup_{k=0}^m (t_k, t_{k+1}). \end{aligned}$$

In view of non-increasing property of  $G_1'$ , we get

$$\begin{aligned} &g_1(\alpha_1(\frac{1}{2})) - (p\alpha_1(0) - q\alpha_1'(0)) \\ &= g_1(\alpha_1(\frac{1}{2})) - r_1(\alpha_1(\frac{1}{2}); \alpha_0, \beta_0) \\ &= g_1(\alpha_1(\frac{1}{2})) - g_1(\alpha_0(\frac{1}{2})) - G_1'(\beta_0(\frac{1}{2}))(\alpha_1(\frac{1}{2}) - \alpha_0(\frac{1}{2})) \\ &\quad - \psi_1(\alpha_0(\frac{1}{2})) + \psi_1(\alpha_1(\frac{1}{2})) \\ &= G_1(\alpha_1(\frac{1}{2})) - G_1(\alpha_0(\frac{1}{2})) - G_1'(\beta_0(\frac{1}{2}))(\alpha_1(\frac{1}{2}) - \alpha_0(\frac{1}{2})) \\ &= (G_1'(c_1) - G_1'(\beta_0(\frac{1}{2}))) (\alpha_1(\frac{1}{2}) - \alpha_0(\frac{1}{2})) \geq 0, \end{aligned}$$

$(\alpha_0(\frac{1}{2}) \leq c_1 \leq \alpha_1(\frac{1}{2}))$  which yields  $p\alpha_1(0) - q\alpha_1'(0) \leq g_1(\alpha_1(\frac{1}{2}))$ . Similarly, we have  $p\alpha_1(1) + q\alpha_1'(1) \leq g_2(\alpha_1(\frac{1}{2}))$ . Using (3.3) and the nondecreasing property of  $V_k'$ , we obtain

$$\begin{aligned} &\Delta\alpha_1'(t_k) \\ &= h_k(\alpha_1(t_k), \alpha_1'(t_k); \alpha_0, \beta_0, \alpha_0') \\ &= v_k(\alpha_0(t_k), \alpha_0'(t_k)) + V_k'(\beta_0(t_k))(\alpha_1(t_k) - \alpha_0(t_k)) \\ &\quad - \phi_k(\alpha_1(t_k), \alpha_1'(t_k)) + \phi_k(\alpha_0(t_k), \alpha_0'(t_k)) \\ &\geq v_k(\alpha_1(t_k), \alpha_1'(t_k)) + V_k'(\alpha_1(t_k))(\alpha_0(t_k) - \alpha_1(t_k)) \\ &\quad + \phi_k(\alpha_1(t_k), \alpha_1'(t_k)) - \phi_k(\alpha_0(t_k), \alpha_0'(t_k)) \\ &\quad + V_k'(\beta_0(t_k))(\alpha_1(t_k) - \alpha_0(t_k)) - \phi_k(\alpha_1(t_k), \alpha_1'(t_k)) + \phi_k(\alpha_0(t_k), \alpha_0'(t_k)) \\ &= v_k(\alpha_1(t_k), \alpha_1'(t_k)) + (V_k'(\beta_0(t_k)) - V_k'(\alpha_1(t_k)))(\alpha_1(t_k) - \alpha_0(t_k)) \\ &\geq v_k(\alpha_1(t_k), \alpha_1'(t_k)), \quad k = 1, 2, \dots, m. \end{aligned}$$

Hence  $\alpha_1$  is a lower solution of (3.1.1)-(3.1.3). Similarly, for  $t \in \bigcup_{k=0}^m (t_k, t_{k+1})$ , it can be shown that

$$\beta_1''(t) \leq f(t, \beta_1(t), \beta_1'(t)),$$

$$p\beta_1(0) - q\beta_1'(0) \geq g_1(\beta_1(\frac{1}{2})), \quad p\beta_1(1) + q\beta_1'(1) \geq g_2(\beta_1(\frac{1}{2})),$$

and for  $k = 1, 2, \dots, m$ ,

$$\Delta\beta_1'(t_k) \leq v_k(\beta_1(t_k), \beta_1'(t_k)).$$

Thus,  $\beta_1$  is an upper solution of (3.1.1)-(3.1.3). Hence, by Theorem 2.1, we have

$$(3.18) \quad \alpha_1 \leq \beta_1.$$

Combining (3.13), (3.17) and (3.18) yields

$$\alpha_0(t) \leq \alpha_1(t) \leq \beta_1(t) \leq \beta_0(t), \quad t \in \bigcup_{k=0}^m (t_k, t_{k+1}).$$

Let us define the following sequences of functions  $\{Q_i\}$ ,  $\{J_i\}$ ,  $\{r_{il}\}$ ,  $\{R_{il}\}$ ,  $\{h_{kl}\}$  and  $\{H_{kl}\}$  by

$$\begin{aligned} Q_i(t, x_1, x_2) &= Q(t, x_1, x_2; \alpha_i, \beta_i, \alpha_i') \\ &= f(t, \alpha_i(t), \alpha_i'(t)) + F_x(t, \beta_i(t))(x_1 - \alpha_i(t)) \\ &\quad - \phi_1(t, x_1, x_2) + \phi_1(t, \alpha_i(t), \alpha_i'(t)), \\ J_i(t, x_1, x_2) &= J(t, x_1, x_2; \beta_i, \beta_i') = f(t, \beta_i(t), \beta_i'(t)) + F_x(t, \beta_i(t))(x_1 - \beta_i(t)) \\ &\quad - \phi_1(t, x_1, x_2) + \phi_1(t, \beta_i(t), \beta_i'(t)), \\ h_{kl} &= h_k(x_1, x_2; \alpha_i, \beta_i, \alpha_i') \\ &= v_k(\alpha_i(t_k), \alpha_i'(t_k)) + V_k'(\beta_i(t_k))(x_1 - \alpha_i(t_k)) \\ &\quad - \phi_k(x_1, x_2) + \phi_k(\alpha_i(t_k), \alpha_i'(t_k)), \\ H_{kl} &= H_k(x_1, x_2; \beta_i, \beta_i') = v_k(\beta_i(t_k), \beta_i'(t_k)) + V_k'(\beta_i(t_k))(x_1 - \beta_i(t_k)) \\ &\quad - \phi_k(x_1, x_2) + \phi_k(\beta_i(t_k), \beta_i'(t_k)), \\ r_{il} &= r_i(x(\frac{1}{2}); \alpha_i, \beta_i) = g_i(\alpha_i(\frac{1}{2})) + G_i'(\beta_i(\frac{1}{2}))(x(\frac{1}{2}) - \alpha_i(\frac{1}{2})) \\ &\quad + \psi_i(\alpha_i(\frac{1}{2})) - \psi_i(x(\frac{1}{2})), \quad i = 1, 2, \\ R_{il} &= R_i(x(\frac{1}{2}); \beta_i) = g_i(\beta_i(\frac{1}{2})) + G_i'(\beta_i(\frac{1}{2}))(x(\frac{1}{2}) - \beta_i(\frac{1}{2})) \\ &\quad + \psi_i(\beta_i(\frac{1}{2})) - \psi_i(x(\frac{1}{2})), \quad i = 1, 2. \end{aligned}$$

Now, by induction, it can be proved that

$$\alpha_0 \leq \alpha_1 \leq \dots \leq \alpha_l \leq \alpha_{l+1} \leq \beta_{l+1} \leq \beta_l \leq \dots \leq \beta_1 \leq \beta_0.$$

For that, we consider the following BVPs with impulse

$$(3.19) \quad x''(t) = Q(t, x(t), x'(t); \alpha_l, \beta_l, \alpha'_l), \quad t_k < t < t_{k+1}, \quad k = 0, \dots, m,$$

$$(3.20) \quad \begin{aligned} px(0) - qx'(0) &= r_1(x(\frac{1}{2}); \alpha_l, \beta_l), \\ px(1) + qx'(1) &= r_2(x(\frac{1}{2}); \alpha_l, \beta_l), \end{aligned}$$

and for  $k = 1, \dots, m$ ,

$$(3.21) \quad \begin{aligned} \Delta x(t_k) &= u_k, \\ \Delta x'(t_k) &= h_k(x(t_k), x'(t_k); \alpha_l, \beta_l, \alpha'_l), \end{aligned}$$

and

$$(3.22) \quad x''(t) = J(t, x(t), x'(t); \beta_l, \beta'_l), \quad t_k < t < t_{k+1}, \quad k = 0, \dots, m,$$

$$(3.23) \quad \begin{aligned} px(0) - qx'(0) &= R_1(x(\frac{1}{2}); \beta_l), \\ px(1) + qx'(1) &= R_2(x(\frac{1}{2}); \beta_l), \end{aligned}$$

and for  $k = 1, \dots, m$ ,

$$(3.24) \quad \begin{aligned} \Delta x(t_k) &= u_k, \\ \Delta x'(t_k) &= H_k(x(t_k), x'(t_k); \alpha_l, \beta_l, \beta'_l). \end{aligned}$$

Assume that  $\alpha_0 \leq \alpha_l \leq \beta_l \leq \beta_0$  for  $l > 1$  and we will show that  $\alpha_0 \leq \alpha_{l+1} \leq \beta_{l+1} \leq \beta_0$ . Since  $\alpha_l$  is a solution of (3.19)-(3.21) with  $\beta_{l-1} \geq \alpha_l$ , it follows from (3.1) that

$$\begin{aligned} \alpha_l''(t) &= Q(t, \alpha_l(t), \alpha'_l(t); \alpha_{l-1}, \beta_{l-1}, \alpha'_{l-1}) \\ &= f(t, \alpha_{l-1}(t), \alpha'_{l-1}(t)) + F_x(t, \beta_{l-1}(t))(\alpha_l(t) - \alpha_{l-1}(t)) \\ &\quad + \phi_1(t, \alpha_{l-1}(t), \alpha'_{l-1}(t)) - \phi_1(t, \alpha_l(t), \alpha'_l(t)) \\ &\geq f(t, \alpha_l(t), \alpha'_l(t)) + F_x(t, \alpha_l(t))(\alpha_{l-1}(t) - \alpha_l(t)) \\ &\quad + \phi_1(t, \alpha_l(t), \alpha'_l(t)) - \phi_1(t, \alpha_{l-1}(t), \alpha'_{l-1}(t)) \\ &\quad + F_x(t, \beta_{l-1}(t))(\alpha_l(t) - \alpha_{l-1}(t)) + \phi_1(t, \alpha_{l-1}(t), \alpha'_{l-1}(t)) \\ &\quad - \phi_1(t, \alpha_l(t), \alpha'_l(t)) \\ &\geq f(t, \alpha_l(t), \alpha'_l(t)) + (F_x(t, \alpha_l(t)) - F_x(t, \alpha_l(t)))(\alpha_{l-1}(t) - \alpha_l(t)) \\ &= f(t, \alpha_l(t), \alpha'_l(t)) \\ &= Q(t, \alpha_l(t), \alpha'_l(t); \alpha_l, \beta_l, \alpha'_l). \end{aligned}$$

Using (3.2), we obtain

$$\begin{aligned}
 & p\alpha_l(0) - q\alpha'_l(0) \\
 &= r_1(\alpha_l(\tfrac{1}{2}); \alpha_{l-1}, \beta_{l-1}) \\
 &= g_1(\alpha_{l-1}(\tfrac{1}{2})) + G'_1(\beta_{l-1}(\tfrac{1}{2}))(\alpha_l(\tfrac{1}{2}) - \alpha_{l-1}(\tfrac{1}{2})) + \psi_1(\alpha_{l-1}(\tfrac{1}{2})) - \psi_1(\alpha_l(\tfrac{1}{2})) \\
 &\leq g_1(\alpha_l(\tfrac{1}{2})) + G'_1(\alpha_l(\tfrac{1}{2}))(\alpha_{l-1}(\tfrac{1}{2}) - \alpha_l(\tfrac{1}{2})) + \psi_1(\alpha_l(\tfrac{1}{2})) - \psi_1(\alpha_{l-1}(\tfrac{1}{2})) \\
 &\quad + G'_1(\beta_{l-1}(\tfrac{1}{2}))(\alpha_l(\tfrac{1}{2}) - \alpha_{l-1}(\tfrac{1}{2})) + \psi_1(\alpha_{l-1}(\tfrac{1}{2})) - \psi_1(\alpha_l(\tfrac{1}{2})) \\
 &\leq g_1(\alpha_l(\tfrac{1}{2})) + (G'_1(\alpha_l(\tfrac{1}{2})) - G'_1(\alpha_l(\tfrac{1}{2}))) (\alpha_{l-1}(\tfrac{1}{2}) - \alpha_l(\tfrac{1}{2})) \\
 &= g_1(\alpha_l(\tfrac{1}{2})) = r_1(\alpha_l(\tfrac{1}{2}); \alpha_l, \beta_l).
 \end{aligned}$$

Similarly, it can be shown that

$$p\alpha_l(1) + q\alpha'_l(1) \leq r_2(\alpha_l(\tfrac{1}{2}); \alpha_l, \beta_l).$$

Using (3.3), the nondecreasing property of  $V'_k$  and the fact that  $\beta_{l-1} \geq \alpha_l$ , we obtain

$$\begin{aligned}
 \Delta\alpha'_l(t_k) &= h_k(\alpha_l(t_k), \alpha'_l(t_k); \alpha_{l-1}, \beta_{l-1}, \alpha'_{l-1}) \\
 &= v_k(\alpha_{l-1}(t_k), \alpha'_{l-1}(t_k)) + V'_k(\beta_{l-1}(t_k))(\alpha_l(t_k) - \alpha_{l-1}(t_k)) \\
 &\quad - \phi_k(\alpha_l(t_k), \alpha'_l(t_k)) + \phi_k(\alpha_{l-1}(t_k), \alpha'_{l-1}(t_k)) \\
 &\geq v_k(\alpha_l(t_k), \alpha'_l(t_k)) + V'_k(\alpha_l(t_k))(\alpha_{l-1}(t_k) - \alpha_l(t_k)) \\
 &\quad - \phi_k(\alpha_{l-1}(t_k), \alpha'_{l-1}(t_k)) + \phi_k(\alpha_l(t_k), \alpha'_l(t_k)) \\
 &\quad + V'_k(\beta_{l-1}(t_k))(\alpha_l(t_k) - \alpha_{l-1}(t_k)) - \phi_k(\alpha_l(t_k), \alpha'_l(t_k)) \\
 &\quad + \phi_k(\alpha_{l-1}(t_k), \alpha'_{l-1}(t_k)) \\
 &= v_k(\alpha_l(t_k), \alpha'_l(t_k)) + (V'_k(\beta_{l-1}(t_k)) - V'_k(\alpha_l(t_k)))(\alpha_l(t_k) - \alpha_{l-1}(t_k)) \\
 &\geq v_k(\alpha_l(t_k), \alpha'_l(t_k)) = h_k(\alpha_l(t_k), \alpha'_l(t_k); \alpha_l, \beta_l, \alpha'_l), \quad k = 1, \dots, m.
 \end{aligned}$$

Thus,  $\alpha_l$  is a lower solution of (3.19)-(3.21). Similarly, we can show that  $\beta_l$  is an upper solution of (3.19)-(3.21). Hence, by Theorem 2.2, there exists a solution  $\alpha_{l+1}$  of (3.19)-(3.21) satisfying

$$\alpha_0 \leq \dots \leq \alpha_l \leq \alpha_{l+1} \leq \beta_l \leq \dots \leq \beta_0.$$

In the same manner, we can show that

$$\alpha_0 \leq \dots \leq \alpha_l \leq \beta_{l+1} \leq \beta_l \leq \dots \leq \beta_0,$$

where  $\beta_{l+1}$  is a solution of (3.22)-(3.24) and  $\alpha_l, \beta_l$  are lower and upper solutions of (3.22)-(3.24). Making use of the earlier arguments, it can be shown that  $\alpha_{l+1}$  and  $\beta_{l+1}$  are respectively lower and upper solutions of the BVP with impulse

(1.1)-(1.3) and by Theorem 2.1, it follows that  $\alpha_{l+1} \leq \beta_{l+1}$ . Hence, we arrive at the conclusion

$$\alpha_0 \leq \alpha_1 \leq \dots \leq \alpha_l \leq \alpha_{l+1} \leq \beta_{l+1} \leq \beta_l \leq \dots \leq \beta_1 \leq \beta_0.$$

We now apply Kamke convergence criterion to show that each of the two sequences  $\{\alpha_l\}$  and  $\{\beta_l\}$  converges in  $B$  to  $x$ , the unique solution of the BVP with impulse (1.1)-(1.3).

The Kamke Convergence Theorem does not apply directly to either the sequence  $\{\alpha_l\}$  or  $\{\beta_l\}$  as neither  $Q_l$  nor  $J_l$  converges uniformly on compact sets to  $f$ . In order to check this, we note that

$$Q_l(t, x_1, x_2) = f(t, x_1, x_2) + F_x(t, \beta_l(t))(x_1 - \alpha_l(t)) + F(t, \alpha_l(t)) - F(t, x_1),$$

and

$$J_l(t, x_1, x_2) = f(t, x_1, x_2) + F_x(t, \beta_l(t))(x_1 - \beta_l(t)) + F(t, \beta_l(t)) - F(t, x_1).$$

Define

$$\widehat{Q}_l(t, x_1, x_2) = f(t, x_1, x_2) + F_x(t, \beta_l(t))(\alpha_{l+1}(t) - \alpha_l(t)) + F(t, \alpha_l(t)) - F(t, \alpha_{l+1}(t)),$$

$$\widehat{J}_l(t, x_1, x_2) = f(t, x_1, x_2) + F_x(t, \beta_l(t))(\beta_{l+1}(t) - \beta_l(t)) + F(t, \beta_l(t)) - F(t, \beta_{l+1}(t)),$$

so that  $\alpha_{l+1}$  is the unique solution of the BVP with impulse (1.1)-(1.3) with  $f = \widehat{Q}_l$  and each  $v_k = h_{kl}$ ,  $g_i = r_{il}$ ,  $i = 1, 2$ . The Kamke Convergence Theorem now does apply. Employing the arguments used in the proof of Theorem 2.2, we conclude that the sequence  $\{\alpha_l\}$  converges in  $B$  to  $x$ , the unique solution of (1.1)-(1.3). Similarly, the sequence  $\{\beta_l\}$  converges in  $B$  to the unique solution  $x$  of the BVP with impulse (1.1)-(1.3).

Now, we show the quadratic convergence of the sequences. For that, let us define  $\gamma_n(t) = \beta_n(t) - x(t)$ ,  $\chi_n(t) = x(t) - \alpha_n(t)$  and set  $e_n = \max\{\|\gamma_n\|_B, \|\chi_n\|_B\}$ . We will only prove the quadratic convergence of the sequence  $\{\gamma_n\}$  as that of  $\{\chi_n\}$  is similar. Using  $(A_2)$ , we find that

$$\begin{aligned} p\gamma_{n+1}(0) - q\gamma'_{n+1}(0) &\leq g_1(\beta_{n+1}(\frac{1}{2})) - g_1(x(\frac{1}{2})) \\ &= g'_1(\eta_1(\frac{1}{2}))\gamma_{n+1}(\frac{1}{2}) \leq \gamma_{n+1}(\frac{1}{2}), \end{aligned}$$

$(x(\frac{1}{2}) \leq \eta_1(\frac{1}{2}) \leq \beta_{n+1}(\frac{1}{2}))$  which can alternatively be written as

$$\gamma'_{n+1}(0) \geq \frac{p}{q}\gamma_{n+1}(0) - \frac{1}{q}\gamma_{n+1}(\frac{1}{2}) \geq 0 \text{ if } p \geq \frac{\gamma_{n+1}(\frac{1}{2})}{\gamma_{n+1}(0)}.$$

Also, for  $x(\frac{1}{2}) \leq \eta_2(\frac{1}{2}) \leq \beta_{n+1}(\frac{1}{2})$ , we have

$$\begin{aligned} p\gamma_{n+1}(1) + q\gamma'_{n+1}(1) &\leq g_2(\beta_{n+1}(\frac{1}{2})) - g_2(x(\frac{1}{2})) \\ &= g'_2(\eta_2(\frac{1}{2}))\gamma_{n+1}(\frac{1}{2}) \leq \gamma_{n+1}(\frac{1}{2}), \end{aligned}$$

which implies that

$$\gamma'_{n+1}(1) \leq \frac{1}{q}\gamma_{n+1}\left(\frac{1}{2}\right) - \frac{p}{q}\gamma_{n+1}(1) \leq 0 \text{ if } p \geq \frac{\gamma_{n+1}\left(\frac{1}{2}\right)}{\gamma_{n+1}(1)}.$$

Hence we conclude that  $\gamma'_{n+1}(0) \geq 0$  and  $\gamma'_{n+1}(1) \leq 0$  when  $p = \max\left\{\frac{\gamma_{n+1}\left(\frac{1}{2}\right)}{\gamma_{n+1}(0)}, \frac{\gamma_{n+1}\left(\frac{1}{2}\right)}{\gamma_{n+1}(1)}\right\}$ . For  $t \in \bigcup_{k=0}^m (t_k, t_{k+1})$ , the repeated application of the mean value theorem yields

$$\begin{aligned} \gamma''_{n+1}(t) &= \beta''_{n+1}(t) - x''(t) \\ &= J(t, \beta_{n+1}(t), \beta'_{n+1}(t); \beta_n, \beta'_n) - f(t, x(t), x'(t)) \\ &= f(t, \beta_n(t), \beta'_n(t)) + F_x(t, \beta_n(t))(\beta_{n+1}(t) - \beta_n(t)) \\ &\quad - \phi_1(\beta_{n+1}(t), \beta'_{n+1}(t)) + \phi_1(t, \beta_n(t), \beta'_n(t)) - F(t, x(t)) \\ &\quad + \phi_1(t, x(t), x'(t)) \\ &= F(t, \beta_n(t)) - \phi_1(t, \beta_n(t), \beta'_n(t)) + F_x(t, \beta_n(t))(\beta_{n+1}(t) - \beta_n(t)) \\ &\quad - \phi_1(t, \beta_{n+1}(t), \beta'_{n+1}(t)) + \phi_1(t, \beta_n(t), \beta'_n(t)) - F(t, x(t)) \\ &\quad + \phi_1(t, x(t), x'(t)) \\ &= F(t, \beta_n(t)) - F(t, x(t)) + F_x(t, \beta_n(t))(\beta_{n+1}(t) - \beta_n(t)) \\ &\quad + \phi_1(t, x(t), x'(t)) - \phi_1(t, \beta_{n+1}(t), \beta'_{n+1}(t)) \\ &= F_x(t, \xi_1(t))\gamma_n(t) - F_x(t, \beta_n(t))\gamma_n(t) + F_x(t, \beta_n(t))\gamma_{n+1}(t) \\ &\quad - \phi_{1x}(t, \xi_2(t), \xi_3(t))\gamma_{n+1}(t) - \phi_{1x'}(t, \xi_2(t), \xi_3(t))\gamma'_{n+1}(t), \end{aligned}$$

where  $x(t) \leq \xi_1(t) \leq \beta_n(t)$ ,  $x(t) \leq \xi_2(t) \leq \beta_{n+1}(t)$  and  $x'(t) \leq \xi_3 \leq \beta'_{n+1}(t)$ . Thus, there exists  $\xi_1(t) \leq \xi_4(t) \leq \beta_n(t)$  such that

$$\begin{aligned} \gamma''_{n+1}(t) &= F_{xx}(t, \xi_4(t))\gamma_n(t)(\xi_1(t) - \beta_n(t)) \\ &\quad + (F_x(t, \beta_n(t)) - \phi_{1x}(t, \xi_2(t), \xi_3(t)))\gamma_{n+1}(t) \\ &\quad - \phi_{1x'}(t, \xi_2(t), \xi_3(t))\gamma'_{n+1}(t) \\ &\geq -F_{xx}(t, \xi_4(t))\gamma_n^2(t) + f_{x'}(t, \xi_2(t), \xi_3(t))\gamma'_{n+1}(t), \end{aligned}$$

where we have used the monotonicity of  $F_x$ . In particular, there exists  $M > 0$  such that

$$(3.25) \quad \gamma''_{n+1}(t) - f_{x'}(t, \xi_2(t), \xi_3(t))\gamma'_{n+1}(t) \geq -Me_n^2,$$

where  $M > \max_i \max_{(t,x) \in D_i} F_{xx}(t, x)$ , and for  $i = 0, \dots, m$ ,

$$D_i = \{(t, x) : t_i \leq t \leq t_{i+1}, \alpha_0(t) \leq x \leq \beta_0(t)\}.$$

Similarly, there exist appropriate  $\xi_5$  and  $\xi_6$  such that

$$(3.26) \quad \Delta\gamma'_{n+1}(t_k) - v_{kx'}(\xi_5(t_k), \xi_6(t_k))\gamma'_{n+1}(t_k) \geq -Me_n^2, \quad k = 1, \dots, m.$$



Multiplying both sides of (3.25) by  $\mu(t)$ , we get

$$(3.27) \quad \mu(t)\gamma''_{n+1}(t) - \mu(t)f_{x'}(t, \xi_2(t), \xi_3(t))\gamma'_{n+1}(t) = (\gamma'_{n+1}(t)\mu(t))' \geq -M\mu(t)e_n^2,$$

where  $\mu(t)$  denote the integrating factor associated with (3.25) and is given by

$$\mu(t) = \exp\left(-\int_0^t f_{x'}(s, \xi_2(s), \xi_3(s))ds\right).$$

Integrating both sides of (3.27) from  $t$  to 1, we get

$$\gamma'_{n+1}(1)\mu(1) - \gamma'_{n+1}(t)\mu(t) \geq -Me_n^2 \int_t^1 \mu(s)ds, \quad t_m \leq t \leq 1.$$

In view of  $\gamma'_{n+1}(1) \leq 0$ , the above expression reduces to

$$\gamma'_{n+1}(t) \leq Me_n^2 \int_t^1 \frac{\mu(s)}{\mu(t)} ds.$$

Since  $\gamma_{n+1}$  converges to 0 in  $B$ , it ultimately follows that  $(s, \zeta_2(s), \zeta_3(s))$  belongs to

$$\widehat{D} = \{(s, x_1, x_2) : t_m \leq s \leq 1, \alpha_0(s) \leq x_1 \leq \beta_0(s), x'(s) - 1 \leq x_2 \leq x'(s) + 1\}.$$

Thus, we can bound  $\mu(t)$  away from both 0 and  $\infty$  for sufficiently large  $n$ . In particular, for  $t_m \leq t \leq 1$ , there exists  $N_1 > 0$  such that

$$(3.28) \quad \gamma'_{n+1}(t) \leq N_1 e_n^2.$$

Applying (3.26) at  $t_m$ , we obtain

$$\begin{aligned} & \Delta\gamma'_{n+1}(t_m) - v_{mx'}(\xi_5(t_m), \xi_6(t_m))\gamma'_{n+1}(t_m) \\ &= \gamma'_{n+1}(t_m^+) - \gamma'_{n+1}(t_m) - v_{mx'}(\xi_5(t_m), \xi_6(t_m))\gamma'_{n+1}(t_m) \geq -Me_n^2. \end{aligned}$$

In view of  $(A_3)$  and taking  $M_1 > 0$  as the bound for  $v_{mx'}$  away from both 0 and  $\infty$ , we get

$$(3.29) \quad \gamma'_{n+1}(t_m^-) \geq -M_1 e_n^2.$$

Using (3.27) and (3.29), there exists some  $N_2 > 0$  such that  $\gamma'_{n+1}(t) \leq N_2 e_n^2$  for  $t_{m-1} \leq t \leq t_m$ . Arguing as before, we can find a suitable form of (3.29) at  $t_{m-1}$ . Continuing this procedure inductively, we can find  $N > 0$  for sufficiently large  $n$  such that

$$(3.30) \quad \gamma'_{n+1}(t) \leq Ne_n^2, \quad t \in \bigcup_{k=0}^m [t_k, t_{k+1}].$$

Since  $\gamma_{n+1}(t) \geq 0$  and  $\gamma_{n+1} \in C[0, 1]$ , we integrate (3.30) from 0 to  $t$  and for sufficiently large  $n$ , we have  $\gamma_{n+1}(t) - \gamma_{n+1}(0) \leq Ne_n^2$ . As  $\gamma_{n+1}(0) \geq 0$ , therefore

$$(3.31) \quad 0 \leq \gamma_{n+1}(t) \leq Ne_n^2.$$

Integrating (3.27) from 0 to  $t$  ( $\leq t_1$ ), we obtain

$$\gamma'_{n+1}(t)\mu(t) - \gamma'_{n+1}(0) \geq -Me_n^2 \int_0^t \mu(s)ds.$$

As  $\gamma'_{n+1}(0) \geq 0$ , it follows that for  $0 \leq t \leq t_1$ , there exists  $N_1 > 0$  such that

$$\gamma'_{n+1}(t) \geq -Me_n^2 \int_0^t \frac{\mu(s)}{\mu(t)} ds \geq -N_1 e_n^2$$

for sufficiently large  $n$  which is analogous to (3.28). In a similar manner, we can choose  $N$  large enough such that for  $t \in \bigcup_{k=0}^m [t_k, t_{k+1}]$  and for sufficiently large  $n$ , we have

$$(3.32) \quad \gamma'_{n+1}(t) \geq -N_1 e_n^2.$$

From the estimates (3.30), (3.31) and (3.32), it follows that the sequence  $\{\beta_n\}$  converges to  $x$  quadratically in  $B$ . This completes the proof.  $\square$

#### 4. Concluding remarks

We have developed the generalized quasilinearization method for a class of impulsive three-point boundary value problems with nonlinear boundary conditions. It has been shown that the concavity assumption on the nonlinearities in the boundary conditions can be relaxed on the pattern similar to the one employed to relax the convexity condition on nonlinearity involved in the differential equation. Several interesting results appear as a special case of the work established in this paper, for example, if we take  $p = 1$ ,  $q = 0$ ,  $g_1(x(\frac{1}{2})) = a$  and  $g_2(x(\frac{1}{2})) = b$ , the results of [17] appear as a special case of our problem. In case we take  $g_1(x(\frac{1}{2})) = a$  and  $g_2(x(\frac{1}{2})) = b$ , our results correspond to an impulsive two-point boundary value problem with separated boundary conditions. Moreover, the classical method of quasilinearization for impulsive nonlinear three-point boundary value problems can be recorded by taking  $\phi(t, x) \equiv 0$  and  $\psi_i(x) \equiv 0$  in  $(A_1)$  and  $(A_2)$  of Theorem 3.1.

**Acknowledgment.** The authors are very grateful to the referee for his/her valuable suggestions that led to the improvement of the original manuscript.

#### References

- [1] A. R. Abd-Ellateef Kamar and Z. Drici, *Generalized quasilinearization for systems of nonlinear differential equations with periodic boundary conditions*, Dyn. Contin. Discrete Impuls. Syst. Ser. A Math. Anal. **12** (2005), no. 1, 77–85.
- [2] B. Ahmad, *A quasilinearization method for a class of integro-differential equations with mixed nonlinearities*, Nonlinear Anal. Real World Appl. **7** (2006), no. 5, 997–1004.
- [3] B. Ahmad and J. J. Nieto, *Existence and approximation of solutions for a class of nonlinear impulsive functional differential equations with anti-periodic boundary conditions*, Nonlinear Analysis doi:10.1016/j.na.2007.09.018.

- [4] B. Ahmad and S. Sivasundaram, *Rapid unilateral monotone scheme for impulsive hybrid nonlinear boundary value problems*, Int. J. Hybrid Systems **4** (2004), 345–354.
- [5] C. Altafini, A. Speranzon, and K. H. Johansson, *Hybrid control of a truck and trailer vehicle*, In C. J. Tomlin and M. R. Greenstreet (Eds), Hybrid Systems: Computation and Control, volume 2289, Lecture Notes in Computer Science. Springer Verlag, New York, 2002.
- [6] R. Alur, C. Courcoubetis, and D. Dill, *Model-checking for real-time systems*, Fifth Annual IEEE Symposium on Logic in Computer Science (Philadelphia, PA, 1990), 414–425, IEEE Comput. Soc. Press, Los Alamitos, CA, 1990.
- [7] P. J. Antsaklis and A. Nerode, *Hybrid Control Systems: An Introductory Discussion to the Special Issue*, IEEE Trans. Automat. Control **43** (1998), no. 4, 457–460.
- [8] C. Bai and D. Yang, *Existence of solutions for second-order nonlinear impulsive with periodic boundary value conditions*, Bound. Value Probl. Vol. 2007, Article ID 41589, 13 pages, doi:10.1155/2007/41589.
- [9] A. Balluchi, L. Benvenuti, M. Di Benedetto, C. Pinello, and A. Sangiovanni-Vincentelli, *Automotive engine control and hybrid systems: Challenges and opportunities*, Proceedings of the IEEE **7** (2000), 888–912.
- [10] A. Balluchi, P. Soueres, and A. Bicchi, *Hybrid feedback control for path tracking by a bounded-curvature vehicle*, In M. Di Benedetto and A. L. Sangiovanni-Vincentelli (Eds), Hybrid Systems: Computation and Control, volume 2034, Lecture Notes in Computer Science. Springer Verlag, New York, 2001.
- [11] R. Bellman, *Methods of Nonlinear Analysis*, Vol. 2. Academic Press, New York, 1973.
- [12] R. Bellman and R. Kalaba, *Quasilinearization and Nonlinear Boundary Value Problems*, Amer. Elsevier, New York, 1965.
- [13] M. Benchohra, J. Henderson, and S. Ntouyas, *Impulsive Differential Equations and Inclusions*, Contemporary Mathematics and Its Applications, 2. Hindawi Publishing Corporation, New York, 2006.
- [14] J. Chen, C. C. Tisdell, and R. Yuan, *On the solvability of periodic boundary value problems with impulse*, J. Math. Anal. Appl. **331** (2007), no. 2, 902–912.
- [15] M. Choisy, J. F. Guegan, and P. Rohani, *Dynamics of infectious diseases and pulse vaccination: teasing apart the embedded resonance effects*, Phys. D **223** (2006), no. 1, 26–35.
- [16] W. Ding, M. Han, and J. Mi, *Periodic boundary value problem for the second-order impulsive functional differential equations*, Comput. Math. Appl. **50** (2005), no. 3–4, 491–507.
- [17] V. Doddaballapur, P. W. Elloe, and Y. Zhang, *Quadratic Convergence of approximate solutions of two-point boundary value problems with impulse*, Proceedings of the Third Mississippi State Conference on Difference Equations and Computational Simulations (Mississippi State, MS, 1997), 81–95 (electronic), Electron. J. Differ. Equ. Conf., 1, Southwest Texas State Univ., San Marcos, TX, 1998.
- [18] A. d’Onofrio, *On pulse vaccination strategy in the SIR epidemic model with vertical transmission*, Appl. Math. Lett. **18** (2005), no. 7, 729–732.
- [19] P. Egbunonu and M. Guay, *Identification of switched linear systems using subspace and integer programming techniques*, Nonlinear Anal. Hybrid Syst. **1** (2007), no. 4, 577–592.
- [20] P. W. Elloe and B. Ahmad, *Positive solutions of a nonlinear nth order boundary value problem with nonlocal conditions*, Appl. Math. Lett. **18** (2005), no. 5, 521–527.
- [21] P. W. Elloe and S. G. Hristova, *Method of the quasilinearization for nonlinear impulsive differential equations with linear boundary conditions*, Electron. J. Qual. Theory Differ. Equ. **2002** (2002), no. 10, 1–14.
- [22] P. Elloe and Y. Gao, *The method of quasilinearization and a three-point boundary value problem*, J. Korean Math. Soc. **39** (2002), no. 2, 319–330.

- [23] P. W. Eloe and Y. Zhang, *A quadratic monotone iteration scheme for two-point boundary value problems for ordinary differential equations*, *Nonlinear Anal.* **33** (1998), no. 5, 443–453.
- [24] S. Engell, G. Frehse, and E. Schnieder, *Lecture Notes In Control and Information Sciences; Modelling, Analysis and Design of Hybrid Systems*, Springer Verlag, Heidelberg, 2002.
- [25] S. Engell, S. Kowalewski, C. Schultz, and O. Strusberg, *Continuous-discrete interactions in chemical process plants*, *Proceedings of the IEEE* **7** (2000), 1050–1068.
- [26] S. Gao, L. Chen, J. J. Nieto, and A. Torres, *Analysis of a delayed epidemic model with pulse vaccination and saturation incidence*, *Vaccine* **24** (2006), 6037–6045.
- [27] C. P. Gupta, *A second order m-point boundary value problem at resonance*, *Nonlinear Anal.* **24** (1995), no. 10, 1483–1489.
- [28] C. P. Gupta and S. Trofimchuck, *A priori estimates for the existence of a solution for a multi-point boundary value problem*, *J. Inequal. Appl.* **5** (2000), no. 4, 351–365.
- [29] I. T. Kiguradze and A. G. Lomtatidze, *On certain boundary value problems for second-order linear ordinary differential equations with singularities*, *J. Math. Anal. Appl.* **101** (1984), no. 2, 325–347.
- [30] B. Krogh and N. Lynch, *Hybrid Systems: Computation and Control*, *Lecture Notes in Computer Science*, Vol. 1790, Springer Verlag, New York, 2000.
- [31] V. Lakshmikantham, *An extension of the method of quasilinearization*, *J. Optim. Theory Appl.* **82** (1994), no. 2, 315–321.
- [32] ———, *Further improvement of generalized quasilinearization method*, *Nonlinear Anal.* **27** (1996), no. 2, 223–227.
- [33] V. Lakshmikantham, D. D. Bainov, and P. S. Simeonov, *Theory of Impulsive Differential Equations*, World Scientific, Singapore, 1989.
- [34] V. Lakshmikantham and A. S. Vatsala, *Generalized Quasilinearization for Nonlinear Problems*, Kluwer Academic Publishers, Dordrecht, 1998.
- [35] ———, *Generalized quasilinearization versus Newton's method*, *Appl. Math. Comput.* **164** (2005), no. 2, 523–530.
- [36] X. Liu, *Advances in Impulsive Differential Equations*, *Dyn. Contin. Discrete Impuls. Syst. Ser. A Math. Anal.* **9** (2002), no. 3, 313–462.
- [37] J. Lygeros, D. N. Godbole, and S. Sastry, *A verified hybrid controller for automated vehicles*, *IEEE Trans. Automat. Control* **43** (1998), no. 4, 522–539.
- [38] V. B. Mandelzweig and F. Tabakin, *Quasilinearization approach to nonlinear problems in physics with application to nonlinear ODEs*, *Comput. Phys. Comm.* **141** (2001), no. 2, 268–281.
- [39] D. L. Peypne and C. G. Cassandras, *Optimal control of hybrid systems in manufacturing*, *Proceedings of the IEEE* **7** (2000), 1108–1123.
- [40] Y. V. Rogovchenko, *Impulsive evolution systems: main results and new trends*, *Dynam. Contin. Discrete Impuls. Systems* **3** (1997), no. 1, 57–88.
- [41] A. M. Samoilenko and N. A. Perestyuk, *Impulsive Differential Equations*, World Scientific Publishing Co., Inc., River Edge, NJ, 1995.
- [42] S. Tang and L. Chen, *Density-dependent birth rate, birth pulses and their population dynamic consequences*, *J. Math. Biol.* **44** (2002), no. 2, 185–199.
- [43] C. Tomlin, G. Pappas, and S. Sastry, *Conflict resolution for air traffic management: a study in multiagent hybrid systems*, *IEEE Trans. Automat. Control* **43** (1998), no. 4, 509–521.
- [44] F. Vaandrager and J. Van Schuppen, *Hybrid Systems: Computation and Control*, *Lecture Notes in Computer Science*, Vol. 1569, Springer Verlag, New York, 1999.
- [45] P. Varaiya, *Smart cars on smart roads: problems of control*, *IEEE Trans. Automat. Control* **38** (1993), no. 2, 195–207.

- [46] I. Yermachenko and F. Sadyrbaev, *Quasilinearization and multiple solutions of the Emden-Fowler type equation*, Math. Model. Anal. **10** (2005), no. 1, 41–50.
- [47] S. T. Zavalishchin and A. N. Sesekin, *Dynamic Impulse Systems*, Theory and applications. Mathematics and its Applications, 394. Kluwer Academic Publishers Group, Dordrecht, 1997.

BASHIR AHMAD  
DEPARTMENT OF MATHEMATICS  
FACULTY OF SCIENCE  
KING ABDULAZIZ UNIVERSITY  
P.O. Box. 80203  
JEDDAH 21589, SAUDI ARABI  
*E-mail address:* bashir\_qau@yahoo.com

AHMED ALSAEDI  
DEPARTMENT OF MATHEMATICS  
FACULTY OF SCIENCE  
KING ABDULAZIZ UNIVERSITY  
P.O. Box. 80203  
JEDDAH 21589, SAUDI ARABI  
*E-mail address:* aalsaedi@hotmail.com

DOA'A GAROUT  
DEPARTMENT OF MATHEMATICS  
FACULTY OF SCIENCE  
KING ABDULAZIZ UNIVERSITY  
P.O. Box. 80203  
JEDDAH 21589, SAUDI ARABI  
*E-mail address:* doaa\_ag@hotmail.com