

Experimental and Theoretical Studies on a Thiocarbamide Derivative Containing Schiff Base Groups

Pu Su Zhao,^{*} De Long Shao, Jing Zhang, Yi Wei,[†] and Fang Fang Jian[‡]

Jiangsu Key Laboratory for Chemistry of Low-Dimensional Materials, Huaivn Normal University, Huaian, Jiangsu, 223300, P. R. China. *E-mail: zhaopusu@163.com

[†]New Materials & Function Coordination Chemistry Laboratory, Qingdao University of Science and Technology, Qingdao Shandong 266042, P. R. China

Received February 20, 2009. Accepted May 28, 2009

Key Words: Synthesis, Crystal structure, Density functional calculations, Electronic absorption spectra, Thermodynamic property

Thiocarbamide group is a key element with common structure in a variety of natural and synthetic compounds with interesting biological or chemical properties, and therefore has been known for its important medicinal,¹⁻³ bioorganic,⁴ supramolecular chemistry⁵ applications. Most recently, thiocarbamide derivatives have been used for asymmetric synthesis⁶ and they play an important role as chiral catalysts for highly enantioselective michael reactions⁷ and a new type of herbicides for weed control.⁸ Sparked by above reasons, our group persists in synthesizing and making studies on thiocarbamide derivatives.⁹⁻¹³ On the other hand, Schiff base ligand possess N-donor with favorable coordination ability and easily constructs hydrogen bonds in supramolecular chemistry.¹⁴ In addition, Schiff base complexes have been extensively studied in great detail as a result of their prospective applications in catalysis, magnetic properties, molecular architectures and materials chemistry by coordination chemists at all times.¹⁵⁻¹⁸ Herein, we report the experimental results about the title compound, 1,5-bis[1-(4-nitrophenyl)ethylidene]thiocarbonohydrazide, which is a typical thiocarbamide derivative containing Schiff base groups. We also report the theoretical calculational results about the title compound. Since with the recent advances in compute hardware and software, it is possible to correctly describe physico-chemical properties of relatively small molecules accuracy¹⁹⁻²² using density functional theory (DFT).²³⁻²⁵ The comparisons between the experiments and the calculations about the title compound have also been revealed in the paper.

Crystal structure of the title compound crystallizes in triclinic system, space group *P*-1, with lattice parameters $a = 7.858(4)$ Å, $b = 16.170(7)$ Å, $c = 16.842(7)$ Å, $\alpha = 102.540(9)^\circ$, $\beta = 100.005(8)^\circ$, $\gamma = 101.432(8)^\circ$, $V = 1994.3(16)$ Å³, $M_r = 836.87$ (C₃₄H₃₆N₁₂O₁₀S₂), $Z = 2$, $D_c = 1.394$ g/cm³, $\mu = 0.204$ mm⁻¹, $F(000) = 872$, GOF = 0.926, $R_1 = 0.1139$, $wR_2 = 0.2387$. A displacement ellipsoid plot with the atomic numbering scheme is shown in Figure 1.

The structure of the title compound contains two crystallographically independent water molecules and two 1,5-bis[1-(4-nitrophenyl)ethylidene] thiocarbonohydrazide molecules, hereafter named T1 [containing C(1) ~ C(6) phenyl ring] and T2 [containing C(18) ~ C(23) phenyl ring]. In T1 and T2, all

of the bond lengths and bond angles are different. For example, S=C bond length of 1.682(12) Å in T1 is longer than that of 1.658(11) Å in T2 and both of them are shorter than that found in the similar structure [1.704(3)].¹⁰ In T1 and T2, despite some differences, all of the bond lengths and bond angles in the phenyl rings are in the normal range. As for the N-C single bonds connected with the N-N bonds, in T1, the two bond lengths of N-C are 1.332(12) Å and 1.364(12) Å, respectively, while in T2, the two N-C bond lengths are 1.329(12) Å and 1.345(11) Å, respectively. These four N-C bond lengths are corresponding to those found in a similar structure of 1,5-bis(9-anthracylidene)thiocarbonohydrazide,²⁶ where the two N-C bond lengths are 1.332(2) and 1.355(3) Å, respectively. In view of the bond lengths of N-N and C=N in T1 and T2 (see supporting information S1), all of which are also in agreement with those in the above cited structure [N-N 1.380(2) and 1.381(2) Å, C=N 1.250(3) and 1.266(2) Å].²⁶ Four phenyl rings in T1 and T2 are almost planar each other. The dihedral angles between the two phenyl rings in T1 is 15.34(2)° and in T2 is 10.49(2)°. The dihedral angles between the phenyl ring of C(1) ~ C(6) in T1 and the two phenyl rings in T2 are 16.65(2) and 6.42(2)°, respectively. The phenyl ring C(12) ~ C(17) in T1 forms the dihedral angles with the two phenyl rings in T2 being 7.29 (2) and 9.21(2)°, respectively.

In the crystal lattice, there are some intermolecular and intramolecular supramolecular interactions and the two water molecules take part in the formation of these supramolecular interactions (see supporting information S2), which stabilize




Figure 1. The molecular structure with the atomic numbering for the title compound.

the crystal structure in the solid state.

One molecular structure of the title compound was selected to be optimized at DFT-B3LYP/6-31G* level of theory. Comparisons between the theoretical values with the experimental ones indicate that most of the optimized bond lengths are slightly larger than the experimental values, as the theoretical calculations are performed for isolated a molecule in gaseous phase and the experimental results are for a molecule in a solid state. The geometry of the solid-state structure is subject to intermolecular forces, such as van der Waals interactions, crystal packing forces and hydrogen-bond forces. The biggest bond length differences between the experimental and the predicted values in T1 is 0.035 Å for N(4)-C(9) bond and in T2 is 0.057 Å for N(10)-C(26) bond. In view of the bond angle difference between experimental and theoretical values, the biggest difference in T1 is 4.7° for C(9)-N(3)-N(2) bond angle and in T2 is 3.6° for C(32)-C(33)-C(34) bond angle. The differences maybe since that, in the crystal structure, all the atoms of N(4), N(10), N(2), N(3) and C(33) take part in the formation of hydrogen bonds (see S2), while in theoretical calculations, there was none hydrogen bond considered. Despite of these differences, the optimized geometry can reproduce the crystal structure on the whole.

Based on the optimized structure of the title compound at B3LYP/6-31G* level of theory, the NPA atomic charge distributions for the non-hydrogen atoms were calculated and the results are listed in Table 1. The atomic numbering scheme is the same as that in T1 molecule.

Seen from Table 1, on the one hand, for NPA atomic charge distributions of the non-hydrogen atoms, atomic electronegativity plays important roles. Namely, between the two connecting atoms, the atom having bigger electronegativity will carry negative charges, while the atom having smaller electronegativity will carry positive charges. For example, when a carbon atom is connected with a nitrogen atom, the carbon atomic charges are positive values and the nitrogen atomic charges are negative values, since the nitrogen atom has bigger electronegativity than the carbon atom. However, when a nitrogen atom is joined with an oxygen atom, the nitrogen atom has positive charges and the oxygen atom has negative charges, since the oxygen atom has bigger electronegativity than the nitrogen atom. So, for the title compound, the atoms of C(7), C(9) and C(11) have positive atomic charges and N(2), N(3), N(4) and N(5) have negatively atomic charges.

As regards two carbon atoms joined to two nitro-groups, atoms C(1) and C(15) carry positive charges. N(1) and N(6) also carry positive charges, while four oxygen atoms carry negative charges. C(8) and C(10) in methyl groups and other carbon atoms in two phenyl rings all carry negative charges, since the hydrogen atom has smaller electronegativity than the carbon atom. On the other hand, for the non-hydrogen atoms in Table 1, although atoms of N(3), N(4), C(8) and C(1) in two methyl groups and the some carbon atoms in two phenyl rings have bigger negative charge values, steric effect hinders them to coordinate with metallic ions further. For the atoms of S(1), N(2), N(5) and four oxygen atoms in two nitro-groups, they all carry negative atomic charges and there is none steric effect hindered them, which results in these atoms being able to coordinate with metallic ions further. Namely, the title compound is a good multidentate ligand and it has the potential to construct interesting complexes with various metallic ions, which supports one of our original ideas of the synthesis.


Electronic absorption spectra were calculated by using TD-DFT method at B3LYP/6-31G* level based on the optimized geometry. The experimental electronic absorption spectra values and theoretical electronic absorption spectra information are both given in Table 2. Seen from Table 2, the experiment and theoretical calculations both obtain two absorption bands: one is narrow and another is broad. In experiments, the broad band is from 245 nm to 255 nm and the narrow peak at 349 nm, while contrary to the experiments, in calculations, the narrow peak is at 242 nm and the broad band is from 380 nm to 386 nm. In addition, compared with the experimental data, although the predicted narrow peak has some blue shifts and the predicted broad band has some red shifts, the intensity order of the two predicted bands are the same as the experiments. Namely, TD-DFT method at B3LYP/6-31G* level can predict the electron transition spectra approximately. The detailed electronic transition modes corresponding to each absorption peak are also listed in Table 2. Based on the optimized structure, molecular orbital coefficients analyses indicate that the frontier molecular orbitals are mainly composed of *pv* and *pz* atomic orbitals and electronic absorption spectra aforementioned are mainly assigned to *n-π** and *π-π** electron transitions. Figure 2 shows some frontier molecular orbitals surfaces of the title compound. Seen from Figure 2, except the two methyl groups, the other groups, such as the

Table 1. NPA atomic charge distributions for the non-hydrogen atoms

Atom	Charges(<i>e</i>)	Atom	Charges(<i>e</i>)	Atom	Charges(<i>e</i>)
S(1)	-0.14813	C(1)	0.05929	C(12)	-0.06046
N(2)	-0.29106	C(2)	-0.20297	C(13)	-0.19051
N(3)	-0.41642	C(3)	-0.21675	C(14)	-0.20808
N(4)	-0.41840	C(4)	-0.05821	C(15)	0.05901
N(5)	-0.23079	C(5)	-0.20007	C(16)	-0.20637
C(9)	0.21749	C(6)	-0.20716	C(17)	-0.22020
N(1)	0.51114	C(7)	0.23447	C(11)	0.23199
N(6)	0.51113	C(8)	-0.75325	C(10)	-0.75242
O(1)	-0.38145	O(2)	-0.38109	O(3)	-0.38213
O(4)	-0.38635				

Table 2. Experimental and theoretical electronic absorption spectra values

Exp.		Calc. at B3LYP/6-31G* level		
Wavelength (nm)	log ϵ	Wavelength (nm)	Oscillator strength	Transition
245-255	3.34	242	0.1101	HOMO-2 \rightarrow LUMO+4
349	3.48	380	0.2967	HOMO-2 \rightarrow LUMO HOMO-1 \rightarrow LUMO+1 HOMO-1 \rightarrow LUMO HOMO-2 \rightarrow LUMO+1
		386	0.4794	HOMO-2 \rightarrow LUMO HOMO \rightarrow LUMO+1

**Figure 2.** Molecular orbitals active in the electronic transitions of the title compound.**Table 3.** Thermodynamic properties of the title compound at different temperatures

T (K)	$C_{p,m}^0$ (J·mol ⁻¹ ·K ⁻¹)	S_m^0 (J·mol ⁻¹ ·K ⁻¹)	H_m^0 (kJ·mol ⁻¹)
200.0	305.42	647.23	36.80
298.1	421.09	790.90	72.50
300.0	423.20	793.51	73.28
400.0	529.63	930.22	121.06
500.0	617.83	1058.23	178.59
600.0	688.37	1177.35	244.03
700.0	744.74	1287.86	315.79
800.0	790.40	1390.38	392.63

S=C bond, two nitro-groups, two phenyl rings, two C=N bonds and two N-N bonds, all take part in the electronic transitions, which are corresponding to the intermolecular $n-\pi^*$ and $\pi-\pi^*$ electron transitions.

On the basis of statistical thermodynamic and vibrational analysis at B3LYP/6-31G* level, the standard thermodynamic functions: heat capacity ($C_{p,m}^0$), entropy (S_m^0) and enthalpy (H_m^0) were obtained and listed in Table 3. The scale factor for the frequencies is 0.96. As observed from Table 3, all the values of $C_{p,m}^0$, S_m^0 and H_m^0 increase with the increase of temperature from 200.0 to 800.0 K, which is attributed to the enhancement of the molecular vibration while the temperature increases.

The correlations linear equations between these thermodynamic properties and temperatures T are as follows:

$$C_{p,m}^0 = 20.6011 + 1.5761T - 7.694 \times 10^{-4} T^2 \quad (R^2 = 0.9999)$$

$$S_m^0 = 330.1593 + 1.6749T - 4.3757 \times 10^{-4} T^2 \quad (R^2 = 0.999999)$$

$$H_m^0 = -20.8604 + 0.1976T + 4.017 \times 10^{-4} T^2 \quad (R^2 = 0.9998)$$

These equations could be used for the further studies of the title compound.

Experimental Section and Computational Method

Synthesis. All chemicals were obtained from a commercial source and used without further purification.

p-Nitroacetophenone (0.02 mol) and thiocarbonohydrazide (0.01 mol) were mixed in ethanol solution (50 mL) and stirred with refluxing, 4 h later, many light-yellow solids were observed and then, the reaction was stopped and the mixture was cooled to room temperature. The yellow solids were obtained by filtration and dried at room temperature. Yield 89%, mp. 204-205 °C. IR: ν 3426 (m), 3109 (m), 2966 (m), 1579 (m), 1516 (vs), 1340 (vs), 1276 (m), 1224 (s), 1109 (s), 1064 (m), 875 (s), 789 (w), 753 (m), 692 (w), 609 (w), 543 (w) cm⁻¹. Found: C, 48.79; H, 4.34; N, 20.09%. Calc. for C₃₃H₂₆N₁₂O₁₀S₂: C, 48.66; H, 4.23; N, 20.18%. ¹H NMR (400 MHz, CDCl₃) δ 1.28 (s, 6H, -CH₃), 7.28 (s, 2H, -NH), 7.91-8.32 (m, 8H, -C₆H₄).

For the compound, single crystals suitable for X-ray measurements were obtained by recrystallization from acetic ether at room temperature.

Crystal structure determination. The diffraction data were collected on a Enraf-Nonius CAD-4 diffractometer with graphite-monochromated Mo-K α radiation ($\lambda = 0.71073$ Å, $T = 295(2)$ K). The technique used was ω scan with limits 1.58 to 25.00°. The structure of the title compound was solved by direct method and refined by least squares on F^2 by using the SHELXTL²⁷ software package. All non-hydrogen atoms were anisotropically refined. The approximate positions of the four water H atoms were firstly obtained from a difference map. Then, the four atoms were placed in 'ideal' positions. The other hydrogen atom positions were fixed geometrically at calculated distances and allowed to ride on the parent carbon and nitrogen atoms. The molecular graphics were plotted using SHELXTL. Atomic scattering factors and anomalous dispersion corrections were taken from International Tables for X-ray Crystallography.²⁸

Computational methods. The crystal structure of the title

compound was used as initial molecular geometry, then it was optimized by using MM+ molecular modeling and semi-empirical AM1 method²⁹ (HYPERCHEM 6.0, Hypercube, Ont., Canada). In the next step, B3LYP (Becke's three parameter hybrid functional using the LYP correlation functional) calculations at basis set 6-31G* by the Berny method³⁰ were performed with the Gaussian 03 software package.³¹ Vibrational frequencies calculated ascertain the structure was stable (no imaginary frequencies). The thermodynamic properties of the title compound at different temperatures were calculated on the basis of vibrational analyses. Electronic absorption spectra were calculated by using time-dependent density functional theory (TDDFT)^{32,34} method at 6-31G* basis set. Natural Bond Orbital (NBO) analyses were performed on the optimized structures.³⁵ All calculations were performed on a DELL PE 2850 server and a Pentium IV computer using the default convergence criteria.

Supplementary Data. Crystallographic data for the structure reported here have been deposited with Cambridge Crystallographic Data Center (Deposition No. CCDC- 719724). The data can be obtained free of charge *via* www.ccdc.cam.ac.uk/conts/retrieving.html (or from the CCDC, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +441223 336033; e-mail: deposit@ccdc.cam.ac.uk). Some selected geometric parameters for both experimental and calculational data can be obtained from the author directly.

Acknowledgments. This work was supported by Jiangsu Key Laboratory for Chemistry of Low-Dimensional Materials P. R. China (JSKC08047) and Fund of Huaiyin Normal University (07HSBS004, 08HSJSK003).

References

- Di Grandi, M. J.; Curran, K. J.; Feigelson, G.; Prasad, A.; Ross, A. A.; Visalli, R.; Fairhurst, J.; Feld, B.; Bloom, J. D. *Bioorg. Med. Chem. Lett.* **2004**, *14*, 4157.
- Han, T.; Cho, J. H.; Oh, C. H. *Eur. J. Med. Chem.* **2006**, *41*, 825.
- Kaymakcioglu, B. K.; Rollas, S.; Korcegez, E.; Aricioglu, F. *Eur. J. Pharm. Sci.* **2005**, *26*, 97.
- Rostom, S. A. F. *Bioorg. Med. Chem.* **2006**, *14*, 6475.
- Heck, R.; Marsura, A. *Tetrahedron Lett.* **2003**, *44*, 1533.
- Cao, C. L.; Ye, M. C.; Sun, X. L.; Tang, Y. *Org. Lett.* **2006**, *8*, 2901.
- Liu, K.; Cui, H. F.; Nie, J.; Dong, K. Y.; Li, X. J.; Ma, J. A. *Org. Lett.* **2007**, *9*, 923.
- Wang, X. M.; Zhu, T.; Zheng, L. Y.; Li, Y. X.; Zheng, J. F.; Zhang, S. Q.; Bai, J. *Chin. J. Org. Chem.* **2006**, *26*, 660.
- Jian, F. F.; Zhao, P. S.; Hou, Y. X. *Chin. J. Struct. Chem.* **2004**, *23*, 1256.
- Jian, F. F.; Zhao, P. S.; Hou, Y. X. *Struct. Chem.* **2005**, *16*, 361.
- Jian, F. F.; Zhao, P. S.; Bai, Z. S.; Zhang, L. *Struct. Chem.* **2005**, *16*, 635.
- Jian, F. F.; Wang, K. F.; Zhao, P. S.; Zhuang, R. R.; Zheng, J. *Struct. Chem.* **2006**, *17*, 539.
- Zhao, P. S.; Li, R. Q.; Zhang, L. L.; Jian, F. F. *Polish J. Chem.* **2006**, *80*, 2031.
- Li, C. H.; Wang, Q.; Xu, Y. Q.; Hu, C. W. *Chin. J. Struct. Chem.* **2008**, *27*, 187.
- Kasai, K.; Aoyagi, M.; Fujita, M. *J. Am. Chem. Soc.* **2000**, *122*, 2140.
- Kitagawa, S.; Kitaura, R.; Noro, S. I. *Angew. Chem., Int. Ed.* **2004**, *43*, 2334.
- Hoshino, N.; Ito, T.; Nihei, M.; Oshio, H. *Inorg. Chem. Commun.* **2003**, *6*, 377.
- Khandar, A. A.; Nejati, K. *Polyhedron* **2000**, *19*, 607.
- Foresman, J. B.; Frisch, A. *Exploring Chemistry with Electronic Structure Methods*, 2nd ed.; Gaussian, Inc.: Pittsburgh, PA, 1996.
- Ruud, K.; Helgaker, T.; Kobayashi, R.; Jorgensen, P.; Bak, K. L. *J. Chem. Phys.* **1994**, *100*, 8178.
- Hehre, W. J.; Radom, L.; Schleyer, P. R.; Pople, J. A. *Ab Initio Molecular Orbital Theory*; Wiley: New York, 1986.
- Parr, R. G.; Yang, W. T. *Density Functional Theory of Atoms and Molecules*; Oxford University Press: New York, 1989 and references therein.
- Ziegler, T. *Chem. Rev.* **1991**, *91*, 651 and references therein.
- Wolinski, K.; Hinton, J. F.; Pulay, P. *J. Am. Chem. Soc.* **1990**, *112*, 8251.
- Schreckenbach, G.; Ziegler, T. *J. Phys. Chem.* **1995**, *A 99*, 606.
- Han, F.; Bao, Y. H.; Yang, Z. G.; Fyles, T. M.; Zhao, J. Z.; Peng, X. J.; Fan, J. L.; Wu, Y. K.; Sun, S. G. *Chem. Eur. J.* **2007**, *13*, 2880.
- Sheldrick, G. M. *SHELXTL, v5 Reference Manual*; Siemens Analytical X-Ray Systems: Madison, WI, 1997.
- Wilson, A. J. *International Table for X-Ray Crystallography*; Kluwer Academic: Dordrecht, The Netherlands, 1992; Vol. C, Tables 6.1.1.4 (pp 500-502) and 4.2.6.8 (pp 219-222), respectively.
- Dewar, M. J. S.; Zoebisch, E. G.; Healy, E. F. *J. Am. Chem. Soc.* **1985**, *107*, 3902.
- Peng, C.; Ayala, P. Y.; Schlegel, H. B.; Frisch, M. J. *J. Comput. Chem.* **1996**, *17*, 49.
- Frisch, M. J.; Trucks, G. W.; Schlegel, H. B.; Scuseria, G. E.; Robb, M. A.; Cheeseman, J. R.; Montgomery, J. A.; Vreven, Jr., T.; Kudin, K. N.; Burant, J. C.; Millam, J. M.; Iyengar, S. S.; Tomasi, J.; Barone, V.; Mennucci, B.; Cossi, M.; Scalmani, G.; Rega, N.; Petersson, G. A.; Nakatsuji, H.; Hada, M.; Ehara, M.; Toyota, K.; Fukuda, R.; Hasegawa, J.; Ishida, M.; Nakajima, T.; Honda, Y.; Kitao, O.; Nakai, H.; Klene, M.; Li, X.; Knox, J. E.; Hratchian, H. P.; Cross, J. B.; Adamo, C.; Jaramillo, J.; Gomperts, R.; Stratmann, R. E.; Yazyev, O.; Austin, A. J.; Cammi, R.; Pomelli, C.; Ochterski, J. W.; Ayala, P. Y.; Morokuma, K.; Voth, G. A.; Salvador, P.; Dannenberg, J. J.; Zakrzewski, V. G.; Dapprich, S.; Daniels, A. D.; Stram, M. C.; Farkas, O.; Malick, D. K.; Rabuck, A. D.; Raghavachari, K.; Foresman, J. B.; Ortiz, J. V.; Cui, Q.; Baboul, A. G.; Clifford, S.; Cioslowski, J.; Stefanov, B. B.; Liu, G.; Liashenko, A.; Piskorz, P.; Komaromi, I.; Martin, R. L.; Fox, D. J.; Keith, T.; Al-Laham, M. A.; Peng, C. Y.; Nanayakkara, A.; Challacombe, M.; Gill, P. M. W.; Johnson, B.; Chen, W.; Wong, M. W.; Gonzalez, C.; Pople, J. A. Gaussian, Inc.: Wallingford, CT, 2004.
- Petersilka, M.; Gossmann, U. J.; Gross, E. K. U. *Phys. Rev. Lett.* **1966**, *76*, 1212.
- Bauernschmitt, R.; Ahlrichs, R. *Chem. Phys. Lett.* **1996**, *256*, 454.
- Jamorski, C.; Casida, M. E.; Salahub, D. R. *J. Chem. Phys.* **1996**, *104*, 5134.
- Runge, E.; Gross, E. K. U. *Phys. Rev. Lett.* **1984**, *52*, 997.