

Xenografted Tumorigenesis in the oral vestibule of nude mice by Snail transfection: Histological and immunohistochemical study

Moon-Key Kim^{1,5}, Eun-Ha Lee⁴, Jin Kim^{2,4}, Jong-In Yook^{2,4}, In-Ho Cha^{2,3}

¹Department of Oral and Maxillofacial Surgery, National Health Insurance Corporation Ilsan Hospital,

²Oral Cancer Research Institute, ³Department of Oral and Maxillofacial Surgery,

⁴Department of Oral Pathology, College of Dentistry, Yonsei University,

⁵Department of Dental Science, Graduate School, Yonsei University

Abstract (J. Kor. Oral Maxillofac. Surg. 2009;35:199-204)

Purpose: The purpose of this study is to investigate the epithelial-mesenchymal transition (EMT) induced by Snail transcription factor and Snail-transfected *in vivo* tumors with histopathological features.

Materials and methods: We induced *in vivo* xenografted tumorigenesis in the oral vestibules of nude mice by a Snail transfected HaCaT cell line and investigated morphological and immunohistochemical features in Snail expressive tumors.

Results: We identified tumor masses in 14 out of 15 nude mice in the HaCaT-Snail cell inoculation group, but no tumors were present in any of the HaCaT cell inoculation group. Induced tumors showed features of poorly differentiated carcinoma with invasion to neighboring muscles and bones. The HaCaT-Snail tumors showed decreased expressions of E-cadherin and cytokeratin, but showed increased expressions of vimentin and N-cadherin.

Discussion: The Snail transfected xenograft can improve productivity of malignant tumors, show various histopathological features including invasive growth, and aid in the investigation of tumor progression and the interaction with surrounding tissues.

Key words: E-cadherin, neoplastic cell transformation, Snail transcription factor, xenograft model

(원고접수일 2009.6.2. / 1차수정일 2009.6.12. / 2차수정일 2009.6.27. / 게재확정일 2009.7.13.)

Introduction

The epithelial-mesenchymal transition (EMT) has been observed during malignant transformation in some epithelial tumors¹. E-cadherin, which is located in adherens junction and mediates homophilic cell-cell adhesion, is repressed during EMT². E-cadherin is also involved in maintenance of the cytoskeleton through intracellular binding to actins via catenins³. Since the reduction of its expression has been reported to be associated with tumor invasiveness and prognosis^{4,5}, the E-cadherin gene is also called an invasion suppressor gene⁶. The E-cadherin loses its expression when an advanced epithelial tumor has its ability to invasion⁷.

Meanwhile, N-cadherin, which expressed in nerve cells and fibroblasts and is involved in heterophilic cell-cell adhesion⁸, is called an invasion promoter¹⁰ since there are some relationships between tumor progression and an increase of its expression¹¹. N-cadherin is often up-regulated in situations where E-cadherin has been repressed^{11,12}.

The Snail transcription factor, first described by Bouley et al¹³, induces EMT by down-regulation of E-cadherin. Snail has 4-6 C₂H₄ zinc finger domains and binds to CACCTG (or CAGGTG) specifically in promoter regions as a transcriptional repressor. More than 50 homologues have been reported, and Snail-transfected cells show repression of E-cadherin. Snail binds to the promoter of E-cadherin and represses its expression, resulting in loss of cell-cell adhesion and phenotypic changes, with additional effects on cell survival and migration.

The role of Snail in tumorigenesis as well as tumor invasion and metastasis has been studied in breast, gastric, and hepatic cancers. Yokoyama et al¹⁴ reported that repression of E-cadherin was observed in Snail mRNA expressive oral squamous cell carcinoma culture cells and Takkunen et al¹⁵ reported that

차인호

120-752 서울특별시 서대문구 성산로 250
연세대학교 치과대학 구강악안면외과학교실

In-Ho Cha

Department of Oral and Maxillofacial Surgery, College of Dentistry, Yonsei University
250 Seongsan-ro, Seodaemun-gu, Seoul, 120-752, Korea
Tel: 82-2-2228-3140 Fax: 82-2-364-0992
E-mail: cha8764@yumc.yonsei.ac.kr

*This work was supported by the Korea Research Foundation Grant funded by the Korean Government (KRF-2008-005-J00803).

EMT in oral squamous cell carcinoma cells could result from Snail, a E-cadherin repressor.

We developed a Snail transfected cell line derived from HaCaT cell line, which is a spontaneously immortalized keratinocyte cell line derived from normal adult human skin. It was called HaCaT because it had developed from human adult skin ketatinocytes during prolonged cultivation at a reduced Ca_2+ concentration and elevated temperature¹⁶⁾. We used this cell line for a xenografted tumor model in nude mice to investigate morphological and immunohistochemical changes in Snail expressive tumors.

Materials and methods

1. Plasmid

A human Snail cDNA was subcloned into the pBabePuro retroviral vector to generate pBabePuro-Snail-FLAG plasmid (Fig. 1). The amphotropic Phoenix cell line was cultured in Dulbecco's modified Eagle's medium (DMEM; Gibco BRL, Grand island, NY) supplemented with 1 mM sodium pyruvate, 100 units/ml penicillin, 100 μ g/ml streptomycin, and 10% fetal calf serum (FBS). The plasmid was transfected into the amphotropic Phoenix retrovirus packaging cell line by FuGENE™6 (Boehringer Mannheim, Roche Diagnostics, Laval, PQ) to generate a retroviral stock.




Fig. 1. Schematic diagram of pBabePuro-Snail with no tag ("Addgene plasmid 19292" which quoted from <http://www.addgene.org/pgvec1?f=c&identifier=19292&atqx=snail&cmd=findpl>).

2. Cell culture

The HaCaT cell line was cultured in a medium consisting of a 3:1 ratio of DMEM and Ham's nutrient mixture F12 (Gibco BRL, Grand island, NY), supplemented with 10% FBS, 1×10^{-10} M cholera toxin, 4 mg/ml hydrocortisone, 5 μ g/ml transferring, and 2×10^{-11} M triiodothyronine (T3) under 5% CO_2 atmosphere at 37°C.

The virus-containing supernatant was obtained from the retroviral stock and applied to the cultured cell line.

3. Western blot analysis

A cell lysate was obtained from the transfected cell line and was extracted with a buffer solution of 50 mM Tris-HCl (pH 8.0), 150 mM NaCl, 0.1% SDS (sodium docecyl sulfate, Sigma, Saint Louis, MO), 1% NP₄O, and 1x Protease inhibitors (Roche, Indianapolis, IN). The proteins were resolved by 12% and 8% SDS-PAGE and transferred to nitrocellulose membranes. The membranes was first blocked by 5% skim milk for 2 hours, and then applied with the primary antibodies, anti-FLAG (monoclonal, Sigma-Aldrich, Saint Louis, MO), anti-E-cadherin (polyclonal, Santa Cruz, Santa Cruz, CA), and anti- β -actin (polyclonal, Sigma-Aldrich, Saint Louis, MO). The membranes were washed with phosphate buffered saline tween-20 (PBST) including 20 mM Tris-HCl (pH 7.5), 100 mM NaCl, 0.1% Tween 20 for three times, and then the bound antibodies were applied with horseradish peroxidase conjugated secondary antibodies, respectively and visualized with the ECL system (Pierce).

4. Immunofluorescence

The cultured cells were transferred a chambered glass slide with 2×10^4 cells/chamber and fixed with 5% acetic acid in ethanol at -20°C for 10 minutes. The cells were blocked by 2 mg/ml bovine serum albumin in phosphate-buffered saline (BSA-PBS) for 30 minutes and the primary antibodies for cytokeratin AE 1/3 (1:50, monoclonal, DAKO, Glostrup, Denmark), E-cadherin (1:100, monoclonal, BD Bioscience, San Diego, CA), and vimentin (1:100, monoclonal, DAKO, Glostrup, Denmark) were applied for 30 minutes at 4°C. After washing, they were stained with Alexa Fluor 594 dye (Molecular Probes Inc., Eugene, OR) conjugated secondary antibodies and fixed with ice-cold methanol for 10 minutes. They were then examined with a fluorescence microscope under FluoroGuard anti-fade mounting reagent (Bio-Rad, Hercules, CA).

5. *In vivo* tumorigenesis

The control HaCaT cells and transfected HaCaT-Snail cells were obtained at 5×10^6 cells/100 μ l concentrations and 0.1 ml was injected into buccal vestibules of nude mice. The mice were sacrificed 8 weeks after inoculation to excise the resulting tumor mass, including surrounding tissues.

6. Histological Studies

The specimens were obtained and fixed in 10% neutral formalin for 24 hours, then embedded in paraffin blocks, from which 4 μ m thick sections were prepared and transferred to glass slide.

7. Immunohistochemistry

The slides were deparaffinized in xylene for 30 minutes and rehydrated by sequential transfer to a 95%, 90%, 70% ethyl alcohol and distilled water. They were then immersed in 3% hydrogen peroxide for 20 minutes, to block the endogenous peroxidase activity. We used the Histostan-Plus kit (Zymed, South San Francisco, CA) to perform the immunohistochemical staining. Primary antibodies for this staining were E-cadherin (1:500 dilution, monoclonal, BD Bioscience, San Diego, CA), N-cadherin (1:100 dilution, monoclonal, Santa Cruz, Santa Cruz, CA), cytokeratin AE 1/3 (1:100, dilution, monoclonal, DAKO, Glostrup, Denmark), and vimentin (1:100 dilution, monoclonal, DAKO, Glostrup, Denmark). We washed the slides with phosphate-buffered saline (PBS) at every step. Finally, the slides were developed with DAB (3,3'-diaminobenzidine, Vector), and were counter-stained with Mayer's hematoxylin (Fisher Scientific, Pittsburgh, PA). PBS solutions were used as negative controls and underlying connective tissues in the normal oral mucosa were used as positive controls.

Results

1. Western blot analysis

FLAG expressions were confirmed in Snail-transfected cell lines and HaCaT-Snail cells showed repression of E-cadherin (Fig. 2).

2. Phase contrast and fluorescence microscopy




Fig. 2. The results of Western blotting after Snail transfection (left: HaCaT cell line; right: HaCaT-Snail cell line).

Table 1. Tumorigenesis after inoculation of HaCaT and HaCaT-Snail cell lines

Group	Mass formations
HaCaT	0 / 10
HaCaT-Snail	14 / 15

HaCaT-Snail cells showed loss of cell-cell adhesions in the phase contrast microscope and null expressions of cytokeratin and E-cadherin, and positive expressions of vimentin, in immunofluorescence microscopy (Fig. 3).

3. *In vivo* Tumorigenesis

We identified tumor masses in 14 out of 15 nude mice in the HaCaT-Snail cell inoculation group, but no tumors were present in any of the 10 nude mice in the control group (Table 1). Tumor masses in buccal vestibules extended to the lower face of nude mice (Fig. 4).

4. Histological Features

Tumors showed features of poorly differentiated carcinoma with central necrosis (Fig. 5), with invasion to neighboring muscles and bones (Fig. 6).

5. Immunohistochemistry

The HaCaT-Snail tumors showed decreased expressions of E-cadherin and cytokeratin, but showed increased expressions of vimentin and N-cadherin. E-cadherin was only slightly expressed while vimentin was substantially expressed. Cytokeratin showed negative expressions in almost all areas except some necrotic cells (Fig. 7).




Fig. 3. Photographs of the phase contrast and fluorescence microscopy images, showing positive expressions as red colors.



Fig. 4. Photographs showing the tumor mass within the buccal cheek and vestibule of a nude mouse, extending to the lower face




Fig. 5. Photographs showing the histological features of tumor mass (magnification, left: X 10 ; right: X 400). There are central necrotic areas in the tumor (left).




Fig. 6. Photographs showing invasions to neighboring muscles and bones (magnification, both: X 40).




Fig. 7. Photographs of immunohistochemical staining showing decreased expressions of E-cadherin and cytokeratin and increased expressions of vimentin and N-cadherin (magnification, cytokeratin and N-cadherin: X 40; E-cadherin and vimentin: X 100).

Discussions

Tumorigenesis is now known to be a sequential multistep processes of genetic alterations^{17,18}. As a result, subclones which have some advantages in survival can proceed to proliferation and reach potentials of growth, invasion and metastasis¹⁹. During these processes, epithelial tumor cells lose their cell-cell adhesions, have increased mobilities, and degrade their underlying connective tissue²⁰. For these biochemical properties, they have to undergo epithelial-mesenchymal transition at first and reduce their E-cadherin expressions.

E-cadherins are bound to E-cadherins of neighboring cells to maintain extracellular cell-cell adhesions and are also bound to actin, via catenins, to maintain intracellular cytoskeletons.

β -catenins are known to have some important roles in tumorigenesis²¹. They act as transcription factors that control the Lef/Tcf promoter, stabilize intracellular cytoskeletons, and bind to E-cadherins^{22,23}.

Snail acts on the transcriptional control region of E-cadherin to repress its expression and increases the concentration of free β -catenins in the cytosol to activate a variety of tumor related genes²⁴.

Cano et al.²⁵ and Peinado et al.²⁶ reported subcutaneous formations in nude mice by Snail transfected cell lines. We also confirmed tumorigenesis in HaCaT cells, which used to be non-tumorigenic immortal keratinocytes, by Snail transfection. Snail is known to be degraded through ubiquitination via GSK-3 β mediated phosphorylation and is so unstable that it has a short half-life of about 25 minutes²⁷. We made a stabilized Snail transfectant that could induce E-cadherin repression.

We also observed increased expressions of N-cadherin. Hazan et al. reported increased expressions of N-cadherin in E-cadherin repressed tumor cells with high invasiveness^{28,29}. N-cadherin is expressed in some tumors and is involved in cell mobility and invasion. It involves heterophilic cell-cell adhesion as well as homophilic cell-cell adhesion and its level could be increased by Snail in this study.

We investigated the *in vivo* state of Snail induced tumors in nude mice. The *in vivo* tumor model can have various phenomena, including central necrosis, invasion to surrounding tissues, and neoangiogenesis, as seen in naturally occurring tumors. In this study, we histologically observed partial necrotic regions and invasions to the underlying bones and muscles, and while we immunohistochemically saw uneven staining patterns of E-cadherin and vimentin within the tumor mass. This may be a result of formations of certain different intracлонаl variants or

subclones in accordance with their environments.

We also observed that the tumors had the ability to infiltrate into their surrounding tissues, but we failed to find any distant metastatic lesions in any other organs. Certain other factors should be considered as inducing distant metastasis.

In conclusion, improved productivity of tumors can be obtained by Snail-transfected xenografts and various histopathological features, including invasive growth, can be observed with these *in vivo* tumors. This animal model can aid in the study of tumor progression and the interaction with surrounding tissues.

References

1. Arias AM. Epithelial mesenchymal interactions in cancer and development. *Cell* 2001;105:424-31.
2. Savagner P. Leaving the neighborhood: Molecular mechanisms involved during epithelial-mesenchymal transition. *Bioessays* 2001;23:912-23.
3. Weis WI, Nelson WJ. Re-solving the cadherin-catenin-actin conundrum. *J Biol Chem* 2006;281(47):35593-7.
4. Bracke ME, Van Roy FM, Mareel M. The E-cadherin/catenin complex in invasion metastasis. In *Attempts to Understand Metastasis Formation I* (Günther U, Birchmeier W, eds), 1996, 123-61, Springer, Berlin
5. Mareel M, Boterberg T, Noe V, Van Hoorde L, Vermeulen S, Bruyneel E, Bracke M. E-cadherin/catenin/cytoskeleton complex: a regulator of cancer invasion. *J Cell Physiol*. 1997;173(2): 271-4.
6. Vleminckx K, Vakaet L Jr, Mareel M, Fiers W, Van Roy FM. Genetic manipulation of E-cadherin expression by epithelial tumor cells reveals an invasion suppressor role. *Cell* 1991;66(1): 107-119.
7. Campbell RJ, Pignatelli M. Molecular histology in the study of solid tumours. *J Clin Pathol Mol Pathol* 2002;55:80-2.
8. Derycke LD, Bracke ME. N-cadherin in the spotlight of cell-cell adhesion, differentiation, embryogenesis, invasion and signaling. *Int J Dev Biol* 2004;48:463-76
9. Cavallaro U. N-cadherin as an invasion promoter: a novel target for antitumor therapy? *Curr Opin Investg Drugs* 2004;5(12): 1274-8.
10. Rosivatz E, Becker I, Specht K, Fricke E, Lubber B, Busch R, Hofler H, Becker KF. Differential expression of the epithelial-mesenchymal transition regulators snail, SIP1, and twist in gastric cancer. *Am J pathol* 2002;161(5):1881-91.
11. Tomita K, Bokhoven A, van Leenders GJL, Ruijter ETG, Jansen CFJ, Bussemakers MJG, Shalken JA. Cadherin switching in human prostate cancer progression. *Cancer Res* 2000;60:3650-4.
12. Hazan RB, Phillips GR, Qiao RF, Norton L, Aaronson SA. Exogenous expression of N-cadherin in breast cancer cells induced cell migration, invasion, and metastasis. *J Cell Biol* 2000;148:779-90.
13. Bouley JL, Dennefeld C, Alberga A. The Drosophila developmental gene snail encodes a protein with nucleic acid binding fingers. *Nature* 1987;330(6146):395-8.
14. Yokoyama K, Kamata N, Hayashi E, Hoteiya T, Ueda N, Fujimoto R, Nagayama M. Reverse correlation of E-cadherin and snail expression in oral squamous cell carcinoma cells in vitro. *Oral Oncol*. 2001;37: 65-71.
15. Takkunen M, Grenman R, Hukkanen M, García de Herreros A, Virtanen I. Snail-dependent and -independent epithelial-mesenchymal transition in oral squamous carcinoma cells. *J*

- Histochem Cytochem 2006; 54(11):1263-75.
16. Fusenig NE, Boukamp P. Multiple stages and genetic alterations in immortalization malignant transformation, and tumor progression of human skin keratinocytes. *Mol Carcinog* 1998; 23(3): 144-58.
 17. Vogelstein B, Kinzler KW. Cancer genes and the pathways they control. *Nat Med* 2004;10(8):789-99.
 18. Karakosta A, Golias Ch, Charalabopoulos A, Peschos D, Batistatou A, Charalabopoulos K. Genetic models of human cancer as a multistep process. Paradigm models of colorectal cancer, breast cancer, and chronic myelogenous and acute lymphoblastic leukaemia. *J Exp Clin Cancer Res.* 2005;24(4):505-14.
 19. Loeb LA, Loeb KR, Anderson JP. Multiple mutations and cancer. *Proc Natl Acad Sci USA* 2003;100(3):776-81.
 20. Lyons AJ, Jones J. Cell adhesion molecules, the extracellular matrix and oral squamous cell carcinoma. *Int J Oral Maxillofac Surg* 2007;36(8):671-9.
 21. Fushs SY, Ougolkov AV, Spiegelman VS, Minamoto T. Oncogenic beta-catenin signaling networks in colorectal cancer. *Cell Cycle* 2005;4(11):1522-39.
 22. Brabletz T, Hlubek F, Spaderna S, Schmalhofer O, Hiendlemeyer E, Jung A, Kirchner T. Invasion and metastasis in colorectal cancer: Epithelial-mesenchymal transition, mesenchymal-epithelial transition, stem cells and β -catenin. *Cells Tissue Organs* 2005; 179:56-65.
 23. Hoppler S, Kavanagh CL. Wnt signaling: variety at the core. *J Cell Sci* 2007;120:385-93.
 24. Nelson WJ, Nusse R. Convergence of Wnt, β -catenin, and cadherin pathways. *Science* 2004;303:1483-87.
 25. Cano A, Perez-Moreno MA, Rodrigo I, Locascio A, Blanco MJ, del Barrio MG, Portillo F, Nieto MA. The transcription factor Snail controls epithelial-mesenchymal transitions by repressing E-cadherin expression. *Nat Cell Biol* 2000;2:76-83.
 26. Peinado H, Marin F, Cubillo E, Stark HJ, Fusenig N, Nieto MA, Cano A. Snail and E47 repressors of E-cadherin induce distinct invasive and angiogenic properties *in vivo*. *J Cell Sci* 2004;117(17):2827-39.
 27. Zhou BP, Deng J, Xia W, Xu J, Li YM, Gunduz M, Hung MC. Dual regulation of Snail by GSK-3 β -mediated phosphorylation in control of epithelial-mesenchymal transition. *Nat Cell Biol* 2004;6(10):931-40.
 28. Hazan RB, Kang L, Whooley BP, Borgen PI. N-cadherin promotes adhesion between invasive breast cancer cells and the stroma. *Cell Adhes Commun* 1997;4:399-411.
 29. Hazan RB, Qiao R, Keren R, Badano I, Suyama K. Cadherin switch in tumor progression. *Ann N Y Acad Sci* 2004;1014:155-63.