

Blood Concentration of Polycyclic Aromatic Hydrocarbons from Non-Occupational Exposure in farming towns near metropolitan Busan, Korea : Environmental Tobacco Smoke and Roasted Food Intake as Influential Factors

Chan-Seok Moon^{1*} · Eun Mi Jo² · Chae-Kwan Lee³ · Jung Man Kim² · Young Seoub Hong² · Bu-Soon Son⁴ · Jong-Min Paik¹

¹Dept. of Industrial Health, College of Applied Science, Catholic University of Pusan, Busan, Korea

²Dept. of Preventive Medicine, Dong-A University, Busan, Korea

³Institute of Industrial Medicine, Inje University College of Medicine, Busan, Korea

⁴Dept. of Environment Health Science, Soonchunhyang University, Chungnam, Korea

부산인근 농촌지역의 비직업적 노출에 의한 혈중 다환방향족탄화수소 농도: 간접흡연과 구운음식 섭취를 중심으로

문찬석^{1*} · 조은미² · 이채관³ · 김정만² · 홍영섭² · 손부순⁴ · 백종민¹

¹부산가톨릭대학교 산업보건학과

²부산 동아대학교 예방의학교실

³부산 인제대학교 산업의학연구소

⁴순천향대학교 환경보건학과

연구목적: 본 연구는 일반인의 비직업적 노출에 대한 혈중 16종의 다환방향족탄화수소(PAHs)의 농도 분석하여 PAHs의 백그라운드 노출 농도를 평가하는 것이다.

내용 및 방법: 농촌지역에 거주(비직업적 노출로서 16종 다환방향족탄화수소가 특정한 환경오염을 받지 않은 지역에 거주)하는 156명의 연구 참여자의 혈중 16가지 다환방향족탄화수소 농도의 기하평균치를 간접흡연군, 구운음식물섭취군과 각대조군을 비교 하였다. 혈중 16종의 다환방향족탄화수소의 농도는 질량분석기를 장착한 가스 크로마토그래피법과 헤드스페이스 마이크로고체상 추출법에 의하여 분석하였다.

결과: 연구참여자 중, 간접흡연자들에서 4종의 다환방향족탄화수소 아세나프틸렌($p<0.01$), 아세나프텐($p<0.1$), 플

로렌($p<0.01$), 피렌($p=0.05$)가 대조군 보다 유의하게 높았다. 또한 구운육류와 생선 섭취군에서는 벤조(a)피렌($p<0.1$)이 대조군 보다 유의하게 높았다.

결론: 간접흡연과 구운음식물 섭취는 비직업적 노출에서 다환방향족탄화수소 노출원의 가능성이 높다고 추정된다. 혈중 다환방향족탄화수소는 한국일반인의 비직업적 노출 농도로 사용 가능하다고 생각된다.

Key word: Non-occupational exposure, polycyclic aromatic hydrocarbons, blood, environmental tobacco smoking, roasted food intake

접수일: 2010년 4월 21일, 채택일: 2010년 6월 18일

* 교신저자: 문찬석(부산광역시 금정구 부곡3동 9번지 부산가톨릭대학교 산업보건학과,
Tel.: +82-51-510-0633, Fax.: +82-51-510-0638, E-Mail: csmoon@cup.ac.kr)

I . Introduction

Polycyclic aromatic hydrocarbons (PAHs) are widely-distributed and hazardous environmental pollutants that are discharged into the environment from the versatile sources such as motor vehicle emissions, fossil fuels, industrial plants. They are produced during the incomplete combustion of coal, gas, oil, and wood, and can also be found in tobacco smoke, charbroiled meats, and smoked foods (International Agency for Research on Cancer, 1985; International Agency for Research on Cancer, 1987). Industries in exposure to PAHs is likely to occur include those using coke ovens and coal tar, iron and steel works, aluminum works, foundries, carbon electrode and carbon black manufacture, asphalt manufacture and use, etc. (Redmond et al., 1976; Palmer and Scott, 1981; Armstrong et al., 1986; NIEHs, 2000; Campo et al., 2006; Unwin et al., 2006; Buratti et al., 2007). Many of these PAHs shown carcinogenic activity, proven in animals, and are thought to have a similar impact on humans (Boffetta et al., 1997; International Agency for Research on Cancer, 1973; International Agency for Research on Cancer, 1983; International Agency for Research on Cancer, 1984; International Agency for Research on Cancer, 1984a; National Toxicology Program, 2001). Background exposures to PAHs for the general population typically come from sources such as tobacco smoke or environmental tobacco smoke (ETS), charbroiled or smoked foods, and automobile exhaust fumes (WHO, 1984; Grimmer et al., 1987; Liroy et al., 1988; Scherer et al., 1990; Scherer et al., 2000; Kim et al., 2001; Hu et al., 2006).

As representative biological exposure marker, urinary 1-hydroxypyrene (1-OHP) for environmental PAHs exposure has been somewhat limitation for the explanation of environmental PAHs exposure (WHO, 1984). Moreover, the marker is not mostly influenced by respiratory intake but by food intake and tobacco smoking among general population (Grimmer et al., 1987; Liroy et al., 1988; Scherer et al., 1990; Jongeneelen et al., 2001). In these sense, urinary 1-OHP is total exposure marker for pyrene exposure via respiration and food intake so that PAHs exposure sources are not clear whether exposure source is from ambient air or from food intake. More effective exposure markers than urinary 1-OHP are necessary to PAHs exposure via respiratory intake. Blood PAHs are, therefore, thought to more effective exposure marker than 1-OHP as urinary metabolite.

A venous blood sample is a most effective and sensitive biological matrix in cases of low-level exposure to volatile organic compounds (Kawai et al., 1992; Kawai et al., 1992a; Kawai et al., 1994; Kawai

et al., 1996; Ikeda, 1999). PAHs in blood have been used as exposure markers for long-term low-level exposure. However, few reports are available for the evaluation of the PAHs in blood for non-occupational exposure. Moreover, a large-scale national field survey (cohort study) for the health effects on residents living near industrial sites was conducted in six Korean industrial areas every year for 20 years, in order to clarify the exposure to PAHs from the combustion of coal, gas, oil etc. The detection of PAHs in blood is required a sensitive and rapid method to handle a large number of blood samples in a very short detection period. The present study was initiated to examine the background levels of US EPA 16 PAHs (naphthalene, acenaphthylene, acenaphthene, fluorene, phenanthrene, anthracene, fluoranthene, pyrene, benzo(a)anthracene, chrysene, benzo(b)fluoranthene, benzo(k)fluoranthene, benzo(a)pyrene, indeno(1,2,3-cd)pyrene, dibenz(a,h)anthracene, benzo(g,h,i)perylene) in venous blood samples. The blood concentration of 16 PAHs samples were compared exposed group to ETS and nonsmoker group, and group of roasted food intake with non-intake group of these foods.

II . Materials and Methods

1. Survey participants and study design

This survey was conducted from February to April 2008 in several farming towns near metropolitan Busan, Korea, where there is no known industrial pollution from PAHs. The participants were 158 healthy non-smoking participants (68 man and 90 women), with an age range of 21-68 years (mean \pm SD, 43.8 \pm 19.4), (Table 1). The subjects were divided into subgroups of nonsmokers exposed to ETS and non-smoker, or intake of roasted meats or fish and non-intake of these foods. Then, significant increases in 16 PAH concentrations in blood samples were evaluated.

In selecting the group of nonsmokers exposed to ETS, the participants were asked to describe their exposure to ETS in terms of minutes per day, and were accepted if they had an ETS exposure of up to 1 minute a day. Roasted meats or fish involved charbroiled, briquette-roasted, butane gas-roasted, city gas (LNG)-roasted or electronic grill-roasted meats or fish. Moreover, the questionnaires were fully explained the participants.

2. Sample collection and analysis of the 16 PAHs

Venous blood samples and questionnaires were collected from the participants with informed consents provided in writing. Heparinized vacuum tubes for blood sampling (Vacutainer, Becton Dickinson, N.J., USA) were employed (Moon et al., 1995) with due caution to avoid the contamination of PAHs. Then, each blood sample was stored at -18 °C until it was analyzed.

PAHs in blood were analyzed using headspace-solid phase microextraction (CTC CombiPAL automated sample injectors for

gas chromatography, Agilent Technologies, Inc., USA) coupled with gas chromatography-mass spectrometry [HS-SPME/GC-MS (GCMS-QP2010, Shimadzu, Japan)] according to published methods (Aguinaga et al., 2007; Campo et al., 2006), with slight modifications made. Whole blood (1 ml) was transferred to a vial containing 100 mg of NaCl and K₂CO₃ and 1 ppm of internal standard solution (phenanthrene-d₁₀ and pyrene-d₁₀) was added. Analytes were extracted from the headspace of the samples using a 65 μm PDMS/DVB SPME fiber (Supelco) with agitation. To increase absorption efficiency to the fiber, an absorption temperature of

Table 1. The general characteristics of the study participants *

	Man (68)		Women (90)		Total (158) [†]	
	AM ± ASD	Min.-Max.	AM ± ASD	Min.-Max.	AM ± ASD	Min.-Max.
Age (years)	44.8 ± 21.3	(21-67)	43.1 ± 17.7	(21-68)	43.8 ± 19.4	(21-68)
Height (cm)	167.4 ± 6.8	(150.0-181.0)	157.8 ± 6.1	(141.0-171.5)	162.0 ± 8.0	(141.0-181.0)
Weight (kg)	63.7 ± 11.4	(41.0-98.0)	56.3 ± 9.8	(40.0-93.0)	59.5 ± 11.1	(40.0-98.0)
Body Mass Index	22.6 ± 3.3	(17.6-32.5)	22.6 ± 3.6	(15.0-33.7)	22.6 ± 3.5	(15.0-33.7)

* healthy non-smoker.

† number of participants

Table 2. Analytical condition of subjected 16 PAHs

	Retention time (min)	Selected ion (m/z)	Regression parameter*		Detection limit (μg/l)
			α	β	
Naphthalene	10.111	128, 129, 127	0.0064	0.0983	0.8
Acenaphthylene	14.214	152, 151, 153	0.0025	0.0107	0.7
Acenaphthene	14.690	154, 153, 152	0.0179	0.0989	0.8
Fluorene	16.003	166, 165, 167	0.0216	0.0127	0.7
Phenanthrene	18.441	178, 179, 176	0.0104	0.0579	0.9
Anthracene	18.560	178, 176, 179	0.0984	0.0507	0.9
Fluoranthene	21.478	202, 101, 203	0.0563	0.0147	0.8
Pyrene	22.026	202, 200, 203	0.0319	0.0193	0.9
Benzo(a)anthracene	25.078	228, 229, 226	0.0296	0.0218	0.8
Chrysene	25.172	228, 226, 229	0.0568	0.0286	1.0
Benzo(b)fluoranthene	27.650	252, 253, 125	-0.0172	0.0107	0.9
Benzo(k)fluoranthene	27.683	252, 253, 125	0.0772	0.0319	0.9
Benzo(a)pyrene	28.367	252, 253, 125	0.0327	0.0133	0.9
Indeno(1,2,3-cd)pyrene	30.864	276, 138, 227	-0.0054	0.0074	1.1
Dibenz(a,h)anthracene	30.895	278, 139, 279	-0.0045	0.0231	1.3
Benzo(g,h,i)perylene	30.926	276, 138, 277	-0.0183	0.0178	1.3

* α and β are parameters of a regression line of $Y = \alpha + \beta X$, where X is the blood concentration (μg/l) and Y is the value for mass chromatogram. $r > 0.99$ in all of 16 PAHs.

Table 3. Geometric mean and maximum concentration of the 16 PAHs in blood samples ($\mu\text{g/l}$)

	Man (No. 68)	Women (No. 90)	Total [maximum] (No. 158)
Naphthalene	27.66	23.26	25.20 [918.45]
Phenanthrene	4.08	4.61	4.36 [64.66]
Anthracene	4.26	4.16	4.21 [61.65]
Acenaphthylene	3.80	3.10	3.41 [325.44]
Pyrene	1.72	1.82	1.77 [75.08]
Benz(a)anthracene	1.44	1.39	1.41 [65.89]
Chrysene	1.91	1.49	1.67 [24.08]
Dibenzo(a,h)anthracene	2.35	2.85	2.60 [26.15]
Benzo(g,h,i)perylene	1.98	2.71	2.34 [17.22]
Fluoranthene	1.71	1.76	1.76 [150.31]
Acenaphthene	1.35	1.48	1.42 [5.98]
Benzo(a)pyrene	1.32	1.36	1.34 [34.64]
Benzo(k)fluoranthene	1.98	1.67	1.81 [242.61]
Fluorine	0.97	0.99	0.98 [227.97]
Indeno(1,2,3-cd)pyrene	1.16	1.07	1.11 [68.83]
Benzo(b)fluoranthene	-	-	-

Student t-test between man and women were not statistical significant with all of 16 analytes ($p > 0.05$).

90 °C for 50 min was applied. The analytes were then desorbed from the fiber for 5 min in the injector, operating in a splitless mode at a temperature of 250 °C, equipped with an inlet liner for SPME (internal diameter 0.75mm, Supelco).

The GC conditions used were as follows: HP-5 capillary GC column (30 m, 0.25 μm inner diameter, 0.25 μm film thickness), helium carrier gas at a flow rate of 1 ml/min; gas chromatograph oven temperature programmed from 40 °C (2 min initial hold) to 230 °C at 15 °C/min (then 2 min. hold), then to 230 °C at 5 °C/min (then 6 min hold), and finally to 300 °C at 10 °C/min (then 15 min hold). The MS conditions were 280 °C in transfer line temperature and 200 °C in ion source temperature. The detector was operated in the selected ion monitoring mode (SIM) for nominal molecular ions (M/Z) (Table 2).

3. Statistical analysis

The statistical analysis was performed using SPSS 17.0 for Windows. Arithmetic means (AM) and arithmetic standard deviations (ASD) were employed for age, height, weight and body mass index (BMI) because these measures were assumed to be

distributed normally. A log-normal distribution was assumed for the 16 PAHs (naphthalene, acenaphthylene, acenaphthene, fluorene, phenanthrene, anthracene, fluoranthene, pyrene, benzo(a)anthracene, chrysene, benzo(b)fluoranthene, benzo(k)fluoranthene, benzo(a)pyrene, indeno(1,2,3-cd)pyrene, dibenz(a,h)anthracene, and benzo(g,h,i)perylene), so that geometric means (GMs) and geometric standard deviations (GSD) were taken to express the distribution. Student's t-test was employed for comparison between the groups of man and women, nonsmokers exposed to ETS and the control group, and intake of roasted meat or fish and the control group.

To calculate the GM and GSD, the value below the detection limit (DL) was assumed as one-half of the DL. DLs in values from GC/MS ranged from 0.7 to 1.3 $\mu\text{g/l}$ for the 16 PAHs.

III. Results

1. Geometric means (GMs) and maximum of the 16 PAHs in blood




Fig. 1 The comparison of GM of blood concentration between the group of ETS smoker (Group A), and control group (Control)

The 16 PAHs measured in the blood samples of the study participants are summarized in Tables 3. GMs of 16 PAHs between man and women were not significantly different ($p > 0.05$). GMs of naphthalene in blood were $25.20 \mu\text{g/l}$ in total, $27.66 \mu\text{g/l}$ in man, and $23.26 \mu\text{g/l}$ in women, respectively. That of Phenanthrene in blood were $4.46 \mu\text{g/l}$ in total, $4.08 \mu\text{g/l}$ in man, and $4.61 \mu\text{g/l}$ in women. Anthracene were $4.21 \mu\text{g/l}$ in total, $4.26 \mu\text{g/l}$ in man, and $4.16 \mu\text{g/l}$ in women. Acenaphthylene were $3.41 \mu\text{g/l}$ in total, $3.80 \mu\text{g/l}$ in man and 3.10 in women, respectively. Most of the 16 PAHs analytes showed low blood concentration, however, maximums of naphthalene ($918.45 \mu\text{g/l}$), acenaphthylene ($325.44 \mu\text{g/l}$), fluoranthene ($150.31 \mu\text{g/l}$), benzo(k)fluoranthene ($242.61 \mu\text{g/l}$), fluorine ($227.97 \mu\text{g/l}$) shown to high levels of blood concentration. Pyrene, benzo(a)pyrene, indeno(1,2,3-cd)pyrene as pyrenes were $1.77 \mu\text{g/l}$ (GM in total), $75.08 \mu\text{g/l}$ (maximum) in pyrene, $1.34 \mu\text{g/l}$ (GM in total), and $34.64 \mu\text{g/l}$ (maximum) in benzo(a)pyrene, and $1.11 \mu\text{g/l}$, $68.83 \mu\text{g/l}$ (maximum) in indeno(1,2,3-cd)pyrene. These 3 PAHs analytes were not showed to high blood concentration

2. Blood PAH concentration and nonsmokers exposed to ETS.

In environmental tobacco smoking (ETS), survey participants were evaluated for blood PAH concentrations, as shown in Fig.1. The numbers for nonsmokers exposed to ETS, and nonsmokers (control group) were 18 and 138, respectively. Among the 16 PAHs, 4 PAHs analytes [acenaphthylene ($p < 0.01$), acenaphthene ($p < 0.1$),




Fig. 2 The comparison of GM of blood concentration between the group of intake of roasted meats or fish for the past 3 days (Group B), and control group (Control)

fluorene ($p < 0.01$), pyrene ($p < 0.05$)] in GM of blood PAHs were higher in the group with exposure to ETS than in the control group. Anthracene, benzo(a)pyrene, and benz(a)anthracene also had higher GMs in the group exposed to ETS, but with no statistical significance.

3. Blood PAH concentration and intake of roasted meats or fish

Participants (69) had consumed roasted meats or fish in the previous three days (Group B), and 19 participants had not (control) intake these roasted foods. Figure 2 highlights the comparison between Group B and the control group for blood concentrations of the 16 PAHs. Among the geometric means of the 16 PAHs, only benzo(a)pyrene was significantly higher in Group B (intake of roasted meats or fish in the previous three days) than in the control group ($p < 0.1$). Acenaphthylene, phenanthrene, anthracene, fluoranthene, pyrene, benz(a)anthracene, and chrysene were higher in Group B than in the control group, however, there was no statistical significance ($p > 0.1$).

IV. Discussion

In the study, very low level 16 PAHs in blood could efficiently detected for the evaluation of non-occupational exposure. The head space-solid phase microextraction (HS-SPME) method was applied

for the detection of 16 PAHs in blood. This method did not require large volumes of organic solvents for extraction, which can be time consuming and may involve a multi-step process that could result in the loss of some analytes (King et al., 2003). Moreover, simplicity, low cost and high sensitivity make this method very useful for quantitative analysis of volatile or semi-volatile compounds (Cam et al., 2004; Ahn et al., 2001). By using the head space analysis method, the GC column is influenced by the direct injection of the sample to the GC injection port, rather than by the sample contents, so that a large number of samples provide efficient qualitative analysis. Moreover, the solvent-extraction and concentration required by existing methods requires a longer experimental time, a large amount of solvents and man-power for the experiment, so that the method is thought to be very limited in cases with large samples and reduced analytical working periods.

GM, and the maximum of PAHs in blood are elucidated in Tables 3. The GM of Naphthalene was 25.20 $\mu\text{g/l}$ in total among the PAH analytes. However, there was no significant variation in the GM for blood concentration between man and women. The maximum was 1118.49 $\mu\text{g/l}$ for man and women combined. If the participant had not been exposed to smoking or ETS or motor vehicle emissions, or any other occupational exposure sources, the other source of exposure to naphthalene for non-smokers is likely to be mothballs (used in toilets and to protect clothing), which represents a continuous exposure. More studies are necessary to evaluate the impact of indoor air pollutants on the general Korean population.

One of the main sources of 16 PAHs exposure for the general population is cigarette smoke (Hu et al., 2006; Ding et al., 2006). This study showed that there was an effect of ETS in the passive smoking group because 4 analytes, acenaphthylene ($p<0.01$), acenaphthene ($p<0.1$), fluorene ($p<0.01$), and pyrene ($p<0.05$), among the 16 PAHs showed the significant increase in the blood samples. The participants showed the significant increase in the 4 PAHs were mainly exposed a particulate phase of the ETS. In the report from Lu and Zhu (2007), the particulate phase showed fluorene, phenanthrene, benzo(g,h,i)perylene at high concentrations. On the other hand, the vapor phase showed high concentrations of acenaphthene, acenaphthylene, fluorene, phenanthrene, fluoranthene, and pyrene (Lu and Zhu, 2007). Therefore, as a result of this study, 4 PAHs from the blood regarded as an indicator of the exposure of ETS. Benzo(a)pyrene was related to smoking or to ETS (Fagundes et al., 2006). The results of the study, however, showed a very low detection percentage (12.7 % in total). Further studies of dose-response relations between blood PAHs and exposure to ETS

are needed to clarify the exposure source and route for the general Korean population.

Participants who had consumed roasted meats or fish showed a significant increase in benzo(a)pyrene levels (Fig. 2). Reinik et al., 2007 reported that the maximum acceptable concentration for benzo(a)pyrene of 5 $\mu\text{g/kg}$ was exceeded in 3.4 % of samples. These samples included commercial, cured meat products, and home grilled meats in Estonia. In non-smoking Japanese university students, pyrene, benzo(k)fluoranthene, and benzo(a)pyrene levels primarily came from food intake (Suzuki and Yoshinaga, 2007). Moreover, benzo(a)pyrene was identified as a principal dietary source in the adult Spanish population (Ibáñez et al., 2005). When the results of the current study were compared to recent reports, blood benzo(a)pyrene was closely related to meat intake or the method used for cooking the meat (Sinha et al., 2005; Kazerouni et al., 2001; Phillips, 1999). Thus it is assumed that levels of benzo(a)pyrene in the blood are influenced by exposure to roasted meats or fish. No trials were possible in the present study, however, to make quantitative assessment of roasted food intake. Further studies are necessary to identify such factors quantitatively as being influential sources of PAHs exposure to general population.

V. Conclusion

The blood PAHs were efficiently detected by GC-MS in non-occupational exposed subjects. ETS is an influential factor, which it could affect the increase of blood PAHs [i.e., four PAHs, acenaphthylene ($p<0.01$), fluorene ($p<0.01$), acenaphthene ($p<0.1$), and pyrene($p=0.1$)]. The roasted meat and fish ingestion was also found to be influential factors, which were increased the concentrations of benzo(a)pyrene in the blood. The concentrations of blood PAHs in this study are regarded some of the background exposure level in the general Korean population. In relation to the dietary ingestion of 16 PAHs, the further study of Korean food is needed to clarify the daily exposure and the dose-response relationship in background exposure.

VI. Acknowledgments

This work was supported by the Korea Research Foundation Grant funded by the Korean Government (MOEHRD) (KRF-2007-331-E00055).

REFERENCES

- Aguinaga N, Campillo N, Vinas P, Hernandez-Cordoba M. Determination of 16 polycyclic aromatic hydrocarbons in milk and related products using solid-phase microextraction coupled to gas chromatography-mass spectrometry. *Analytica Chimica Acta* 2007;596: 285-290
- Ahn Y-G, Seo J-B, Hong J. Comparison solid phase microextraction with Purge & trap on the GC/MS analysis of volatile organic compounds in biota samples. *Analytical Science & Technology* 2001;145: 392-399 (in Korean)
- Armstrong BG, Tremblay CG, Cyr D, Theriault GP. Estimating the relationship between exposure to tar volatiles and the incidence of bladder cancer in aluminum smelter workers. *Scandinavian Journal of Work Environmental & Health* 1986; 12: 486-493
- Boffetta P, Jourenkova N, Custavsson P. Cancer risk from occupational and environmental exposure to polycyclic aromatic hydrocarbons. *Cancer Causes & Control* 1997; p. 444-472
- Buratti M, Campo L, Fustinoni S, Cirila PE, Martinotti I et al. Urinary hydroxylated metabolites of polycyclic aromatic hydrocarbons as biomarkers of exposure in asphalt workers. *Biomarkers* 2007;123: 221-239
- Cam D, Gagni S, Lombardi N, Punin MO. Solid-phase microextraction and gas chromatography-mass spectrometry for the determination of polycyclic aromatic hydrocarbons in environmental solid matrices. *Journal of Chromatographic Science* 2004;426: 329-35
- Campo L, Addario L, Buratti M, Scibetta L, Longhi O et al. Biological monitoring of exposure to polycyclic aromatic hydrocarbons by determination of unmetabolized compound in urine. *Toxicology Letters* 2006;162: 132-138
- Ding YS, Yan XJ, Jain RB, Lopp E, Tavakoli A et al. Determination of 14 polycyclic aromatic hydrocarbons in mainstream smoke from U.S. brand and non-U.S. brand cigarettes. *Environmental Science & Technology* 2006;404: 1133-1138
- Fagundes R B, Abnet C C, Strickland P T, Kamangar F, Roth M J et al. Higher urinary 1-hydroxy glucuronide (1-OHPG) is associated with tobacco smoke exposure and drinking mate in healthy subjects from Rio Grande do Sul, Brazil. *BMC Cancer* 2006;6: 1-7
- Gimmer G, Naujack K W, Dettbarn G. Gas chromatographic determination of polycyclic aromatic hydrocarbons, azarenes, aromatic amines in the particle and vapor phase of mainstream and side stream smoke of cigarettes. *Toxicology Letters* 1987;35: 117-124
- Hu Y, Zhou Z, Xue X, Li X, Fu J et al. Sensitive biomarker of polycyclic aromatic hydrocarbons (PAHs): urinary 1-hydroxypyrene glucuronide in relation to smoking and low ambient levels of exposure. *Biomarkers* 2006;11: 306-318
- Ibáñez R, Agudo A, Berenguer A, Jakszyn P, Tormo M J et al. Dietary intake of polycyclic aromatic hydrocarbons in a Spanish population. *Journal of Food Protection* 2005;68: 2190-2195
- Ikedo M. Solvents in urine as exposure markers. *Toxicology Letters* 1999;108: 99-106
- International Agency for Research on Cancer (). Polynuclear aromatic compounds, Part 4: Bitumens, coal-tars and derived products, shale-oils and soots. In IARC monographs on the evaluation of carcinogenic risks to humans. Lyon, France: International Agency for Research on Cancer; 1985;Vol. 35: p. 271
- International Agency for Research on Cancer . Polynuclear aromatic compounds, Part 2 Carbon Blacks, mineral oils (lubricant base oils and derived products) and some nitroarenes. In: IARC Monographs on the evaluation of carcinogenic risks to humans). Lyon, France: International Agency for Research on Cancer; 1984.; Vol. 33: p.245
- International Agency for Research on Cancer. Certain polycyclic aromatic hydrocarbons and heterocyclic compounds. In: IARC monographs on the evaluation of carcinogenic risks to humans Vol. 3 Lyon, France: International Agency for Research on Cancer; 1973. p. 271
- International Agency for Research on Cancer. Monographs of the evaluation of the carcinogenic risk of chemicals to humans. Polycyclic aromatic hydrocarbons. Part 1. Chemical, environmental and experimental data, In: IARC monographs on the evaluation of carcinogenic risks to humans Vol. 32. Lyon, France: International Agency for Research on Cancer; 1983
- International Agency for Research on Cancer. Overall evaluation of carcinogenic risks to humans. In IARC monographs on the evaluation of carcinogenic risks to humans. Supplement 7. Lyon, France: International Agency for Research on Cancer; 1987
- International Agency for Research on Cancer. Polynuclear aromatic compounds, Part 3 Industrial exposures in aluminium production, coal gasification, coke production, and iron and

- steel foundin. In IARC Nonographs on the evaluation of carcinogenic risks to humans Lyon, France: International Agency for Research on Cancer; 1984a; Vol. 34: (p.219)
- Jongeneelen FJ. Benchmark guideline for urinary 1-hydroxypyrene as biomarker of occupational exposure to polycyclic aromatic hydrocarbons. *Ann Occup Hyg* 2001;45: 3-13
- Kawai T, Mizunuma K, Okada Y, Horiguchi S, Ikeda M et al. Toluene itself as the best urinary marker of toluene exposure. *International Archives of Occupational & Environmental Health* 1996;68: 289-297
- Kawai T, Mizunuma K, Yasugi T, Horiguchi S, Ikeda M et al. Toluene in blood as a marker of choice for low-level exposure to toluene. . *International Archives of Occupational & Environmental Health* 1994;66: 309-315
- Kawai T, Yasugi T, Mizunuma K, Horiguchi S, Iguchi H et al. Comparative evaluation of urinalysis and blood analysis as means of detecting exposure to organic solvents at low concentrations. . *International Archives of Occupational & Environmental Health* 1992a; 64: 223-234
- Kawai T, Yasugi T, Mizunuma K, Horiguchi S, Ikeda M Urinalysis vs. blood analysis, as a tool for biological monitoring of solvent exposure. *Toxicology Letters* 1992;63: 333-343
- Kazerouni N, Shnha R, Hsu C-H, Greenberg A, Rothman N. Analysis of 200 food items for benzo(a)pyrene and estimation of its intake in an epidemiologic study. *Food & Chemical Toxicology*, 2001;39: 423-436
- Kim J, Cho S-H, Kang J-W, Kim Y-D, Nan H-M, Lee C-H, Lee H, Kawamoto T. Urinary 1-hydroxypyrene and 2-naphthol concentrations in male Koreans. *International Archives of Occupational & Environmental Health* 2001;74: 59-62
- King AJ, Readman JW, Zhou JL. The application of solid-phase micro-extraction (SPME) to the analysis of polycyclic aromatic hydrocarbons (PAHs). *Environmental Geochemistry & Health* 2003;25: 69-75.
- Lioy PJ, Waldman JM, Greenberg AR. The total human environmental exposure study (THESS) to benzo(a)pyrene: Comparison of the inhalation and food pathways. *Archives of Environmental Health* 1988;43: 304-312
- Lu H and Zhu L. Pollution patterns of polycyclic aromatic hydrocarbons in tobacco smoke. *Journal of Hazardous Materials* 2007;A139: 193-198
- Moon C-S, Zhang Z-W, Shimbo S, Watanabe T, Moon D-H, Lee C-U, Lee B-K, Ahn K-D, Lee S-H, Ikeda M. Dietary intake of cadmium and lead among general population in Korea. *Environmental Research* 1995;71: 46-54
- National Institute of Environmental Health Sciences. 9th report on carcinogens. Research Triangle Park, NC, U.S., Department of Health and Human Services, Public Health Service, National Toxicology Program; 2000
- National Toxicology Program. 9th Report on carcinogens. Revised edition. U.S. Department of Health and Human Services, Public Health Service, Research Triangle Park, North Carolina: National Toxicology Program; 2001
- Palmer WG, Scott WD. Lung cancer in ferrous foundry workers: a review. *American Industrial Hygiene Association Journal* 1981;42: 329-340
- Phillips DH. Polycyclic aromatic hydrocarbons in the diet. *Mutation Research-Genetic Toxicology & Environmental Mutagenesis* 1999;443: 139-147
- Redmond CK, Strobino BR, Cypress RH. Cancer experience among coke by-product workers. *Annals of the New York Academy of Sciences* 1976;271: 102-115
- Reinik M, Tame T, Roasto M, Junkam K, Tenno T et al. Polycyclic aromatic hydrocarbons (PAHs) in meat products and estimated PAH intake by children and the general population in Estonia. *Food Additives & Contaminants* 2007;24: 427-437
- Scherer G, Conze C, VonMeyerinck L, Sorsa M, Adlkofer F et al. Importance of exposure to gaseous and particulate phase components of tobacco smoke in active and passive smokers. *International Archives of Occupational & Environmental Health* 1990;62: 459-466
- Scherer G, Frank S, Riedel K, Meger-Kossien I, Renner T. et al. Biomonitoring of exposure to polycyclic aromatic hydrocarbons of nonoccupationally exposed persons. *Cancer Epidemiology Biomarkers & Prevention* 2000;9: 373-380
- Sinha R, Peters U, Cross A J, Kulldorff M, Weissfeld J L et al. Meat, Meat cooking methods and preservation, and risk for colorectal adenoma. *Cancer Research* 2005;65: 8034-8041
- Suzuki K, Yoshinaga J. Inhalation and dietary exposure to polycyclic aromatic hydrocarbons and urinary 1-hydroxypyrene in non-smoking university students. *International Archives of Occupational & Environmental Health* 2007;81: 115-121
- Unwin J, Cocker J, Scobbie E, Chambers H. An assessment of occupational exposure to polycyclic aromatic hydrocarbons in the UK. *Annals of Occupational Hygiene* 2006;50: 395-403
- WHO. Guideline for drinking-water quality Vol. 1 Recommendation. Geneva; 1984: p.67