

Clinical Article

The Effect of Phosphodiesterase-4-Specific Inhibitor in the Rat Model of Spinal Nerve Ligation

Sung Hoon Kim, M.D., Ph.D.,¹ Bit-Na-Ri Park, M.D.,¹ Seok Won Kim, M.D., Ph.D.²

Department of Rehabilitation,¹ College of Medicine, Wonju Christian Hospital, Yonsei University, Wonju, Korea

Department of Neurosurgery,² College of Medicine, Chosun University, Gwangju, Korea

Objective : Peripheral neuropathy is characterized by hyperalgesia, spontaneous burning pain, and allodynia. The purpose of this study was to investigate the effect of rolipram, a phosphodiesterase-4-specific inhibitor, in a segmental spinal nerve ligation model in rats.

Methods : Both the L5 and L6 spinal nerves of the left side of the rats were ligated. Phosphodiesterase-4 inhibitor (rolipram) and saline (vehicle) were administered intraperitoneally. We measured mechanical allodynia using von Frey filaments and a nerve conduction study.

Results : The mechanical allodynia, which began to manifest on the first day, peaked within 2 days. Multiple intraperitoneal injections of rolipram ameliorated the mechanical allodynia. Furthermore, an intraperitoneal administration of rolipram improved the development of pain behavior and nerve conduction velocity.

Conclusion : This study suggests that the phosphodiesterase-4 inhibitor, rolipram, alleviates mechanical allodynia induced by segmental spinal nerve ligation in rats. This finding may have clinical implications.

Key Words : Peripheral nerve injury · Allodynia · Rolipram.

INTRODUCTION

Spinal radiculopathy and peripheral neuropathy may generate a syndrome characterized by spontaneous pain, and exaggerated responses to light touch and temperature stimuli. The polyneuropathy is an incapacitating complication of chronic drug consumption, characterized by allodynia and pain, primarily in the lower extremities, and is poorly relieved by available treatments²³. Depending on the criteria and patient selection, the incidence of peripheral neuropathy has been reported to range from 10% to 50%²². The pathogenesis of drug-induced neuropathy is an axonal neuropathy characterized by wallerian degeneration of the axons and a reduction in the myelination of nerve fibers³⁶. Controversy surrounds the pathogenic role of trauma or drugs in the development of neuropathy. Studies on rat models have suggested that many causative factors have a direct neurotoxic effect on neuronal organelles and the spinal cord system^{5,16,24}. Oxidative stress and inflammatory stress are known to play a very pivotal role in the experimental animal

models of neuropathic pain. Lee et al.¹⁸ suggested that reactive oxygen species are critical to the development and maintenance of capsaicin-induced pain, particularly in the process of central sensitization in the rat nervous system. Padi and Kulkarni²⁶ demonstrated that chronic administration of minocycline when started early, before peripheral nerve injury, could ameliorate the development of neuropathic pain by inhibiting the release of proinflammatory cytokines and oxidative and nitrosative stress in mononeuropathic rats. A significant increase in lipid peroxidation and decrease in the activity of antioxidant enzymes (superoxide dismutase and catalase) have been observed in the sciatic nerves of diabetic rats with established neuropathic pain³⁰. Dina et al.⁶ demonstrated that hyperalgesia is present in an established model of peripheral neuropathy in the rat and that inflammatory process and protein kinase signaling play a pivotal role in the enhanced nociception. An important action of cAMP is activation of transcription factors including c-AMP-responsive element binding (CREB) protein and nuclear factor- κ B (NF- κ B) p50¹². Phosphorylation of CREB stimulates transcription of cell survival genes²⁰. Phosphorylation of NF- κ B p50 subunit suppresses transcription of genes associated with inflammation, especially the pro-inflammatory cytokines, tumor necrosis factor- α (TNF- α) and interleukin-1 β (IL-1 β)^{4,12,19,33}. Thus, in the current study, we tested the hypothesis that rolipram, a selective inhibitor of cAMP-specific phosphodiesterase

• Received : March 25, 2011 • Revised : June 16, 2011

• Accepted : August 8, 2011

• Address for reprints : Seok Won Kim, M.D., Ph.D.

Department of Neurosurgery, School of Medicine, Chosun University,
588 Seoseok-dong, Dong-gu, Gwangju 501-717, Korea
Tel : +82-62-220-3126, Fax : +82-62-227-4575

E-mail : ns64902@hanmail.net

(PDE), would play a pivotal role in improving mechanical allodynia and nerve conduction velocity in segmental spinal nerve ligation-induced neuropathic pain in rats.

MATERIALS AND METHODS

Experimental animals

Sixteen male adult Sprague-Dawley rats weighing 200-350 g were used in this study. The animals were housed in two groups in plastic cages with soft bedding and free access to food and water. All animals were acclimated in their cages for 1 week before any experiments were performed. All experimental protocols were approved by the Institutional Animal Care and Use Committee at our institute and carried out in accordance with the National Institutes of Health's Guide for the Care and Use of Laboratory Animals.

Segmental spinal nerve ligation model and drug treatment protocol

Under sodium pentobarbital anesthesia (40 mg/kg, i.p.), the rat was placed in a prone position and the left paraspinal muscles were separated from the spinous processes at the L4-S2 level. The L6 transverse process was carefully removed to identify the spinal nerves. The left L5 and L6 spinal nerves were ligated with 6-0 silk thread. The PDE-4-specific inhibitor, rolipram (Sigma, St. Louis, MO, USA), was first dissolved and then gently mixed with 0.9% physiological saline to a final 10% v/v solution. Physiological saline was used as the vehicle for the control group. According to their specific study group, animals received rolipram or vehicle once daily for a period of three weeks.

Behavioral tests for mechanical allodynia

Behavioral tests were conducted blindly so that the experimenter who conducted the tests did not know the nature of the experimental application. The behavioral tests measured were foot withdrawal thresholds (as an indicator of mechanical allodynia) in response to mechanical stimuli applied to the left hind paws. For each test, the animals were placed in a plastic

chamber (9×9×30 cm) and habituated for at least 10 minutes. The chamber was placed on top of a mesh screen, so that mechanical stimuli could be administered to the plantar surface of the left hind paws. Thresholds were determined by the up-down method⁷⁾ using a set of von Frey monofilaments (von Frey filament values : 3.65, 3.87, 4.10, 4.31, 4.52, 4.74, 4.92, and 5.16; equivalent to : 0.45, 0.74, 1.26, 2.04, 3.31, 5.50, 8.32, and 14.45 g values). Gram (g) means bending force of a set of von Frey monofilaments. A von Frey filament was applied perpendicularly to the most sensitive areas of the plantar surface at the center area of paw or the base of the third or fourth toes with sufficient force to bend the filament slightly for 3-4 seconds. An abrupt withdrawal of the foot during stimulation or immediately after stimulus removal was considered to be a positive response. The first stimulus was always the 4.31 filament. When there was a positive response, the next lower filament was used, and when no response was observed, the next higher filament was applied. This testing pattern continued until responses to the sixth von Frey stimuli from the first change of response (either higher or lower than the first stimulus depending on whether the first response was negative or positive) were measured. The responses were then converted to a 50% threshold value using the formula : $50\% \text{ threshold} = 10(X - kd) / 104$, where X is the value of the final von Frey hair used in log units, k is the tabular value for the pattern of positive or negative responses, and d is the mean differences between stimuli in log units (0.22). When positive or negative responses were still observed at the 3.65 or 5.16 filament, values of 0.3 or 18.6 g were assigned, respectively, by assuming a value of -0.5 for k in these cases.

Electrophysiological findings

A Nicolet Viking IV (Nicolet Biomedical Ins., USA) was used for recording the sensory evoked potential (SEP). The SEP is an objective and reliable electrophysiological measurement that reflects the status of the sensory pathways. All SEP responses were obtained from stimulation of an anesthetized, unrestrained rat. Rats were placed in the prone position on a plastic board with the active electrode implanted 2.5 mm posterior and 2.8




Fig. 1. A dorsal (A) and lateral (B) schematic of the rat skull with electrode placement for the SEP study. The arrow, triangle and circle indicate the bregma, lambda and recording electrode, respectively. SEP : sensory evoked potential.

mm laterally to the bregma and the reference electrode was implanted in the mid frontal bone (Fig. 1). The SEP responses were elicited by activation of the tibial nerve in the left leg. The ground electrode was placed subcutaneously between the stimulation and recording electrode. The left limb was stimulated by positive current pulses of 1.3-1.9 mA magnitude, 0.2 msec duration, and 3 Hz frequency. The signal-to-noise ratio was improved by ensemble averaging of 500 stimulus locked sweeps. The first peak latency was recorded. All animals were evaluated at day 1, 2, 3, 4, 8, 13, 17, and 21.

Statistical analysis

Statistical analyses were done using two-way repeated-measures analysis of variances with two-repeated factors followed by Tukey post hoc test for the experiment of Latin square design or two-way repeated-measures analysis of variances with one repeated time factor followed by Tukey post hoc tests. Results are presented as means \pm SEMs and analyzed using the Sigma Stat program (Systat Software, Inc., Chicago, IL, USA). In all cases, $p < 0.05$ was considered statistically significant.

RESULTS

Segmental spinal nerve ligation model

Mechanical allodynia threshold

The baseline mechanical threshold for all rats before operation was 17.1 g, which was the maximal cutoff point. We used the left hind paw as the site to measure the nociceptive threshold. Beginning at day 1, there was a significantly greater decrease in nociceptive threshold, a peak at 2 days and then a plateau until week 3, the final week of the study ($p < 0.05$) (Fig. 2).




Fig. 2. The time courses of mechanical threshold in L5, 6 spinal nerve ligation induced-neuropathy. Rolipram was injected intraperitoneally once a day. The vehicle group received an equal volume of normal saline. Note that spinal nerve ligation with vehicle application significantly decreased the mechanical threshold at the beginning of day 1. Rolipram significantly ameliorated the mechanical threshold decrease in comparison to the vehicle-injected group ($p < 0.05$). Asterisks indicate values significantly different ($p < 0.05$) from corresponding vehicle values using a two-way repeated measures analysis of variance with a repeated time factor, followed by the Tukey post hoc test.

SEP responses

Beginning on day 1, there was an increase in latency in the vehicle control group, and the increase was statistically significant after 2 days ($p < 0.05$) (Fig. 3).

Role of rolipram, specific PDE-4 inhibitor

We found that an intraperitoneal injection of the phosphodiesterase-4 inhibitor, rolipram, improved mechanical behavior and latency of SEP, compared with the vehicle-injected group ($p < 0.05$) (Fig. 2, 3).

DISCUSSION

The neurologic effects of numerous physical traumas and drugs are complex, encompassing both the central and peripheral nervous system^{28,34,37}. In the peripheral nervous system, the neurologic effect is generally small-fiber death and painful neuropathy^{2,6}. Over time, the pain far outweighs the analgesia, producing a neuropathic pain syndrome with symptoms that have been described as “like tearing flesh off the bones”³³. In the recent years, peripheral neuropathy due to anticancer drugs and ethanol consumption in western industrialized countries has increased among women with an associated increase in the rates of alcohol-related health problems^{8,11}. Thus, understanding the basis of clinical expression of neuropathy is an issue of growing importance and adequate treatment of this symptom may require therapeutic strategies. PDE4 inhibitors have been reported to reduce both proinflammatory cytokine levels, including TNF- α and IL-1 β ^{1,27}, molecules involved in free radical production and oxidative stress, such as iNOS and COX-2, as well as immune cell infiltration into the nervous system²⁹. Free radicals are derivatives of molecular oxygen and nitrogen, and in-




Fig. 3. The time courses of SEP in L5, 6 spinal nerve ligation induced-neuropathy. Rolipram was injected intraperitoneally once a day. The vehicle group received an equal volume of normal saline. Note that spinal nerve ligation with vehicle application significantly increased the latency at the beginning of day 1, peaked on day 2 and then reached a plateau until the final day of the study. Rolipram significantly ameliorated the latency increase compared with the vehicle-injected group ($p < 0.05$). Asterisks indicate values determined significantly different ($p < 0.05$) from the corresponding vehicle values using a two-way repeated measures analysis of variance with a repeated time factor, followed by the Tukey post hoc test. SEP : sensory evoked potential.

clude superoxide, hydroxyl radical, hydrogen peroxide, and peroxynitrite¹⁷). These molecules are ubiquitously present in the body and participate in many normal cellular processes including ion transport, transcription, neurotransmission, and neuromodulation¹⁷). Sources of free radicals include mitochondrial oxidative metabolism to produce adenosine triphosphate and several enzymes such as xanthine oxidase, phospholipase A2, cytochrome P450, monoamine oxidase, and tyrosine hydroxylase. Free radicals are normally removed by antioxidant systems including superoxide dismutase, catalase, glutathione, glutathione peroxidase, ascorbate, and α -tocopherol. Thus, their levels are precisely controlled by antioxidant systems. However, in pathologic conditions, levels of free radicals may increase due to the increased production or decreased antioxidants level^{9,10}). Two proinflammatory cytokines released after a central nervous system lesion are TNF- α and IL-1 β . Numerous studies have documented rapid increases in TNF- α and IL-1 β levels after traumatic brain and spinal cord injury^{15,35}). IL-1 β synergistically acts with TNF- α to induce nerve cell death. These proinflammatory cytokines stimulate inflammatory cells to release damaging reactive oxygen and nitrogen species, raise glutamate levels to excitotoxic levels, impair the ability of glia cells to buffer extracellular potassium, compromise the nervous system and attract inflammatory cells into the nervous system^{13,14,21,31,32}). Once initiated, the inflammatory cascade becomes a toxic positive-feedback loop, further exacerbating nervous system pathology. In the models of nervous system injury, several studies have demonstrated that restoration of cAMP levels improve outcome. In the animal model, application of rolipram to inhibit the degradation of cAMP promotes axon sparing and results in functional outcomes²⁵). In this study, we found that rolipram can prevent degradation of cAMP and improve mechanical behavior and latencies of SEP.

CONCLUSION

Our study shows that rolipram ameliorates pain behavior as measured by mechanical allodynia and SEP in an animal model of segmental spinal nerve ligation. Further, rolipram did not induce gastrointestinal problems or sedation at the given doses, so the behavioral changes are interpreted as analgesia. These results raise several questions for future research to identify the longterm effect and optimal dose of the PDE-4-specific inhibitor rolipram.

• Acknowledgements

The authors would like to thank Hee-Kee Kim, Associate Scientist, Dept. of Anesthesiology, University of Miami Miller school of Medicine, for his excellent behavioral and pharmaceutical advice.

References

1. Atkins CM, Oliva AA Jr, Alonso OF, Pearse DD, Bramlett HM, Dietrich WD : Modulation of the cAMP signaling pathway after traumatic brain

injury. *Exp Neurol* 208 : 145-158, 2007

2. Bosch EP, Pelham RW, Rasool CG, Chatterjee A, Lash RW, Brown L, et al. : Animal models of alcoholic neuropathy : morphologic, electrophysiologic, and biochemical findings. *Muscle Nerve* 2 : 133-144, 1979

3. Brain WR WJ : Disorders of peripheral nerves : Alcohol polyneuritis. In : *Brain's diseases of the nervous system*. London, 1969, pp817-819

4. Cogswell JP, Godlevski MM, Wisely GB, Clay WC, Leesnitzer LM, Ways JP, et al. : NF-kappa B regulates IL-1 beta transcription through a consensus NF-kappa B binding site and a nonconsensus CRE-like site. *J Immunol* 153 : 712-723, 1994

5. Corsetti G, Rezzani R, Rodella L, Bianchi R : Ultrastructural study of the alterations in spinal ganglion cells of rats chronically fed on ethanol. *Ultrastruct Pathol* 22 : 309-319, 1998

6. Dina OA, Barletta J, Chen X, Muterio A, Martin A, Messing RO, et al. : Key role for the epsilon isoform of protein kinase C in painful alcoholic neuropathy in the rat. *J Neurosci* 20 : 8614-8619, 2000

7. Dixon WJ : Efficient analysis of experimental observations. *Annu Rev Pharmacol Toxicol* 20 : 441-462, 1980

8. Fillmore KM : Women's drinking across the adult life course as compared to men's. *Br J Addict* 82 : 801-811, 1987

9. Floyd RA : Antioxidants, oxidative stress, and degenerative neurological disorders. *Proc Soc Exp Biol Med* 222 : 236-245, 1999

10. Floyd RA, Hensley K : Oxidative stress in brain aging. Implications for therapeutics of neurodegenerative diseases. *Neurobiol Aging* 23 : 795-807, 2002

11. Gombert ES : Women and alcohol : use and abuse. *J Nerv Ment Dis* 181 : 211-219, 1993

12. Hou S, Guan H, Ricciardi RP : Phosphorylation of serine 337 of NF-kappaB p50 is critical for DNA binding. *J Biol Chem* 278 : 45994-45998, 2003

13. Hu S, Peterson PK, Chao CC : Cytokine-mediated neuronal apoptosis. *Neurochem Int* 30 : 427-431, 1997

14. Keeling KL, Hicks RR, Mahesh J, Billings BB, Kotwal GJ : Local neutrophil influx following lateral fluid-percussion brain injury in rats is associated with accumulation of complement activation fragments of the third component (C3) of the complement system. *J Neuroimmunol* 105 : 20-30, 2000

15. Kinoshita K, Chatzipanteli K, Vitarbo E, Truettner JS, Alonso OF, Dietrich WD : Interleukin-1beta messenger ribonucleic acid and protein levels after fluid-percussion brain injury in rats : importance of injury severity and brain temperature. *Neurosurgery* 51 : 195-203; discussion 203, 2002

16. Koike H, Iijima M, Sugiura M, Mori K, Hattori N, Ito H, et al. : Alcoholic neuropathy is clinicopathologically distinct from thiamine-deficiency neuropathy. *Ann Neurol* 54 : 19-29, 2003

17. Lander HM : An essential role for free radicals and derived species in signal transduction. *FASEB J* 11 : 118-124, 1997

18. Lee I, Kim HK, Kim JH, Chung K, Chung JM : The role of reactive oxygen species in capsaicin-induced mechanical hyperalgesia and in the activities of dorsal horn neurons. *Pain* 133 : 9-17, 2007

19. Manganiello VC, Murata T, Taira M, Belfrage P, Degerman E : Diversity in cyclic nucleotide phosphodiesterase isoenzyme families. *Arch Biochem Biophys* 322 : 1-13, 1995

20. Mayr B, Montminy M : Transcriptional regulation by the phosphorylation-dependent factor CREB. *Nat Rev Mol Cell Biol* 2 : 599-609, 2001

21. Meda L, Cassatella MA, Szendrei GI, Otvos L Jr, Baron P, Villalba M, et al. : Activation of microglial cells by beta-amyloid protein and interferon-gamma. *Nature* 374 : 647-650, 1995

22. Monforte R, Estruch R, Valls-Solé J, Nicolás J, Villalta J, Urbano-Marquez A : Autonomic and peripheral neuropathies in patients with chronic alcoholism. A dose-related toxic effect of alcohol. *Arch Neurol* 52 : 45-51, 1995

23. Naik AK, Tandan SK, Dudhgaonkar SP, Jadhav SH, Kataria M, Prakash VR, et al. : Role of oxidative stress in pathophysiology of peripheral neuropathy and modulation by N-acetyl-L-cysteine in rats. *Eur J Pain* **10** : 573-579, 2006
24. Narita M, Miyoshi K, Suzuki T : Involvement of microglia in the ethanol-induced neuropathic pain-like state in the rat. *Neurosci Lett* **414** : 21-25, 2007
25. Nikulina E, Tidwell JL, Dai HN, Bregman BS, Filbin MT : The phosphodiesterase inhibitor rolipram delivered after a spinal cord lesion promotes axonal regeneration and functional recovery. *Proc Natl Acad Sci U S A* **101** : 8786-8790, 2004
26. Padi SS, Kulkarni SK : Minocycline prevents the development of neuropathic pain, but not acute pain : Possible anti-inflammatory and antioxidant mechanisms. *Eur J Pharmacol* **601** : 79-87, 2008
27. Pearse DD, Pereira FC, Marcillo AE, Bates ML, Berrocal YA, Filbin MT, et al. : Camp and schwann cells promoted axonal growth and functional recovery after spinal cord injury. *Nat Med* **10** : 610-616, 2004
28. Pentney RJ, Quackenbush LJ : Dendritic hypertrophy in Purkinje neurons of old Fischer 344 rats after long-term ethanol treatment. *Alcohol Clin Exp Res* **14** : 878-886, 1990
29. Schaal SM GM, Golshani R, Ghosh M, Lovera L, Lopez M, Patel M, et al. : Targeting phosphodiesterase-4 after spinal cord injury using pharmacological and molecular approaches. *J Neurotrauma* **25** : 297, 2008
30. Sharma SS, Sayyed SG : Effects of trolox on nerve dysfunction, thermal hyperalgesia and oxidative stress in experimental diabetic neuropathy. *Clin Exp Pharmacol Physiol* **33** : 1022-1028, 2006
31. Soares HD, Hicks RR, Smith D, McIntosh TK : Inflammatory leukocytic recruitment and diffuse neuronal degeneration are separate pathological processes resulting from traumatic brain injury. *J Neurosci* **15** : 8223-8233, 1995
32. Tanaka M, Sotomatsu A, Yoshida T, Hirai S, Nishida A : Detection of superoxide production by activated microglia using a sensitive and specific chemiluminescence assay and microglia-mediated PC12h cell death. *J Neurochem* **63** : 266-270, 1994
33. Vergheze MW, McConnell RT, Strickland AB, Gooding RC, Stimpson SA, Yarnall DP, et al. : Differential regulation of human monocyte-derived TNF alpha and IL-1 beta by type IV cAMP-phosphodiesterase (cAMP-PDE) inhibitors. *J Pharmacol Exp Ther* **272** : 1313-1320, 1995
34. Victor M : **Polyneuropathy due to nutritional deficiency and alcoholism**. Philadelphia : WB Saunder's Co., 1975
35. Vitarbo EA, Chatzipanteli K, Kinoshita K, Truettner JS, Alonso OF, Dietrich WD : Tumor necrosis factor alpha expression and protein levels after fluid percussion injury in rats : the effect of injury severity and brain temperature. *Neurosurgery* **55** : 416-424; discussion 424-425, 2004
36. Yerdelen D, Koc F, Uysal H : Strength-duration properties of sensory and motor axons in alcoholic polyneuropathy. *Neurol Res* **30** : 746-750, 2008
37. Zou J, Rabin RA, Pentney RJ : Ethanol enhances neurite outgrowth in primary cultures of rat cerebellar macroneurons. *Brain Res Dev Brain Res* **72** : 75-84, 1993