

Limited mandibular movements after removal of the mandibular third-molar: use of the anterior bite plane and complementary therapies

Marcelo Palinkas^{1,*}, Regina Maura Arantes Nassar², Mariangela Salles Pereira Nassar¹, Solange Aparecida Bataglion², César Bataglion¹, Cássio Edvard Sverzut³, Takami Hirono Hotta⁴, Simone Cecilio Hallak Regalo²

¹Department of Dentistry, Ribeirão Preto Dental School, University of São Paulo, Ribeirão Preto, SP, Brazil; ²Department of Morphology, Stomatology and Physiology, Ribeirão Preto Dental School, University of São Paulo, Ribeirão Preto, SP, Brazil; ³Department of Oral and Maxillofacial Surgery and Periodontology; ⁴Department of Dental Materials and Prosthodontics, Ribeirão Preto Dental School, University of São Paulo; Ribeirão Preto, SP, Brazil

ABSTRACT

The traumatic removal of the mandibular third molar may promote post surgical consequences such as orofacial pain and limited mandibular movements. The aim of this case report is to describe the use of an anterior bite plane and complementary therapies (low level laser therapy and acupuncture) to treat the muscular dysfunction and the painful symptoms. A 33 year-old male patient who had a severe malocclusion and signs and symptoms of temporomandibular dysfunction was submitted to an initial clinical examination: electromyography of the masticatory muscles and IRM of the temporomandibular joint. After treatments, the results showed reduced pain symptoms and an increase of the mandibular movements and adequated electromiographic activities. The authors concluded that the combination therapies may be used as an alternative treatment because it satisfied the functional requirements of the patient and provided an asymptomatic clinical condition.

Keywords mandibular movements, anterior bite plane, acupuncture, low level laser therapy, electromyography

INTRODUCTION

Temporomandibular disorders (TMD) is the term used to refer to dysfunctions characterized by pain in the region of the temporomandibular joints, muscles of mastication, and periauricular area, limitations and deviations in the mandibular movements, joint noises and an altered occlusal relation (Dworkin et al., 1990).

The etiology of TMD has multifactorial factors related to the development of TMD including the loss of posterior occlusal support with a consequent alteration in the vertical dimensions of occlusion, muscular hyperactivity and adjacent structures, postural changes, occlusal relation, emotional stress, and oral habits (LeResche et al., 1997; Hotta et al., 2003). The third molar surgery also could contribute to the number of patients with chronic orofacial pain (Juhl et al., 2009) and has been associated with a risk of TMD (Plesh et al., 1999; Huang et al., 2002; Huang et al., 2006; Huang et al., 2008) although this surgery is the most common procedure performed by oral surgeons (Amarillas-Escobar et al., 2010).

Treatment of TMD is basically odontologic and may involve anterior splints (Torii and Chiwata, 2010) occlusal splints, splints with posterior occlusal support, occlusal adjustment, removable therapeutic partial prostheses (Hotta et al., 1997), oral rehabilitation or orthodontic treatment, although therapeutic support regimens in the areas of psychology, phonoaudiology, and physical therapy may be associated

depending on the needs of each patient (Hotta et al., 2003). Acupuncture (Cho and Whang, 2010; Hotta et al., 2010) and low level laser therapy (Mazzetto et al., 2010) have been used to treat limited mandibular movements and particularly painful symptoms of the masticatory muscles and temporomandibular joint.

Acupuncture is a therapeutic method based on Traditional Chinese Medicine, which consists of the insertion of small, solid needles, usually made of stainless steel, into specific body points in order to improve health or modify painful states. This therapy was shown to be effective in patients with chronic pain and has been used to prevent and treat several disorders, having become a treatment option in dentistry by promoting antiinflammatory and analgesic actions (Smith et al., 2007; Rancan et al., 2009).

Electromyographical analysis of the masseter and temporalis muscles may be used to compare the muscular activities during several mandibular positions and clinical conditions (Hotta et al., 2010).

This article reports a clinical case in which a combination of physical therapy and odontologic resources were used to treat limited mandibular movements and painful symptoms after third molar surgical removal.

METHODS AND RESULTS

This clinical case was conducted after the approval of the Ethics Committee (process number 2006 1971.58.5). Before signing the informed consent, the subject was informed of the study procedures by the investigators. The study was conducted at the School of Dentistry of Ribeirão Preto, University of São Paulo, Brazil.

*Correspondence: Marcelo Palinkas

E-mail: schregalo@forp.usp.br

Received July 13, 2011; Accepted February 13, 2012; Published February 29, 2012

doi: <http://dx.doi.org/10.5667/tang.2011.0020>

© 2012 by Association of Humanitas Medicine



Fig. 1. Linguoersion of the right posterior mandibular teeth.



Fig. 2. Anterior bite plane.



Fig. 3. Low level laser intensity.



Fig. 4. Session of acupuncture.

A 33-year-old man was presented for dental treatment, complaining of difficulty in opening his mouth after extraction of his inferior right third-molar.

Clinical examination showed complete natural dentition with linguoersion of the right posterior mandibular teeth, elongated face, retrognathism (Fig. 1), reduced opening (5 mm, from edge to edge) and masticatory muscle pain. Given the clinical findings, the patient was submitted to electromyography analysis of the masticatory muscles (rest and maximum voluntary contraction positions), before and after treatments (Table 1). These analyses were performed using a MyoSystem- Br1 (São Paulo, Brazil) electromyographer with differential active electrodes (silver bars 10 mm apart, 10 mm long, 2 mm wide, 20 gain, input impedance 10 GX and 130 dB at 60 Hz common mode rejection ratio). Impedance is a function of two components (vectors): the resistance of the tissues themselves, and the additional opposition (reactance) due to the capacitance of membranes, tissue interfaces, and nonionic tissues. The measured resistance is approximately equivalent to that of muscle tissue.

Surface differential active electrodes were placed on the skin, previously cleaned with alcohol, bilaterally on both masseter muscles and temporalis. A ground electrode was also used and fixed on the skin over the frontal region. EMG signals were analogically amplified with a gain of 1000, filtered by a pass-band of 0.01 – 1.5 kHz and sampled by a 12 bit A/D converter with a 2 kHz sampling rate. The signals were digitally filtered by a passband filter of 10 – 500 Hz for data processing (Land and Peregrina, 2003; Cecilio et al., 2010).

After this, the patient was referred to urgency therapy with an anterior bite plane, low level laser intensity and acupuncture

procedures. The use of the anterior bite plane was indicated by two days, uninterrupted (Fig. 2) (Lievens, 1988). The laser apparatus was the GaAlAs laser Twin Laser (MM Optics, São Carlos – SP, Brazil) that operates with a continuous laser beam (780 nm wavelength; 70 mW power output, 60 J/cm², 10 sessions) (Fig. 3) (Cetiner et al., 2006). Finally, the patient was submitted to 5 sessions of acupuncture point applications: Hn3, Ig4 (bi), F3 (bi), E7 (bi), E6 (bi), SJ17 (bi), Vb20 (bi), P7, C7, Vb14 (bi), Id18 (bi), E36 (bi), R3 e Bp6 (bi). The depths of needles insertion were just tissue level (Fig. 4)

Complementary exams were requested and magnetic resonance imaging of the temporomandibular joints was obtained in both closed and open mouth positions. The results showed that the articular disks were well positioned in the mandibular fossae.

The results of the urgency approach showed an increased mouth opening (5 mm to 40 mm), reduction of the dolorous symptoms and reestablishment of the normal stomathognathic system functions such mastication, speech and deglutition.

DISCUSSION

Surgical removal of the third-molar is the most common procedure performed by oral surgeons (Amarillas-Escobar et al., 2010). Some surgeons, even with the existence of risks, often advocate third molar removal during the teenage years (Huang et al., 2008), citing advantages such as incompletely formed roots, better healing, and less morbidity (Phillips et al., 2003).

According to Battista et al. (2007) the surgical trauma of this procedure causes postoperative pain, swelling and trismus.

Table 1. Mean values of electromyographic activities (microvolts/second) of masseter and anterior temporalis muscles during rest position and habitual occlusion, before (1) and after (2) treatments

Muscles	Rest Position				Habitual Occlusion			
	1		2		1		2	
	Right	Left	Right	Left	Right	Left	Right	Left
Masseter	6,31	13,62	5,84	10,96	9,19	25,1	25,56	44,89
Temporalis	8,28	11,84	6,77	8,03	14,97	45,08	27,18	68,19

In the present clinical report, a severe limited open mouth occurred after the surgical removal of right maxillary third-molar in an adult patient and persisted for 56 days.

As the use of low level laser therapy (LLLT) for the treatment of musculoskeletal pain syndromes has become common (Kulekcioglu et al., 2003; Grossi et al., 2007; Mazzetto et al., 2007) and the lasertherapy promotes cellular bio-stimulation, accelerates tissue regeneration and wound healing and decreases pain and swelling (Coluzzi and Convissar, 2004), the authors of the present study selected this therapy to be applied in association with an anterior bite plane (occlusal deprogramming) and acupuncture applications, looking for better results.

Besides the remission of the dolorous symptoms and a significant increase in the open mouth, the subject also noted a reduction in his anxiety symptoms. This result also was shown in an earlier study (Hotta et al., 2010) probably due the acupuncture point IG4.

With reference to electromyographic activities, the results showed that the mean values were changed by the combination therapies and these results were better than those presented by the patient in the initial phase of the treatment in both mandibular positions, rest and habitual occlusion.

In the initial phase, probably due the dolorous symptoms and also the linguoversion of the right posterior mandibular teeth, in habitual occlusion, the electromyographical activities of the masticatory muscles were smaller on this side. After the reduction of the symptoms, the electromyographical activities became stronger, mainly due to the anti-inflammatory action of the laser (Siéssere et al., 2011). On the other hand, in the resting position, the results showed a decreasing of the electromyographical activities as expected (Table 1).

Thus, the authors concluded that, in this case report, the combination of selected therapies was efficient to promote the satisfactory clinical result.

ACKNOWLEDGEMENTS

This research was supported by Ribeirão Preto Dental School, University of São Paulo.

CONFLICT OF INTEREST

The authors have no conflicting financial interests.

REFERENCES

Amarillas-Escobar ED, Toranzo-Fernández JM, Martínez-Rider R, Noyola-Frías MA, Hidalgo-Hurtado JA, Serna VM, Gordillo-Moscoso A, Pozos-Guillén AJ. Use of therapeutic laser after surgical removal of impacted lower third molars. *J Oral Maxillofac Surg.* 2010;68:319-324.

Cecílio FA, Regalo SC, Palinkas M, Issa JP, Siéssere S, Hallak JE, Machado-de-Sousa JP, Semprini M. Ageing and surface EMG activity patterns of masticatory muscles. *J Oral Rehabil.* 2010;37:248-255.

Cetiner S, Kahraman SA, Yuceta S. Evaluation of lowlevel laser therapy in the treatment of temporomandibular disorders. *Photomed Laser Surg.* 2006;24:637-641.

Cho SH, Whang WW. Acupuncture for temporomandibular disorders: a systematic review. *J Orofac Pain.* 2010;24:152-162.

Coluzzi DJ, Convissar RA. Laser in clinical dentistry. *Dent Clin North Am.* 2004;48:10-12.

Dworkin SF, Huggins KH, LeResche L, Von Korff M, Howard J, Truelove E, Sommers E. Epidemiology of signs and symptoms in temporomandibular disorders: clinical signs in cases and controls. *J Am Dent Assoc.* 1990;120:273-281.

Grossi GB, Maiorana C, Garramone RA, Borgonovo A, Creminelli L, Santoro F. Assessing postoperative discomfort after third molar surgery: a prospective study. *J Oral Maxillofac Surg.* 2007;65:901-917.

Hotta PT, Hotta TH, Bataglion C, Bataglion SA, de Souza Coronatto EA, Siéssere S, Regalo SC. Emg analysis after laser acupuncture in patients with temporomandibular dysfunction (TMD). Implications for practice. *Complement Ther Clin Pract.* 2010;16:158-160.

Hotta TH, Bataglion A, Bataglion C, Bezzon OL. Involvement of dental occlusion and trigeminal neuralgia: a clinical report. *J Prosthet Dent.* 1997;77:343-345.

Hotta TH, Vicente MF, dos Reis AC, Bezzon OL, Bataglion C, Bataglion A. Combination therapies in the treatment of temporomandibular disorders: A clinical report. *J Prosthet Dent.* 2003;89:536-539.

Huang GJ, Drangsholt MT, Rue TC, Cruikshank DC, Hobson KA. Age and third molar extraction as risk factors for temporomandibular disorder. *J Dent Res.* 2008;87:283-287.

Huang GJ, LeResche L, Critchlow CW, Martin MD, Drangsholt MT. Risk factors for diagnostic subgroups of painful temporomandibular disorders (TMD). *J Dent Res.* 2002;81:284-288.

Huang GJ, Rue TC. Third-molar extraction as a risk factor for temporomandibular disorder. *J Am Dent Assoc.* 2006;137:1547-1554.

Juhl GI, Jensen TS, Norholt SE, Svensson P. Incidence of symptoms and signs of TMD following third molar surgery: a controlled, prospective study. *J Oral Rehabil.* 2009;36:199-209.

- Kulekcioglu S, Sivrioglu K, Ozcan O, Parlak M. Effectiveness of low-level laser therapy in temporomandibular disorder. *Scand J Rheumatol.* 2003;32:114-118.
- Land MF, Peregrina A. Anterior deprogramming device fabrication using a thermoplastic material. *J Prosthet Dent.* 2003;90:608-610.
- LeResche L, Saunders K, Von Korff MR, Barlow W, Dworkin SF. Use of exogenous hormones and risk of temporomandibular disorder pain. *Pain.* 1997;69:153-160.
- Lievens P. Effects of laser treatment on the lymphatic system and wound healing. *Laser.* 1988;1:12-13.
- Mazzetto MO, Carrasco TG, Bidinelo EF, Pizzo RCA, Mazzetto RG. Low intensity laser application in temporomandibular disorders: a phase I double blind study. *Cranio.* 2007;25:186-192.
- Mazzetto MO, Hotta TH, Pizzo RAC. Measurements of jaw movements and TMJ pain intensity in patients treated with GaAlAs laser. *Braz Dent J.* 2010;21:356-360.
- Phillips C, White RP Jr, Shugars DA, Zhou X. Risk factors associated with prolonged recovery and delayed healing after third molar surgery. *J Oral Maxillofac Surg.* 2003;61:1436-1448.
- Plesh O, Gansky SA, Curtis DA, Pogrel MA. The relationship between chronic facial pain and a history of trauma and surgery. *Oral Surg Oral Med Oral Pathol Oral Radiol Endod.* 1999;88:16-21.
- Rancan SV, Bataglioni C, Bataglioni SA, Bechara OM, Semprini M, Siéssere S, Sousa JP, Crippa JA, Hallak JE, Regalo SC. Acupuncture and temporomandibular disorders: a 3-month follow-up EMG study. *J Altern Complement Med.* 2009;15:1307-1310.
- Siéssere S, de Sousa LG, Issa JP, Iyomasa MM, Pitol DL, Barbosa AP, Semprini M, Sebald W, Bentley MV, Regalo SC. Application of low-level laser irradiation (LLL) and rhBMP-2 in critical bone defect of ovariectomized rats: histomorphometric evaluation. *Photomed Laser Surg.* 2011;29:453-458.
- Smith P, Moss crop D, Davies S, Sloan P, Al-Ani Z. The efficacy of acupuncture in the treatment of temporomandibular joint myofascial pain: a randomised controlled trial. *J Dent.* 2007;35:259-267.
- Torii K, Chiwata I. A case report of the symptom-relieving action of an anterior flat plane bite plate for temporomandibular disorder. *Open Dent J.* 2010;21:218-222.