

RESEARCH ARTICLE

Vitexicarpin Induces Apoptosis in Human Prostate Carcinoma PC-3 Cells through G2/M Phase Arrest

Fan-Min Meng¹, Jing-Bo Yang², Chun-Hui Yang³, Yu Jiang¹, Yong-Feng Zhou¹, Bo Yu¹, Hong Yang^{1*}

Abstract

Vitexicarpin (3', 5-dihydroxy-3, 4', 6, 7-tetramethoxyflavone), a polymethoxyflavone isolated from *Vitex Fructus* (*Vitex rotundifolia* Linne fil.), has long been used as an anti-inflammatory herb in traditional Chinese medicine. It has also been reported that vitexicarpin can inhibit the growth of various cancer cells. However, there is no report elucidating its effect on human prostate carcinoma cells. The aim of the present study was to examine the apoptotic induction activity of vitexicarpin on PC-3 cells and molecular mechanisms involved. MTT studies showed that vitexicarpin dose-dependently inhibited growth of PC-3 cells with an IC_{50} ~28.8 μ M. Hoechst 33258 staining further revealed that vitexicarpin induced apoptotic cell death. The effect of vitexicarpin on PC-3 cells apoptosis was tested using propidium iodide (PI)/Annexin V-FITC double staining and flow cytometry. The results indicated that vitexicarpin induction of apoptotic cell death in PC-3 cells was accompanied by cell cycle arrest in the G2/M phase. Furthermore, our study demonstrated that vitexicarpin induction of PC-3 cell apoptosis was associated with upregulation of the proapoptotic protein Bax, and downregulation of antiapoptotic protein Bcl-2, release of Cytochrome c from mitochondria and decrease in mitochondrial membrane potential. Our findings suggested that vitexicarpin may become a potential leading drug in the therapy of prostate carcinoma.

Keywords: Vitexicarpin - PC - 3 cells - anticancer - apoptosis - G2/M phase arrest

Asian Pacific J Cancer Prev, **13** (12), 6369-6374

Introduction

Prostate cancer is the second most common diagnosed cancer after lung cancer worldwide, and it is the third most common cause of cancer deaths in developed countries. In 2011, 240,890 men were diagnosed with prostate cancer and 33,720 men died of it as estimated by The American Cancer Society 2012. Prostate cancer incidence and mortality is comparatively higher in developed countries. Treatment of prostate cancer depends on age of patients, overall health of individual, and the stage of disease. Current available therapies include active surveillance, surgery, radiation therapy, hormone therapy, chemotherapy, and immunotherapy. However, because prostate cancer can recur in an androgen-insensitive or hormone-refractory form that is not responsive to current therapies, the mortality rate associated with recurrent cases is high (American Cancer Society, 2012). There is great demand for effective novel therapeutic agents.

Vitexicarpin (3', 5-dihydroxy-3, 4', 6, 7-tetramethoxyflavone), a polymethoxyflavone isolated from *Vitex Fructus* (*Vitex rotundifolia* Linne fil.) has long been used as an anti-inflammatory herb in traditional

Chinese medicine (Choudhary et al., 2009; Mesaik et al., 2009). A number of recent pharmacological reports indicated that vitexicarpin induces growth inhibition, cell cycle arrest, and apoptosis in various cancer cells including the human cervical cancer cell line Hela (Chen et al., 2011; Zeng et al., 2012), hepatocellular carcinoma cell line HepG2 (Yang et al., 2011; He et al., 2012), lung epithelial cell line A549 (Koh et al., 2011), and leukemic cell line K562 (Shen et al., 2009). Molecular mechanism studies showed that vitexicarpin induces apoptosis via activation of c-Jun N-terminal kinase (JNK) in Hela cells (Zeng et al., 2012), inactivation NF- κ B in A549 cells (Koh et al., 2011), Caspase-3 activation in K562 cells (Shen et al., 2009), etc. So far, there is no information available on its effect on human prostate carcinoma cells. In the present study, we investigated effect of vitexicarpin on apoptosis in human prostate cancer cell line PC-3 using cytotoxicity assay, cell cycle analysis, and western blotting method.

Materials and Methods

Chemicals and reagents

Vitexicarpin was purchased from the National Institutes

¹School of life sciences, Liaoning Provincial Key Laboratory of Biotechnology and Drug Discovery, Liaoning Normal University, ²College of Laboratory Medicine, Dalian Medical University, Dalian, ³Central Research Laboratory, Jilin University Bethune Second Hospital, Changchun, China *For correspondence: hyanglnnu@gmail.com

for Food and Drug Control (Beijing, China). Characterized grade fetal bovine serum (FBS) was purchased from HyClone (USA). Dimethyl sulfoxide (DMSO) was purchased from Sangon Biontech (Shanghai, China) Co., Ltd. Dulbecco's Modified Eagle's Medium (DMEM), [3-(4,5-Dimethylthiazol-2-yl)-2,5-Diphenyltetrazolium Bromide] (MTT), trypsin, Hoechst 33258, Rhodamine 123, penicillin and streptomycin were purchased from Sigma (Beijing, China). Annexin V-FITC Apoptosis Detection Kit, Cell Cycle and Apoptosis Analysis Kit, Reactive Oxygen Species Assay Kit and BCA Protein Assay kit were purchased from Beyotime Institute of Biotechnology (Shanghai, China). Rabbit anti-human Caspase-3, Bax, Cyclin B1, Cytochrome c, mouse anti-human Bcl-2, β -actin, cyclin-dependent kinase (CDK1) primary antibodies were purchased from Cell Signaling (China). Horseradish peroxidase-conjugated secondary antibodies (Anti-mouse and anti-rabbit) were purchased from Santa Cruz (Beijing, China).

Cell culture

Human prostate cancer PC-3 cell line was purchased from American Type Culture Collection (ATCC, China) and was maintained in our lab. The cells were cultured in DMEM (supplemented with 10% FBS, 100 U/ml penicillin, and 100 μ g/ml streptomycin) at 37 °C, in a CO₂ incubator (5% CO₂ and 95% air, 95% humidity). Newly thawed PC-3 cells were past at least 3 passages before tests.

Freshly isolated mouse splenocytes were prepared from Kunming mice of SPF grade (Central Research Laboratory, Jilin University Bethune Second Hospital, China). The cells were cultured in RPMI 1640 medium supplemented with 10% (v/v) FBS and maintained in a CO₂ incubator (5% CO₂ and 95% air, 95% humidity).

Cell proliferation assay

Effect of vitexicarpin on viability of PC-3 cells was evaluated by MTT assay as reference (Khan et al., 2012). Briefly, PC-3 cells were subcultured in 96-well plates at a density of 10⁵/well and incubated in CO₂ incubator. After ~12h incubation, the cells were exposed to various concentrations of vitexicarpin (3.125, 6.25, 12.5, 25, 50, 100, and 200 μ M) and continuously incubated for 24 h. Following incubation, 20 μ l MTT (5 mg/ml) was added in each well and plate was again incubated at 37 °C. After 4 h, medium was removed carefully and 150 μ l DMSO was added to each well. The plate was shaken gently for 15 min to dissolve formazan crystals. The absorbance nm was recorded at 570 by the microplate reader (Thermo Scientific, Vario Skan Flash, USA). Percentage cell viability was calculated as follow:

$$\text{Cell viability (\%)} = (\text{A570 treated} / \text{A570 control}) \times 100$$

The IC₅₀ Values were calculated using GraphPad Prism 5.

Cell morphological changes

Cells were treated with vitexicarpin for 24 h, and then morphological changes of the cells were observed under a fluorescence microscope (Olympus 1 \times 71, Japan), recorded with a CCD camera (Olympus DP72, Japan) and analyzed

using DP2-BSW software.

Flow cytometry analysis of apoptosis

Apoptosis of PC-3 cells were determined using Annexin V-FITC/PI double staining Apoptosis Detection Kit. PC-3 cells were cultured in 6-well plates, incubated overnight, and then exposed to different concentrations of vitexicarpin for 24h. The cells were trypsinized, washed twice with ice-chilled PBS and collected through centrifugation (1000rpm for 5 min, 4 °C). The cell pellets were resuspended in binding buffer and stained sequentially with Annexin V-FITC and PI as manufacture's instructions. Flow cytometry was performed on Beckman Coulter, Epics XL (USA) and data were recorded using Cell Quest software.

DNA fragmentation analysis by Hoechst 33258 staining

The apoptosis in PC-3 cells was detected by using Hoechst 33258 nuclei staining. Cells were treated with vitexicarpin for 24h and collected by centrifugation at 1000 rpm for 5min. After centrifugation, the cells were washed with PBS, stained with Hoechst 33258 (50 μ g/mL) and incubated at 37 °C for 30 min in the dark. At the end of incubation, cells were washed and resuspended in PBS for the observation of nuclear morphology under fluorescence microscope (Olympus 1 \times 71, Japan), recorded with a CCD camera (Olympus DP72, Japan) and analyzed using DP2-BSW software. Apoptotic cells were defined as cells showing nuclear shrinkage and chromatin condensation.

Flow cytometric analysis of cell cycle

PC-3 cell cycle phase distribution was determined by using PI and flow cytometry as described previously (Khan et al., 2012). After the PC-3 cells were incubated with vitexicarpin for 24 h, the cells were collected and fixed in 70% ice-cold ethanol overnight, rinsed twice with PBS, and then stained with 50 μ g/ml PI (containing 100 μ g/ml RNase A) at 4 °C for 30 min in the dark. The DNA contents were assayed on Beckman Coulter, Epics XL (USA) using Cell Quest software.

Determination of intracellular Reactive Oxygen Species (ROS) generation

Intracellular ROS generations were determined using 2',7'-dichlorofluorescein-diacetate (DCFH-DA) staining method. PC-3 cells cultured in 6-well plates were treated with vitexicarpin for 24 h, and then stained with 10 μ mol/L DCFH-DA at 37 °C for 30 min as the manufacture's instructions. The cells were then collected, rinsed 3 times with PBS, resuspended in 200 μ l PBS. 2',7'-dichlorofluorescein (DCF) fluorescence data were acquired on Beckman Coulter, Epics XL (USA).

Determination of mitochondrial transmembrane potential ($\Delta\Psi_m$)

Rhodamine 123 staining method was used to evaluate $\Delta\Psi_m$ in PC-3 cells. PC-3 cells cultured in 6-well plates were exposed to different concentrations of vitexicarpin for 24 h. The cells were collected in centrifuge tube, resuspended in 500 μ l PBS, and then stained with 10 μ g/ml rhodamine 123 at 37 °C for 30 min in darkroom. After




Figure 1. Effects of Vitexicarpin on Viabilities and Morphological Characteristics of PC-3 Cells. (A) Chemical structure of vitexicarpin. (B) PC-3 cells were treated with indicated concentrations of vitexicarpin for 24 h. Cell survival rates were measured using MTT assay. Data are expressed as Mean \pm SE of three independent experiments from similar results. Morphological changes of PC-3 cells (C) and mouse splenocytes (D) observed under the light microscopy after treated with indicated concentrations of vitexicarpin or DMSO (negative control) for 24 h. Scale bar=50µm

staining, the cells were washed 3 times with PBS, and then resuspended in 200 µl PBS. Rhodamine 123 fluorescence was measured on flow cytometry (Emission wavelength λ_{em} =488, Excitation wavelength λ_{ex} =530). Details were described in reference (Ji et al., 2011).

Western blotting analysis

To elucidate the mechanism of the apoptotic effect of vitexicarpin, we analyzed the apoptosis-related proteins in PC-3 cells. After incubated with (0, 30, and 50 µM) vitexicarpin for 24 h, PC-3 cells were collected, washed twice with PBS, and lysed with lysis buffer. The lysate was centrifuged at 12000 rpm for 15min to get rid of insoluble components. For cytochrome c expression in cytosol, cytosolic fractions were extracted as described previously (Khan et al., 2012). Protein concentrations of the supernatants were determined using bicinchoninic acid (BCA) Protein Assay Kit. ~40 µg of proteins were subjected to SDS-PAGE and transferred to polyvinylidene fluoride (PVDF) membranes (Bio-Rad Laboratories, Inc, USA) followed by Western blotting according to standard protocols. The polyvinylidene fluoride (PVDF) membranes were sequentially blocked with 5% (w/v) nonfat milk for 2h at room temperature, washed with tris-buffered saline-tween solution (TBST). Membranes were then incubated overnight at 4°C with Caspased-3 (1:1000), Bax (1:1000), Cytochrome c (1:1000), Cyclin B1 (1:2000), Bcl-2 (1:500), β -actin (1:500), or cyclin-dependent kinase 1(CDK1) (1:300), respectively. Finally, the PVDF membranes were incubated with horseradish peroxidase (HRP) conjugated secondary antibodies and

signals were detected using ECL Plus Chemiluminescence Kit on X-ray film (Millipore Corporation, Billerica, USA).

Statistical analysis of data

All data are expressed as Mean \pm SEM. One way ANOVA test was used to compare control group and test values. Student's t-test was used to determine significance when only two groups were compared, p values less than 0.05 were considered statistically significant.

Results

Vitexicarpin Reduces Cell Viability and Induces Cell Death in human prostate carcinoma PC-3 cells

Effect of vitexicarpin (chemical structure shown in Figure 1A) on human prostate carcinoma PC-3 cell viabilities were determined using MTT assay. PC-3 cells were incubated with series concentrations of vitexicarpin for 24h. As shown in Figure 1B, vitexicarpin dose-dependently inhibited proliferation of the PC-3 cells with IC₅₀ ~ 28.78 µM. Maximum inhibition activity was achieved by 200 µM vitexicarpin. Cell morphological changes were observed under a light microscope. Cells were treated with 30 and 50 µM vitexicarpin for 24 h. As compared to DMSO control group, vitexicarpin treated group showed a significant decrease in adherent cells which accompanied an increase in floating cells in culture medium. This study also revealed significant changes in cell morphology: in contrast to normal polyclonal structure of normal cells, the vitexicarpin treated PC-3 cells acquired a round and shrunken shape (Figure 1C). We also examined the effect of vitexicarpin on mouse splenocytes using trypan blue method; it showed that vitexicarpin had little toxic effect on the cells (Figure 1D).

Vitexicarpin Induces Apoptotic Cell Death in human prostate carcinoma PC-3 cells

Cell death can be induced by apoptosis and/or necrosis. Apoptotic cell death is characterized by major features like DNA fragmentation and loss of plasma membrane integrity (Schafer, 1998). Hoechst 33258 staining analysis showed that 10, 30 and 50 µM vitexicarpin significantly increased the percentage of cleaved nuclei from 3.96 % (in DMSO control group) to 12.44%, 27.83%, and 54.65 % (Figure 2A). To further investigate vitexicarpin-induced apoptotic effect, PC-3 cells were treated with different concentrations of vitexicarpin for 24 h. The cells were double stained with Annexin V-FITC/PI and analyzed on flow cytometry as described in the Materials and Methods Section. As shown in Figure 2B, treatment of cells with 10, 30, and 50 µM vitexicarpin significantly increased apoptosis rate from 2.07% \pm 0.1925 to 3.87% \pm 0.3031 5.62% \pm 0.3964 and 25.27% \pm 1.1993 respectively.

Vitexicarpin Induces Mitotic Arrest in human prostate carcinoma PC-3 cells

Cell cycle arrest is also a major cause of cell proliferation inhibition. In order to elucidate whether vitexicarpin-induced PC-3 cell growth inhibition is associated with mitotic arrest, cell cycle was profiled by PI staining and flow cytometry method. PC-3 cells were




Figure 2. Apoptosis Induced by Vitexicarpin in PC-3 Cells. (A) Cells were treated with 0, 10, 30 and 50 μM vitexicarpin for 24 h respectively. Nuclear morphological changes were observed using Hoechst 33258 staining and fluorescence microscope. Arrows indicate the condensed and fragmented nuclei. Scale bar = 20 μm. Histograms show number of cleaved nuclei (apoptotic cells) counted microscopically from 100 nuclei. Data are expressed as Mean ± SE of three independent experiments with the similar results. **p<0.01 compared with the control. (B) PC-3 cells were treated with 0, 10, 30, and 50 μM vitexicarpin for 24 h. The cells were then stained with FITC-conjugated Annexin V and PI for flow cytometric analysis. X-axis and y-axis represents Annexin V-FITC and PI staining, respectively. Cell populations shown in the lower right (Annexin V+/PI-) represents apoptotic cells, upper right (Annexin V+/PI+) represents necrotic cells. Data expressed as Mean ± SE of three independent experiments with the similar results. *p<0.05; and **p<0.01 compared with the control

treated with different concentrations of vitexicarpin for 24 h. As shown in Figure 3, vitexicarpin dose-dependently increased the percentage of cells in G2/M phase to 26.97±0.9909%, 35.87±1.0117%, and 58.23±1.0826% versus 18.67±0.6417% in DMSO control group after treated with 10, 30, and 50 μM vitexicarpin, with a corresponding decrease in G0/G1 phase.

Vitexicarpin Induces Increased Generation of ROS in Human Prostate Carcinoma PC-3 cells

Cell apoptosis can be initiated via two distinct but convergent pathways: the death receptors and mitochondrial pathways (Schultz and Harrington, 2003). Mitochondria-mediated intracellular signals pathway is characterized by increased ROS generation, mitochondrial membrane potential (MMP) disruption, and glutathione (GSH) depletion (Schultz and Harrington, 2003; Antico et al., 2012). Recently, several studies showed that vitexicarpin induced apoptosis in cervical cancer cells through ROS-mediated mitochondrial signaling pathways (Gottlieb, 2001; Antico et al., 2012). It is reasonable to ask whether or not vitexicarpin could increase ROS generation in PC-3 cells. So, we measured ROS generation in PC-3 cells by using a Reactive Oxygen Species Assay Kit.




Figure 3. Effect of Vitexicarpin on PC-3 Cell Cycle Progression. (A) Flow cytometry analysis of cell cycle phase distribution in PC-3 cells. The cells were treated with 0, 10, 30, and 50 μM vitexicarpin for 24 h and then stained with PI for flow cytometric analysis. X-axis and y-axis represent DNA content and number of cells, respectively. (B). Data are expressed as Mean ± SE of three independent experiments with the similar results. *p<0.05; and **p<0.01 compared with the control




Figure 4. Effect of Vitexicarpin on ROS Generation in PC-3 Cells. (A) Flow cytometry analysis of intracellular ROS level in PC-3 cells. After treated with 0, 10, 30, and 50 μM vitexicarpin for 24 h, the PC-3 cells were stained with 10 μmol/L DCFH-DA for flow cytometric analysis. (B) Summarized results from (A), DCF fluorescence data are expressed as Mean ± SE of three independent experiments with the similar results. **p<0.01 compared with the control




Figure 5. Effect of Vitexicarpin on MMP in PC-3 Cells. (A) Cells were treated with indicated concentrations of vitexicarpin for 24 h, and then stained with 10 μg/ml Rho-123 at 37 °C for 30 min in darkroom. Rho-123 fluorescence was assayed using flow cytometry. (B) Data are expressed as Mean ± SE of three independent experiments with the similar results. **p<0.01 compared with the control

As shown in Figure 4, ROS levels in PC-3 cells were significantly increased from 5.49±0.6296% (DMSO control group) to 12.89±1.4849%, 17.19±1.0523%, and 24.06±1.616% respectively after treated with 10, 30, or 50 μM vitexicarpin (p<0.01).

Vitexicarpin Disrupts Mitochondrial Membrane Potential (MMP) in Human Prostate Carcinoma PC-3 cells

Because excess ROS production can induce apoptosis by depolarization of MMP, we determined MMP in PC-3 cells by using Rho-123 staining and flow cytometry assay. PC-3 cells were incubated with different concentrations




Figure 6. Effect of Vitexicarpin on Major Cell Cycle Regulators and Mitochondrial Apoptosis Regulators.

PC-3 cells were treated with indicated concentrations vitexicarpin for 24 h. Cell lysates were prepared and subjected to western blotting for CDK1, Cyclin B1 (A) Caspase-3, Bcl-2, Bax and Cytochrome c (B). Gray values were determined using Image J software. Data are expressed as Mean \pm SEM (n=3). *p<0.05; and **p<0.01 compared with the control (C)

of vitexicarpin (10, 30, 50 μ M) for 24 h before Rho-23 staining. The data indicated that vitexicarpin significantly decreased MMP (87.09 \pm 1.5568%, 67.23 \pm 2.2089% & 59.77 \pm 2.2775% respectively vs 96.16 \pm 1.2768% in control group) (Figure 5).

Effect of Vitexicarpin on Major Cell Cycle Regulators and Mitochondrial Apoptosis Regulatory Proteins

To get better insight into vitexicarpin-induced apoptosis in PC-3 cells, we detected the expression of some major cell cycle regulators (CDK1 and Cyclin B1) and mitochondrial apoptosis regulators (Caspase-3, Bax, Cytochrome c, and Bcl-2) using Western blotting analysis. The results showed that vitexicarpin significantly increased the expression of Caspase-3, Bax, and release of Cytochrome c; meanwhile decreased the expression of Bcl-2, CDK1 and Cyclin B1 (Figure 6).

Discussion

Chemotherapy is currently one of the most important available therapies in cancer treatment. Herbal medicines have been proven to be a major source of novel agents with diverse pharmaceutical activities and structures (Mishra and Tiwari, 2011; De et al., 2012; Mondal et al., 2012). Natural products have traditionally provided a rich source of drugs for many diseases, including cancer. Previous studies have indicated that vitexicarpin has inhibitory activities against various cancer cell lines (Shen et al., 2009; Chen et al., 2011; Koh et al., 2011; Yang et al., 2011; He et al., 2012; Zeng et al., 2012), while its effects on human prostate cancer cells remain uninvestigated. In the present paper, we showed that vitexicarpin is able to inhibit growth of human PC-3 prostate cancers. It is fully accepted that cell growth inhibition might be resulted from cell cycle block and cell death. Apoptosis (programmed cell death) and necrosis are the two major types of cell death. To further investigate mechanisms involve in this effect, we performed flow cytometric analysis of apoptosis and cell cycle arrest. We found that vitexicarpin induced cell cycle arrest at G2/M phase. This result is consistent with previous studies that vitexicarpin induced G2/M phase arrest in several types of human cancer cell lines (Shen et al., 2009; He et al., 2012).

Two classes of regulatory molecules, the cyclins and the cyclin-dependent kinases (CDKs), play key roles in cell cycle control. Different cyclin and CDK combinations determine their downstream targeted proteins. Cyclin B1/CDK1 complex is pivotal in regulating cells from G2 to M phase transition (Giono and Manfredi, 2006; Enomoto et al., 2009; Gavet and Pines, 2010). To elucidate the molecular mechanism underlying this G2/M phase blockage, we measured the expressions of Cyclin B1 and CDK1 using Western blotting analysis. The results demonstrated that the expression of these proteins decreased significantly indicating that vitexicarpin arrested the cell cycle at G2 phase in PC-3 cells. Similar results were also found in vitexicarpin treated human hepatocellular carcinoma (HCC) cells (He et al., 2012).

Apoptosis is a main reason for cell growth inhibition, and many cytotoxic agents arrest the cell cycle at specific phase and then induce apoptosis (Muschel and McKenna, 1996; Komata et al., 2003; Maddika et al., 2007; Batsi et al., 2009). So, we performed flow cytometric analysis by using propidium iodide (PI) / Annexin V-FITC double staining methods. It showed that vitexicarpin dose-dependently induced apoptosis in PC-3 cells. Previous studies have indicated that vitexicarpin induced apoptosis in hepatocellular carcinoma HepG2 (Yang et al., 2011; He et al., 2012) and human cervical cancer cells (Chen et al., 2011; Zeng et al., 2012). Also, we found vitexicarpin induced apoptotic cell death in PC-3 cells in a dose-dependent manner.

Over production of ROS induces oxidative damage including; lipid peroxidation, protein oxidation and DNA damage which results in modulation of Bcl-2 family proteins and dissipation of mitochondrial membrane potential, which are characteristic features of mitochondrial apoptosis (Khan et al., 2012). We here examined whether vitexicarpin increases intracellular ROS generation in PC-3 cells. Significant increased ROS generations were found in the vitexicarpin-treated PC-3 cells. Because excessive ROS production can induce depolarization of MMP, next we performed flow cytometric analysis to determine MMP in control and vitexicarpin-treated cells. The data showed that MMP in vitexicarpin-treated PC-3 cells was significantly decreased which is in line well with previous result found in Hela cells (Chen et al., 2011; Zeng et al., 2012).

Apoptosis can be initiated through two distinct but convergent pathways: the death receptors and mitochondrial pathways (Schultz and Harrington, 2003). The process of apoptosis is triggered by a diverse range of cellular signals. A balance between pro-apoptotic (BAX, BID, BIM, PUMA, BAK, or BAD) and anti-apoptotic (Bcl-XL, Bcl-2, BCLW, MCL1, A1, or BOO/DIVA) proteins of the Bcl-2 family controls the mitochondrial apoptosis pathway (Zinkel et al., 2006). To investigate whether vitexicarpin can trigger intrinsic apoptosis in PC-3 cells, we examined effect of the compound on expression of Bcl-2 and Bax proteins using Western blot analysis. The results demonstrated that expression of Bcl-2 gradually decreased while expression of Bax increased in cells treated with vitexicarpin, suggesting that the compound induces apoptosis through intrinsic pathway.

Release of Cytochrome c into cytosol in vitexicarpin-treated cells further support the above findings (Gottlieb, 2001; Antico et al., 2012).

In conclusion, our data demonstrated for the first time that vitaxicarpin induced apoptosis in PC-3 cells which is associated with increased ROS generation, modulation of Bcl-2 family proteins, and disruption of MMP and activation of Caspase-3. Therefore, vitaxicarpin may become a leading drug in the development of chemotherapeutic or chemopreventive of human prostate cancer.

Acknowledgements

The authors are grateful to Dr. Muhammad Khan (Department of zoology, University of the Punjab, P.O. Box No. 54590. Lahore, Pakistan) for critical reading of the manuscript and helpful suggestions. This work was supported by Specialized Research Fund for the Doctoral Program of Higher Education (20112136110002), National Natural Science Fund (No.30973577).

References

- Antico Arciuch VG, Elguero ME, Poderoso JJ, et al (2012). Mitochondrial regulation of cell cycle and proliferation. *Antioxid Redox Sign*, **16**, 1150-80.
- Batsi C, Markopoulou S, Kontargiris E, et al (2009). Bcl-2 blocks 2-methoxyestradiol induced leukemia cell apoptosis by a p27(Kip1)-dependent G1/S cell cycle arrest in conjunction with NF-kappaB activation. *Biochem Pharmacol*, **78**, 33-44.
- Chen D, Cao J, Tian L, et al (2011). Induction of apoptosis by casticin in cervical cancer cells through reactive oxygen species-mediated mitochondrial signaling pathways. *Oncol Rep*, **26**, 1287-94.
- Choudhary MI, Azizuddin, Jalil S, et al (2009). Antiinflammatory and lipoxygenase inhibitory compounds from *Vitex agnus-castus*. *Phytother Res*, **23**, 1336-9.
- De Luca V, Salim V, Atsumi SM, et al (2012). Mining the biodiversity of plants: a revolution in the making. *Science*, **336**, 1658-61.
- Enomoto M, Goto H, Tomono Y, et al (2009). Novel positive feedback loop between Cdk1 and Chk1 in the nucleus during G2/M transition. *J Biol Chem*, **284**, 34223-30.
- Gavet O, Pines J (2010). Progressive activation of CyclinB1-Cdk1 coordinates entry to mitosis. *Dev Cell*, **18**, 533-43.
- Giono LE, Manfredi JJ (2006). The p53 tumor suppressor participates in multiple cell cycle checkpoints. *J Cell Physiol*, **209**, 13-20.
- Gottlieb RA (2001). Mitochondria and apoptosis. *Biol Signal Recept*, **10**, 147-61.
- He L, Yang X, Cao X, et al (2012). Casticin induces growth suppression and cell cycle arrest through activation of FOXO3a in hepatocellular carcinoma. *Oncol Rep*, **29**, 103-8.
- Ji YB, Qu ZY, and Zou X (2011). Juglone-induced apoptosis in human gastric cancer SGC-7901 cells via the mitochondrial pathway. *Exp Toxicol Pathol*, **63**, 69-78.
- Khan M, Ding C, Rasul A, et al (2012). Isoalantolactone induces reactive oxygen species mediated apoptosis in pancreatic carcinoma PANC-1 cells. *Int J Biol Sci*, **8**, 533-47.
- Khan M, Zheng B, Yi F, et al (2012). Pseudolaric Acid B induces caspase-dependent and caspase-independent apoptosis in u87 glioblastoma cells. *Evid-Based Complalt*, Article ID 957568.
- Koh DJ, Ahn HS, Chung HS, et al (2011). Inhibitory effects of casticin on migration of eosinophil and expression of chemokines and adhesion molecules in A549 lung epithelial cells via NF- κ B inactivation. *J Ethnopharmacol*, **136**, 399-405.
- Komata T, Kanzawa T, Takeuchi H, et al (2003). Antitumour effect of cyclin-dependent kinase inhibitors (p16(INK4A), p18(INK4C), p19(INK4D), p21(WAF1/CIP1) and p27(KIP1)) on malignant glioma cells. *Brit J Cancer*, **88**, 1277-80.
- Maddika S, Ande SR, Panigrahi S, et al (2007). Cell survival, cell death and cell cycle pathways are interconnected: implications for cancer therapy. *Drug Resist Updat*, **10**, 13-29.
- Mesaik MA, Azizuddin, Murad S, et al (2009). Choudhary MI. Isolation and immunomodulatory properties of a flavonoid, casticin from *Vitex agnus-castus*. *Phytother Res*, **23**, 1516-20.
- Mishra BB, Tiwari VK (2011). Natural products: an evolving role in future drug discovery. *Eur J Med Chem*, **46**, 4769-807.
- Mondal S, Bandyopadhyay S, Ghosh MK, et al (2012). Natural products: promising resources for cancer drug discovery. *Anti-cancer Agent Me*, **12**, 49-75.
- Muschel RJ, McKenna WG (1996). Alterations in cell cycle control during tumor progression: effects on apoptosis and the response to therapeutic agents. *Curr Top Microbiol*, **213**, 197-213.
- Prostate Cancer Overview. *American Cancer Society*, 02/27/2012.
- Schafer KA (1998). The cell cycle: a review. *Vet Pathol*, **35**, 461-78.
- Schultz DR, Harrington WJ Jr (2003). Apoptosis: programmed cell death at a molecular level. *Semin Arthritis Rheu*, **32**, 345-69.
- Shen JK, Du HP, Yang M, et al (2009). Casticin induces leukemic cell death through apoptosis and mitotic catastrophe. *Ann Hematol*, **88**, 743-52.
- Yang J, Yang Y, Tian L, et al (2011). Casticin-induced apoptosis involves death receptor 5 upregulation in hepatocellular carcinoma cells. *World J Gastroentero*, **17**, 4298-307.
- Zeng F, Tian L, Liu F, et al (2012). Induction of apoptosis by casticin in cervical cancer cells: reactive oxygen species-dependent sustained activation of Jun N-terminal kinase. *Acta Bioch Bioph Sin*, **44**, 442-9.
- Zinkel S, Gross A, Yang E (2006). BCL2 family in DNA damage and cell cycle control. *Cell Death Differ*, **13**, 1351-9.