

Meal skipping relates to food choice, understanding of nutrition labeling, and prevalence of obesity in Korean fifth grade children

Hye-Young Kim¹, Na-Rae Lee¹, Jung-Sug Lee², Young-Sun Choi³, Tong-Kyung Kwak⁴, Hae Rang Chung⁵, Sehyug Kwon⁶, Youn-Ju Choi⁷, Soon-Kyu Lee⁷ and Myung-Hee Kang^{8§}

¹Department of Food & Nutrition, Yongin University, Yongin 449-714, Korea

²FANSA (Food and Nutrition Statistical Analysis), Seoul 153-764, Korea

³Department of Food & Nutrition, Daegu University, Daegu 712-714, Korea

⁴Department of Food & Nutrition, Yonsei University, Seoul 120-749, Korea

⁵Nutrition for the Future Inc., Seoul 151-848, Korea

⁶Department of Statistics, Hannam University, Daejeon 306-791, Korea

⁷Dietary Life Safety Division, KFDA, Cheongwon 363-951, Korea

⁸Department of Food & Nutrition, Daeduk Valley Campus, Hannam University, Daejeon 305-811, Korea

Abstract

This study was performed to investigate the differences in food choice, nutrition labeling perceptions, and prevalence of obesity due to meal skipping in Korean elementary school children. A national survey was performed in 2010 to collect data on food intake frequency, understanding of nutrition labeling, and body mass index from 2,335 fifth grade students in 118 elementary schools selected from 16 metropolitan local governments by stratified cluster sampling. The data were analyzed using the SAS 9.1 and SUDAAN 10.0 packages. Students who consumed three meals for 6-7 days during the past week were classified into the regular meal eating (RM) group (n=1,476) and those who did not were placed into the meal skipping (MS) group (n=859). The daily intake frequency of fruits, vegetables, kimchi, and milk was significantly lower in the MS group compared to that in the RM group ($P<0.001$), whereas the daily intake frequency of soft drinks and instant noodles (ramyeon) was significantly higher in the MS group than that in the RM group ($P<0.05$). The MS group demonstrated a significantly lower degree of understanding with regard to nutrition labeling and high calorie foods containing low nutritional value than that in the RM group. The distribution of obesity based on the percentile criteria using the Korean growth chart was different between the MS and RM groups. The MS group (8.97%) had a higher percentage of obese subjects than that in the RM group (5.38%). In conclusion, meal skipping was related to poor food choice, low perception of nutrition labeling, and a high prevalence of obesity in Korean fifth grade children.

Key Words: Meal skipping, children, food choice, nutrition labeling, obesity

Introduction

Regular meal eating of three meals a day with balanced nutrition is essential for optimal growth and development in children [1]. The family dinner is related to increased intake of healthy foods such as fruits, vegetables, and dairy foods [2]. High-calorie sweet beverages or fast foods are consumed more during meal skipping, because calories that should be ingested in one meal increase [3]. When poor eating behavior continues, excessive energy intake can be accompanied by decreased intake of dietary fiber and micronutrients, resulting in possible threatening of children's health [4].

Consuming regular meals also has a beneficial influence on

study performance and grade of the student [5,6]. In particular, intake of breakfast greatly helps improve memory in children with a risk for nutrient deficiency [7], and the school breakfast program contributes to improve class attendance and academic performance [5,8].

The incidence of obesity in children and adolescents has increased around the world [9]. Obesity in childhood can increase the chance of acquiring metabolic syndrome [10] and the probability of obesity in adulthood [11]. According to the most recent 1998-2007 Korean National Health and Nutrition Examination Survey, the proportion of obese teens doubled from 5.4% to 10.3% [12]. The average incidence of metabolic syndrome in Korean teens is only 4.2%, but that of obese teens

This research was supported by a grant (08082KFDA999) from the Korea Food & Drug Administration in 2010.

Corresponding Author: Myung-Hee Kang, Tel. 82-42-629-8791, Fax. 82-42-629-8808, Email. mhkang@hnu.kr

Received: March 2, 2012, Revised: July 19, 2012, Accepted: July 19, 2012

©2012 The Korean Nutrition Society and the Korean Society of Community Nutrition

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

is significantly higher at 31.3% [13].

Meal skipping is related to obesity in many studies conducted abroad [14-16]. The body mass index (BMI) and body weight of children skipping breakfast is higher than in those who eat breakfast [1,17,18]. Additionally, obese children show higher rates of meal skipping compared to non-obese children [19,20]. The reason for obesity due to meal skipping might be related to the observation that children do not select healthy foods or have poor eating habits [3,21].

Nutrition labeling provides proper nutrition information to consumers and helps with reasonable food selection and health management by labeling the type and amount of nutrients in processed foods [22]. "High-calorie low-nutritional value food" is a type of nutrition labeling in Korea that is marked on processed foods due to concerns of developing obesity or nutritional imbalance [23]. A correct understanding of nutrition labeling may help students to prevent obesity through wise selection of healthy foods.

The relationship between meal skipping and risk of obesity has been studied intensively abroad, but that relationship has not been investigated nationwide in Korean children. Therefore, the purpose of this study was to investigate the relationship between meal skipping and prevalence of obesity in Korean children. We hypothesized that meal-skipping students select healthy foods less frequently and have a lower degree of understanding of nutrition labeling compared to those of regular meal eating students. To test this hypothesis, we divided subjects into a meal skipping group (MS) and a regular meal eating group (RM) and investigated the differences in children's food intake frequency from several food groups, the degree of nutrition labeling understanding, and obesity prevalence.

Subjects and Methods

Subjects and survey methods

This study used data from the 2010 national survey on the "Children's perception and practice level of dietary life" [24]. The survey was conducted in 2,335 (1,141 boys, 48.2% and 1,194 girls, 51.8%) fifth grade students in 118 elementary schools selected from 16 metropolitan local governments by stratified cluster sampling. The survey was conducted between June and July 2010 by personal interview through visits to each school by trained interviewers.

The survey questionnaire included meal intake frequency for breakfast, lunch, and dinner for 1 week and the food intake frequency of fruits, vegetables, kimchi, white milk, flavored milk, pizza, hamburgers, fried chicken, soft drinks, instant noodles, crackers, and chocolate. A visual measure was presented for foods to reduce differences in serving size quantities among individuals [14].

Multiple choice questions on the understanding of nutrition

labeling and high calorie foods with low nutritional value were also included in the survey questionnaire. To investigate the degree of understanding of nutrition labeling, the question "What is the nutrient that obese people should carefully check on the nutrition label of a food package?" was included. In addition, to identify the degree of recognition for 'high calorie foods with low nutritional value', the question "A high calorie food with low nutritional value is a food that has higher calories and lower nutritional value than general foods and that is related to developing obesity or nutritional imbalance. What is the lowest nutrient in a high calorie food with low nutritional value?" was included.

For meal intake frequency, the frequency of 6-7 times per week was converted to 6.5 times/week, 3-5 times per week was converted to 4 times/week, 1-2 times/week to 1.5 times/week, and no intake to 0 time/week. For the evaluation of food intake frequency for fruits, vegetables, milk, fast-food, and soft drinks during the past week, the frequency of more than twice a day was converted to 2 times/day, once per day to 1 time/day, once every other day to 0.5 times/day, 1-2 times a week to 0.21 times/day, and no intake to 0 times/day.

BMI was calculated from self-reported weight and height obtained from the questionnaires. The percentile for BMI-for-age was calculated. Underweight was defined as a BMI < 15th percentile, normal weight was \geq 15th percentile and < 85th percentile, overweight was \geq 85th percentile and < 95th percentile and obesity was \geq 95th percentile using the age- and sex-specific BMI cutoff points proposed by the Korean Centers for Disease Control and Prevention [25].

The subjects were divided into two groups based on intake of a three regular meal per day pattern. The RM group included those who answered "6-7 times a week" to all three questions that asked how often they ate breakfast, lunch, and dinner. The others were placed into the MS group. The number of subjects in the RM group was 1,476 (64.2%) and that in the MS group was 859 (35.8%).

Statistical analysis

All analyses were conducted using a survey weighting to account for the survey design, which consisted of stratified cluster samples. Probability sampling weights were used in conjunction with strata and primary sampling units to apply weight to the data analysis. The collected data were used to calculate the percentage or mean and standard errors using the SAS 9.12 (SAS Institute, Cary, NC, USA) and SUDAAN 10.0 programs. The difference between groups for food intake frequency was analyzed using Student's *t*-test, and the frequency difference between the groups for nutrition labeling and high calorie foods with low nutritional value was analyzed using the χ^2 -test. Statistical significance was verified at the $\alpha = 0.05$ level.

Results

Meal regularity of the subjects

Meal intake frequency of the subjects is shown in Table 1. The average meal intake frequency in all subjects was 5.55 times/week for breakfast, 6.15 times/week for lunch, and 6.14 times/week for dinner. While the RM group had all meals 6-7 times/week, the MS group showed 3.85 times/week for breakfast, 5.53 times/week for lunch, and 5.5 times/week for dinner.

Daily food intake frequency by meal regularity

The daily intake frequency of healthy foods such as fruits, vegetables, and milk is shown in Table 2. The daily food intake

Table 1. Meal intake frequency (times/week) of the students

	Total	MS ¹⁾	RM ²⁾
	2,335 (100.0) ³⁾	859 (35.8)	1,476 (64.2)
Breakfast	5.55 ± 0.05 ²⁾	3.85 ± 0.09	6.50 ± 0.00***
Lunch	6.15 ± 0.03	5.53 ± 0.06	6.50 ± 0.00***
Dinner	6.14 ± 0.03	5.50 ± 0.07	6.50 ± 0.00***

¹⁾ MS: meal skipper; having breakfast, lunch, or dinner ≤ 3-5 days/week

²⁾ RM: regular meal consumer; having breakfast, lunch, and dinner 6-7 days/week

³⁾ N (%)

⁴⁾ Mean ± standard error

*** $P < 0.001$, significantly different between MS and RM groups by t -test.

Table 2. Daily intake frequency of fruits, vegetables, and milk (times/day)

	Total	MS ¹⁾	RM ²⁾
	2,335 (100.0) ³⁾	859 (35.8)	1,476 (64.2)
Fruit	0.71 ± 0.02 ⁴⁾	0.62 ± 0.03	0.76 ± 0.02***
Kimchi	1.32 ± 0.02	1.21 ± 0.03	1.38 ± 0.02***
Other vegetables	1.03 ± 0.02	0.92 ± 0.03	1.09 ± 0.02***
White Milk	0.92 ± 0.01	0.84 ± 0.03	0.97 ± 0.02***
Flavored milk	0.21 ± 0.01	0.23 ± 0.02	0.19 ± 0.01

¹⁾ MS: meal skipper; having breakfast, lunch, or dinner ≤ 3-5 days/week

²⁾ RM: regular meal consumer; having breakfast, lunch, and dinner 6-7 days/week

³⁾ N (%)

⁴⁾ Mean ± standard error

*** $P < 0.001$, significantly different between the MS and RM groups by t -test.

Table 3. Daily intake frequency of fast food, soft drinks, and high-calorie snacks (times/day)

	Total	MS ¹⁾	RM ²⁾
	2,335 (100.0) ³⁾	859 (35.8)	1,476 (64.2)
Pizza	0.08 ± 0.01 ⁴⁾	0.09 ± 0.01	0.08 ± 0.01
Hamburger	0.07 ± 0.01	0.07 ± 0.01	0.06 ± 0.01
Fried chicken	0.12 ± 0.01	0.13 ± 0.01	0.12 ± 0.01
Soft drink	0.22 ± 0.01	0.25 ± 0.01	0.20 ± 0.01***
Instant noodle (Ramyon)	0.20 ± 0.01	0.23 ± 0.01	0.18 ± 0.01**
High-calorie snack (cracker and chocolates)	0.25 ± 0.01	0.26 ± 0.01	0.25 ± 0.01

¹⁾ MS: meal skipper; having breakfast, lunch, or dinner ≤ 3-5 days/week

²⁾ RM: regular meal consumer; having breakfast, lunch, and dinner 6-7 days/week

³⁾ N (%)

⁴⁾ Mean ± standard error

*** $P < 0.001$, significantly different between MS and RM groups by t -test.

frequency was 0.71 times for fruit, 1.32 times for kimchi, and 1.03 times for vegetables, except kimchi. Thus, the intake frequency of all vegetables including kimchi was 2.35 times/day. Daily intake of milk was 0.92 cup (200 ml basis) for white milk and 0.21 cup for flavored milk, adding up to 1.13 cups/day.

The daily intake frequency of fruit was 0.62 times in the MS group, which was significantly lower compared to 0.76 times in the RM group ($P < 0.001$), and the daily intake frequency of kimchi and other vegetables was 1.21 and 0.92 times, respectively, in the MS group, which was lower compared to the 1.38 and 1.09 times, respectively, in the RM group ($P < 0.001$). The average daily intake amounts of white milk were 0.84 cup in the MS group, which was significantly lower compared to the 0.97 cups in the RM group ($P < 0.001$), but that of flavored milk was 0.23 cups in the MS group, which tended to be higher than the 0.19 cups in the RM group, but the difference was not significant.

The daily intake frequency of fast-food, soft drinks, and high-calorie snacks is shown in Table 3. The average daily intake of pizza was 0.08 times, hamburgers 0.07 times, fried chicken 0.12 times, soft drinks 0.22 times, instant noodles 0.20 times, and crackers and chocolate 0.25 times, showing frequent intake of fast-food and snacks in the order of cracker and chocolate > soft drinks > instant noodles > fried chicken > pizza > hamburgers. Eating frequency was slightly higher in the MS group for pizza, hamburgers, fried chicken, crackers, and chocolate, compared to that in the RM group but the difference was not significant. In contrast, the frequency of soft drink and instant noodle (ramyeon) intake was 0.25 and 0.23, respectively, in the MS group, which was significantly higher compared to the 0.2 and 0.18, respectively, in the RM group ($P < 0.01$).

Understanding of nutrition labeling and high calorie foods with low nutritional value

The understanding of nutrition labeling and high calorie foods with low nutritional value is shown in Table 4. Subjects were asked for the nutrient that obese individuals should check carefully to confirm whether they properly understand the

Table 4. Understanding of nutrition labeling and high calorie foods with low nutrition value

	Total	MS ¹⁾	RM ²⁾	χ^2 -value
	2,335 (100.0) ³⁾	859 (35.8)	1,476 (64.2)	
Understanding for nutrition labeling				
Correct	906 (38.7)	293 (35.3)	613 (40.7)	4.46*
Wrong	1,429 (61.3)	566 (64.7)	863 (59.3)	
Understanding for high calorie foods with low nutrition value				
Correct	923 (40.2)	316 (35.6)	607 (42.7)	7.37**
Wrong	1,412 (59.9)	543 (64.4)	869 (57.3)	

¹⁾ RM: Regular meal eating group

²⁾ MS: meal skipping group

³⁾ N (%)

* $P < 0.05$, ** $P < 0.01$, significantly different between the RM and MS groups by χ^2 -test.

Table 5. Anthropometric measurements of the students

	Total	MS ¹⁾	RM ²⁾
	2,335 (100.0) ³⁾	859 (35.8)	1,476 (64.2)
Height (cm)	149.66 ± 1.57	150.63 ± 2.96	149.13 ± 1.83
Weight (kg)	39.52 ± 0.25	40.62 ± 0.47	38.91 ± 0.29**
BMI (kg/m ²)	18.48 ± 0.09	18.78 ± 0.17	18.31 ± 0.11*

¹⁾ MS: meal skipper; having breakfast, lunch, or dinner ≤ 3-5 days/week

²⁾ RM: regular meal consumer; having breakfast, lunch, and dinner 6-7 days/week

³⁾ N (%)

⁴⁾ Mean ± standard error

* $P < 0.05$, ** $P < 0.01$, significantly different between MS and RM groups by t -test.

Table 6. Distribution of obesity in the students

	Total	MS ¹⁾	RM ²⁾	χ^2 -value
	2,335 (100.0) ³⁾	859 (35.8)	1,476 (64.2)	
Underweight	105 (5.36)	35 (5.27)	70 (5.41)	2.71*
Normal	1,781 (77.24)	632 (74.29)	1,149 (78.88)	
Overweight	279 (10.73)	116 (11.46)	163 (10.33)	
Obesity	170 (6.67)	76 (8.97)	94 (5.38)	

¹⁾ MS: meal skipper; having breakfast, lunch, or dinner ≤ 3-5 days/week

²⁾ RM: regular meal consumer; having breakfast, lunch, and dinner 6-7 days/week

³⁾ N (%)

⁴⁾ Underweight was defined as a body mass index (BMI) < 15th percentile, normal ≥ 15th percentile, and < 85th percentile, overweight ≥ 85th percentile and < 95th percentile, and obesity ≥ 95th percentile using the age- and sex-specific BMI cutoff points proposed by the Korean Centers for Disease Control and Prevention.

* $P < 0.05$, significantly different between the MS and RM groups by χ^2 -test.

nutrition labeling and only 39% of the subjects correctly answered. The proportion of correct answers for understanding nutritional labeling was significantly lower in the MS group (35.3%) compared to that in the RM group (40.7%) ($P < 0.05$).

The proportion of correct answers for the question of high calorie foods with low nutritional value was 40.2% and, overall, the correct understanding of high calorie foods with low nutritional value was insufficient in the study subjects. The MS group (35.6%) had a significantly lower degree of understanding for high calorie foods with low nutritional value compared to that in the RM group (42.7%) ($P < 0.01$).

Anthropometric measurements

Anthropometric measurements of the students are shown in Table 5. The average height and weight of the students were 149.7 cm and 39.5 kg, and the average BMI was 18.48 kg/m². No difference in height was observed between the two groups, but weight was significantly ($P < 0.01$) greater in the MS group. BMI in the MS group was also significantly higher than that in the RM group.

The distribution of student obesity is shown in Table 6. A total of 5.36% of the study subjects were underweight, 77.24% were normal (non-obese), 10.73% were overweight, and 6.67% were obese. The distribution of obesity was significantly ($P < 0.05$) different between the two groups and the MS group (8.97%) had a higher percentage of obese subjects than that in the RM group (5.38%).

Discussion

This study was performed to investigate differences in food choice, understanding of nutrition labeling, and prevalence of obesity due to meal skipping in Korean children. The criteria for meal skipping are different among investigators [14] and, in this study, subjects who had at least one meal less than five times/week were assigned to the MS group. The MS group showed an average eating frequency of 3.85 times/week for breakfast and 5.53 and 5.50 times/week for lunch and dinner, respectively. The reason for meal skipping was not asked in this study, but in other studies, the main reason for skipping breakfast was due to “no appetite” or “no time to eat”. “Too much snack consumption” was provided as the reason for skipping lunch and “weight control” was added as the reason for skipping dinner [26-28].

The frequency of three meals/day in Korean elementary students has varied depending on survey year. In a 1998 survey ($n = 2,433$), with the influence of International Monetary Fund in Korea, the proportion of children eating breakfast everyday was 64%, eating lunch was 82.5%, and eating dinner was 54.3% [29]. In a 2002 survey ($n = 544$), the proportion of students eating breakfast everyday was 68.9%, lunch 91.7%, and dinner 81.1%, showing an overall increase of meal intake frequency [30]. In the present 2010 nationwide survey, the proportion of students eating breakfast everyday was 73.5%, eating lunch was 88.0%, and eating dinner was 87.7%. Therefore, the proportion of students eating breakfast and dinner every day has increased, but the students eating lunch every day decreased slightly compared to that in 2002. It is difficult to identify the reasons for the decreased percentage of students eating lunch every day. It was thought that it was not related to weekdays when lunch was eaten at school but was related to weekends when lunch might be skipped due to late breakfast with the family.

The dietary reference intakes for Koreans (KDRI) recommend that elementary school children should eat one serving of fruit, five servings of vegetables, and two servings of milk/day [31]. However, in this study, students reported 0.71 servings of fruit, 2.35 servings of vegetables, and 1.13 cups of milk/day, which is insufficient compared to the amounts recommended for elementary school students by the KDRI. A detailed dietary intake survey was not conducted in this study, but an effort is needed to provide more fruits, vegetables, and dairy products to children during meals in the future.

The MS group had a lower intake frequency of fruits, vegetables, and milk compared to those in the RM group. Other studies have also reported that children skipping meals show insufficient daily intake for fruits and vegetables compared to children eating meals [32], significantly lower mean adequacy ratio for micro-nutrients [33], or decreased dietary variety scores [34].

As meal skipping increases, the quality of meals become poorer, hunger is aggravated, and excessive eating occurs during

the next meal [35,36]. Breakfast skipping is related to increased consumption of soft drinks and French fries [37] as well as increased snack intake [38]. In adolescents, breakfast skipping is related to skipping other meals, which directly worsens improper food intake [39]. Additionally, the possibility of eating a ready-made dinner with lower nutritional quality increases when children do not eat dinner with the family [40]. In this study, the intake frequency of instant noodles (ramyeon) and soft drinks was higher in the MS group than that in the RM group, suggesting the possibility of contributing to high-energy and high-sugar intake.

The degree of accurate nutrition labeling understanding and high calorie foods with low nutritional value was only 38.7% and 40.2%, respectively, of all students surveyed. Thus, more efforts are needed so that children understand nutrition labeling and select more proper foods to manage their health. The degree of an accurate understanding of nutrition labeling was lower in the MS group than that in the RM group. A better understanding of nutrition labeling may help students who skip meals to prevent obesity through wise selection of healthy foods.

Meal skipping can influence the incidence of childhood obesity [14-18]. Several studies on meal skipping have reported the effect of skipping breakfast on health [3,4]. In those studies, children who skipped breakfast had higher BMIs than those who ate breakfast [1,17,18,41]. The 1996-2006 National Health and Nutrition Examination Survey in the USA showed that 9-13 year old children who skipped breakfast had higher BMIs, greater waist circumference, and high prevalence of obesity for their age compared to those who ate breakfast [1,42]. Breakfast skipping from childhood to adulthood increases not only the obesity rate but also fasting blood insulin and cholesterol levels, suggesting that meal skipping can cause obesity and related abnormal blood metabolism [4,43]. In our study, the average BMI in the MS and RM groups was 18.31 and 18.78, respectively, which was within the normal range for children. However, the proportion with a BMI > 95th percentile was 8.97% in the MS group, which was significantly higher than the 5.38% in the RM group, suggesting that meal skipping affects obesity prevalence in Korean children.

This study had several limitations. First, because skipping breakfast, lunch, or dinner was categorized as the MS group and analyzed for differences in food intake frequency pattern with a three-meals-eating group, the characteristics of each meal skipped were not investigated. Second, the BMI was based on self-reported measures of weight and height. Third, the observed difference in food intake frequency and obesity due to meal skipping was based on data from a cross-sectional study. Therefore, causality between breakfast skipping and weight status cannot be established. A follow-up study is needed to assess long-term outcomes of meal regularity on obesity and health of Korean children.

In conclusion, meal skipping was related to obesity in Korean elementary students, which was related to the finding that

students with an irregular meal eating pattern selected unhealthy foods more often and did not understand nutrition labeling compared to those with a regular meal eating pattern. The development and distribution of customized nutrition education is needed to inform students of the importance of regular meal eating, to recommend adequate intake of fruits, vegetables, and dairy products, and to understand nutrition labeling correctly.

References

1. Deshmukh-Taskar PR, Nicklas TA, O'Neil CE, Keast DR, Radcliffe JD, Cho S. The relationship of breakfast skipping and type of breakfast consumption with nutrient intake and weight status in children and adolescents: the National Health and Nutrition Examination Survey 1999-2006. *J Am Diet Assoc* 2010;110:869-78.
2. Videon TM, Manning CK. Influences on adolescent eating patterns: the importance of family meals. *J Adolesc Health* 2003;32:365-73.
3. Dwyer JT, Evans M, Stone EJ, Feldman HA, Lytle L, Hoelscher D, Johnson C, Zive M, Yang M; Child and Adolescent Trial for Cardiovascular Health (CATCH) Cooperative Research Group. Adolescents' eating patterns influence their nutrient intakes. *J Am Diet Assoc* 2001;101:798-802.
4. Smith KJ, Gall SL, McNaughton SA, Blizzard L, Dwyer T, Venn AJ. Skipping breakfast: longitudinal associations with cardiometabolic risk factors in the Childhood Determinants of Adult Health Study. *Am J Clin Nutr* 2010;92:1316-25.
5. Kim HY, Frongillo EA, Han SS, Oh SY, Kim WK, Jang YA, Won HS, Lee HS, Kim SH. Academic performance of Korean children is associated with dietary behaviours and physical status. *Asia Pac J Clin Nutr* 2003;12:186-92.
6. Wesnes KA, Pincock C, Richardson D, Helm G, Hails S. Breakfast reduces declines in attention and memory over the morning in schoolchildren. *Appetite* 2003;41:329-31.
7. Cueto S, Jacoby E, Pollitt E. Breakfast prevents delays of attention and memory functions among nutritionally at-risk boys. *J Appl Dev Psychol* 1998;19:219-33.
8. Meyers AF, Sampson AE, Weitzman M, Rogers BL, Kayne H. School Breakfast Program and school performance. *Am J Dis Child* 1989;143:1234-9.
9. World Health Organization [Internet]. Obesity and overweight. Fact sheet N°311. 2012 [cited 2012 May 13]. Available from: <http://www.who.int/mediacentre/factsheets/fs311/en>.
10. Ribeiro J, Guerra S, Pinto A, Oliveira J, Duarte J, Mota J. Overweight and obesity in children and adolescents: relationship with blood pressure, and physical activity. *Ann Hum Biol* 2003; 30:203-13.
11. Guo SS, Wu W, Chumlea WC, Roche AF. Predicting overweight and obesity in adulthood from body mass index values in childhood and adolescence. *Am J Clin Nutr* 2002;76:653-8.
12. Ministry of Health and Welfare, Korea Centers for Disease Control and Prevention. 2007 Korea National Health and Nutrition Examination Survey. Seoul: Ministry of Health and Welfare; 2008. p.56.
13. Huh K. Metabolic syndrome in Korean children and adolescents; from the Korean NHANES 1998-2007 data analysis [master's thesis]. Busan: Department of Medicine, Inje University; 2010.

14. Rampersaud GC, Pereira MA, Girard BL, Adams J, Metz JD. Breakfast habits, nutritional status, body weight, and academic performance in children and adolescents. *J Am Diet Assoc* 2005;105:743-60.
15. Veltsista A, Laitinen J, Sovio U, Roma E, Järvelin MR, Bakoula C. Relationship between eating behavior, breakfast consumption, and obesity among Finnish and Greek adolescents. *J Nutr Educ Behav* 2010;42:417-21.
16. Horikawa C, Kodama S, Yachi Y, Heianza Y, Hirasawa R, Ibe Y, Saito K, Shimano H, Yamada N, Sone H. Skipping breakfast and prevalence of overweight and obesity in Asian and Pacific regions: a meta-analysis. *Prev Med* 2011;53:260-7.
17. Siega-Riz AM, Popkin BM, Carson T. Trends in breakfast consumption for children in the United States from 1965-1991. *Am J Clin Nutr* 1998;67:748S-756S.
18. Song WO, Chun OK, Obayashi S, Cho S, Chung CE. Is consumption of breakfast associated with body mass index in US adults? *J Am Diet Assoc* 2005;105:1373-82.
19. Ortega RM, Requejo AM, López-Sobaler AM, Quintas ME, Andrés P, Redondo MR, Navia B, López-Bonilla MD, Rivas T. Difference in the breakfast habits of overweight/obese and normal weight schoolchildren. *Int J Vitam Nutr Res* 1998;68:125-32.
20. Boutelle K, Neumark-Sztainer D, Story M, Resnick M. Weight control behaviors among obese, overweight, and nonoverweight adolescents. *J Pediatr Psychol* 2002;27:531-40.
21. Serra-Majem L, Ribas L, Pérez-Rodrigo C, García-Closas R, Peña-Quintana L, Aranceta J. Determinants of nutrient intake among children and adolescents: results from the enKid Study. *Ann Nutr Metab* 2002;46 Suppl 1:31-8.
22. World Health Organization. Nutrition Labels and Health Claims: the Global Regulatory Environment. Geneva: World Health Organization; 2004.
23. Chung HR, Kwak TK, Choi YS, Kim HY, Lee JS, Choi JH, Yi NY, Kwon S, Choi YJ, Lee SK, Kang MH. Development of evaluation indicators for a children's dietary life safety index in Korea. *Korean J Nutr* 2011;44:49-60.
24. Kang MH. In Depth Analysis of Child Food Safety Index and Evaluation of Cognition and Practice Level. KFDA Report. Seoul; 2010.
25. Korea Centers for Disease Control and Prevention. Growth Charts for Koreans. Cheongwon; 2007.
26. Kim KH. Food habits, eating behaviors and food frequency by gender and among Seoul and other regions in upper-grade elementary school children. *Korean J Community Nutr* 2010;15:180-90.
27. Kim MJ, Kim YH. Dietary habits, nutrition knowledge and dietary behaviors of the 3rd grade elementary school students in Ulsan area by sex and skipping breakfast. *J East Asian Soc Diet Life* 2010;20:209-17.
28. Yi BS, Yang IS. An exploratory study for identifying factors related to breakfast in elementary, middle and high school students. *Korean J Community Nutr* 2006;11:25-38.
29. Kim SH. Children's growth and school performance in relation to breakfast. *J Korean Diet Assoc* 1999;5:215-24.
30. Yi BS. A comparative study on dietary life and recognition of diet related factors in elementary, middle and high school students. *J Korean Diet Assoc* 2004;10:364-74.
31. The Korean Nutrition Society. Dietary Reference Intakes for Koreans. Seoul; 2010. p.503-50.
32. Utter J, Scragg R, Mhurchu CN, Schaaf D. At-home breakfast consumption among New Zealand children: associations with body mass index and related nutrition behaviors. *J Am Diet Assoc* 2007;107:570-6.
33. Yeoh YJ, Yoon J, Shim JE. Relation of breakfast intake to diet quality in Korean school-aged children: analysis of the data from the 2001 national health and nutrition survey. *Korean J Community Nutr* 2009;14:1-11.
34. Nicklas TA, Reger C, Myers L, O'Neil C. Breakfast consumption with and without vitamin-mineral supplement use favorably impacts daily nutrient intake of ninth-grade students. *J Adolesc Health* 2000;27:314-21.
35. Miech RA, Kumanyika SK, Stettler N, Link BG, Phelan JC, Chang VW. Trends in the association of poverty with overweight among US adolescents, 1971-2004. *JAMA* 2006;295:2385-93.
36. Lioret S, Volatier JL, Lafay L, Touvier M, Maire B. Is food portion size a risk factor of childhood overweight? *Eur J Clin Nutr* 2009;63:382-91.
37. Lattimore PJ, Halford JC. Adolescence and the diet-dieting disparity: healthy food choice or risky health behaviour? *Br J Health Psychol* 2003;8:451-63.
38. Schlundt DG, Hill JO, Sbrocco T, Pope-Cordle J, Sharp T. The role of breakfast in the treatment of obesity: a randomized clinical trial. *Am J Clin Nutr* 1992;55:645-51.
39. Sjöberg A, Hallberg L, Höglund D, Hulthén L. Meal pattern, food choice, nutrient intake and lifestyle factors in The Göteborg Adolescence Study. *Eur J Clin Nutr* 2003;57:1569-78.
40. Gillman MW, Rifas-Shiman SL, Frazier AL, Rockett HR, Camargo CA Jr, Field AE, Berkey CS, Colditz GA. Family dinner and diet quality among older children and adolescents. *Arch Fam Med* 2000;9:235-40.
41. Barton BA, Eldridge AL, Thompson D, Affenito SG, Striegel-Moore RH, Franko DL, Albertson AM, Crockett SJ. The relationship of breakfast and cereal consumption to nutrient intake and body mass index: the National Heart, Lung, and Blood Institute Growth and Health Study. *J Am Diet Assoc* 2005;105:1383-9.
42. Deshmukh-Taskar P, Nicklas TA, Morales M, Yang SJ, Zakeri I, Berenson GS. Tracking of overweight status from childhood to young adulthood: the Bogalusa Heart Study. *Eur J Clin Nutr* 2006;60:48-57.
43. Timlin MT, Pereira MA. Breakfast frequency and quality in the etiology of adult obesity and chronic diseases. *Nutr Rev* 2007;65:268-81.