

## RECURRENT JACOBI OPERATOR OF REAL HYPERSURFACES IN COMPLEX TWO-PLANE GRASSMANNIANS

IMSOON JEONG, JUAN DE DIOS PÉREZ, AND YOUNG JIN SUH

ABSTRACT. In this paper we give a non-existence theorem for Hopf hypersurfaces in the complex two-plane Grassmannian  $G_2(\mathbb{C}^{m+2})$  with recurrent normal Jacobi operator  $\bar{R}_N$ .

### 1. Introduction

Let  $(\bar{M}, \bar{g})$  be a Riemannian manifold. The Jacobi operator  $\bar{R}_X$ , for any tangent vector field  $X$  at  $x \in \bar{M}$ , is defined by

$$(\bar{R}_X Y)(x) = (\bar{R}(Y, X)X)(x)$$

for any  $Y \in T_x \bar{M}$ . It becomes a self adjoint endomorphism of the tangent bundle  $T\bar{M}$  of  $\bar{M}$ , where  $\bar{R}$  denotes the curvature tensor of  $(\bar{M}, \bar{g})$ . That is, the Jacobi operator satisfies  $\bar{R}_X \in \text{End}(T_x \bar{M})$  and is symmetric in the sense of  $\bar{g}(\bar{R}_X Y, Z) = \bar{g}(\bar{R}_X Z, Y)$  for any vector fields  $Y$  and  $Z$  on  $\bar{M}$ .

Let  $M$  be a hypersurface in a Riemannian manifold  $\bar{M}$ . Now by putting a unit normal vector  $N$  into the curvature tensor  $\bar{R}$  of  $\bar{M}$ , the normal Jacobi operator  $\bar{R}_N$  is defined by

$$\bar{R}_N X = \bar{R}(X, N)N$$

for any tangent vector field  $X$  on  $M$  in  $\bar{M}$ .

Related to the commuting problem with the shape operator for real hypersurfaces  $M$  in quaternionic projective space  $\mathbb{H}P^m$  or in quaternionic hyperbolic space  $\mathbb{H}H^m$ , Berndt [1] has introduced the notion of normal Jacobi operator  $\bar{R}_N \in \text{End}(T_x M)$ ,  $x \in M$ , where  $\bar{R}$  denotes the curvature tensor of the ambient spaces  $\mathbb{H}P^m$  or  $\mathbb{H}H^m$ . He [1] showed that the *curvature adaptedness*, when

---

Received November 11, 2011.

2010 *Mathematics Subject Classification.* 53C40, 53C15.

*Key words and phrases.* real hypersurfaces, complex two-plane Grassmannians, recurrent normal Jacobi operator, distribution, Hopf hypersurfaces.

This work is supported by grant Proj. No. NRF-2011-220-C00002 from National Research Foundation of Korea. The first author is supported by Proj. No. BSRP-2012-0004248, the second by MEC-FEDER Grant MTM 2010-18099 and the third by BSRP-2012-R1A2A2A-01043023 and KNU, 2012.

the normal Jacobi operator commutes with the shape operator  $A$ , is equivalent to the fact that the distributions  $\mathfrak{D}$  and  $\mathfrak{D}^\perp = \text{Span}\{\xi_1, \xi_2, \xi_3\}$  are invariant under the shape operator  $A$  of  $M$ , where  $T_x M = \mathfrak{D} \oplus \mathfrak{D}^\perp$ ,  $x \in M$ . Here,  $\{J_\nu \mid \nu = 1, 2, 3\}$  is a canonical local basis of quaternionic Kähler structure  $\mathfrak{J}$  and  $\xi_\nu = -J_\nu N$ ,  $\nu = 1, 2, 3$ . Moreover, he gave a complete classification of curvature adapted real hypersurfaces in quaternionic projective space  $\mathbb{H}P^m$  and in quaternionic hyperbolic space  $\mathbb{H}H^m$ , respectively.

We say that the normal Jacobi operator  $\bar{R}_N$  is *parallel* on  $M$  if the covariant derivative of the normal Jacobi operator  $\bar{R}_N$  identically vanishes, that is,  $\nabla_X \bar{R}_N = 0$  for any vector field  $X$  on  $M$ .

Parallelness of the normal Jacobi operator means that the normal Jacobi operator  $\bar{R}_N$  is parallel on a real hypersurface  $M$  in ambient space  $\bar{M}$ . This means that the eigenspaces of the normal Jacobi operator are parallel along any curve  $\gamma$  in  $M$ . Here the eigenspaces of the normal Jacobi operator  $\bar{R}_N$  are said to be *parallel* along any curve  $\gamma$  if they are *invariant* with respect to any *parallel displacement* along the curve  $\gamma$ .

The complex two-plane Grassmannian  $G_2(\mathbb{C}^{m+2})$  which consists of all complex two-dimensional linear subspaces in  $\mathbb{C}^{m+2}$  has a remarkable geometric structure. It is known to be the unique compact irreducible Riemannian symmetric space equipped with both a Kähler structure  $J$  and a quaternionic Kähler structure  $\mathfrak{J}$  not containing  $J$  (See [2]). From these two structures  $J$  and  $\mathfrak{J}$ , we have geometric conditions naturally induced on a real hypersurface  $M$  in  $G_2(\mathbb{C}^{m+2})$ : That  $[\xi] = \text{Span}\{\xi\}$  or  $\mathfrak{D}^\perp = \text{Span}\{\xi_1, \xi_2, \xi_3\}$  is invariant under the shape operator. From these two conditions, Berndt and Suh [3] have proved the following:

**Theorem A.** *Let  $M$  be a connected real hypersurface in  $G_2(\mathbb{C}^{m+2})$ ,  $m \geq 3$ . Then both  $[\xi]$  and  $\mathfrak{D}^\perp$  are invariant under the shape operator of  $M$  if and only if*

(A)  *$M$  is an open part of a tube around a totally geodesic  $G_2(\mathbb{C}^{m+1})$  in  $G_2(\mathbb{C}^{m+2})$ , or*

(B)  *$m$  is even, say  $m = 2n$ , and  $M$  is an open part of a tube around a totally geodesic  $\mathbb{H}P^n$  in  $G_2(\mathbb{C}^{m+2})$ .*

The structure vector field  $\xi$  of a real hypersurface  $M$  in  $G_2(\mathbb{C}^{m+2})$  is said to be a *Reeb* vector field. If the *Reeb* vector field  $\xi$  of a real hypersurface  $M$  in  $G_2(\mathbb{C}^{m+2})$  is invariant under the shape operator,  $M$  is said to be a *Hopf hypersurface*. In such a case the integral curves of the *Reeb* vector field  $\xi$  are geodesics (See [4]). Moreover, the flow generated by the integral curves of the structure vector field  $\xi$  for Hopf hypersurfaces in  $G_2(\mathbb{C}^{m+2})$  is said to be a *geodesic Reeb flow*.

In paper [9], Jeong, Kim and Suh considered the notion of *parallel* normal Jacobi operator, that is,  $\nabla_X \bar{R}_N = 0$  for any vector field  $X$  on  $M$  in  $G_2(\mathbb{C}^{m+2})$ , where  $\nabla$  denotes the induced connection from the Levi-Civita connection  $\bar{\nabla}$  of  $G_2(\mathbb{C}^{m+2})$ . They proved a non-existence theorem for Hopf hypersurfaces

in complex two-plane Grassmannian  $G_2(\mathbb{C}^{m+2})$ ,  $m \geq 3$ , with parallel normal Jacobi operator as follows:

**Theorem B.** *There does not exist any connected Hopf hypersurface in  $G_2(\mathbb{C}^{m+2})$ ,  $m \geq 3$ , with parallel normal Jacobi operator.*

On the other hand, in [10] Jeong, Lee and Suh have considered a Lie parallelness of the normal Jacobi operator, that is,  $\mathcal{L}_X \bar{R}_N = 0$ , where  $\mathcal{L}_X$  denotes the Lie derivative along any direction  $X$  on  $M$  in  $G_2(\mathbb{C}^{m+2})$ , and asserted the following:

**Theorem C.** *There does not exist any Hopf hypersurface in complex two-plane Grassmannians  $G_2(\mathbb{C}^{m+2})$  with Lie parallel normal Jacobi operator if the integral curves of  $\mathfrak{D}$  and  $\mathfrak{D}^\perp$  components of the Reeb vector field are totally geodesics.*

The purpose of this paper is to study a generalized condition weaker than parallel normal Jacobi operator for real hypersurfaces in  $G_2(\mathbb{C}^{m+2})$ .

Let  $T$  be a tensor field of type (1,1) on  $M$ .  $T$  is said to be recurrent if there exists a certain 1-form  $\omega$  on  $M$  such that for any vector fields  $X, Y$  tangent to  $M$ ,  $(\nabla_X T)(Y) = \omega(X)T(Y)$ . This notion generalizes the fact of  $T$  being parallel (see [13]).

Hamada [5], [6] investigated real hypersurfaces  $M$  in complex projective space  $\mathbb{C}P^m$  with recurrent shape operator. This means that the eigenspaces of the shape operator are parallel along any curve  $\gamma$  in  $M$ . That is, they are invariant with respect to parallel translation along  $\gamma$ . He proved that there does not exist real hypersurface in complex projective space  $\mathbb{C}P^m$  with recurrent shape operator. In [7] he also proved that there does not exist any real hypersurface  $M$  in complex projective space  $\mathbb{C}P^m$  with recurrent Ricci tensor if the structure vector field  $\xi$  is principal.

For a real hypersurface in complex projective space  $\mathbb{C}P^m$ , Pérez and Santos [15] introduced a new notion of  $\mathfrak{D}$ -recurrent, which is weaker than the structure Jacobi operator being recurrent. Here, the structure Jacobi operator  $R_\xi$  is said to be  $\mathfrak{D}$ -recurrent if it satisfies

$$(\nabla_X R_\xi)(Y) = \omega(X)R_\xi(Y),$$

where  $\omega$  and  $\mathfrak{D}$  respectively denote an 1-form on  $M$  and the orthogonal complement of the Reeb vector field  $\xi$  in  $TM$ , and any vector fields  $X \in \mathfrak{D}$ ,  $Y \in TM$ . Namely, they proved the following:

**Theorem D.** *Let  $M$  be a real hypersurface of  $\mathbb{C}P^m$ ,  $m \geq 3$ . Then its structure Jacobi operator is  $\mathfrak{D}$ -recurrent if and only if it is a minimal ruled real hypersurface.*

Related to the shape operator, in paper due to [12], first they have applied Hamada's results to hypersurfaces in complex two-plane Grassmannians  $G_2(\mathbb{C}^{m+2})$  and next obtained a non-existence property for Hopf hypersurfaces in  $G_2(\mathbb{C}^{m+2})$  with recurrent shape operator.

Motivated by such a recurrent shape operator, and in order to make a generalization of Theorem B, in this paper we introduce a new notion of *recurrent Jacobi operator*, that is, the *recurrent normal Jacobi operator* for a real hypersurface  $M$  in complex two-plane Grassmannians  $G_2(\mathbb{C}^{m+2})$ . A hypersurface  $M$  in  $G_2(\mathbb{C}^{m+2})$  with *recurrent normal Jacobi operator* is defined by

$$(\nabla_X \bar{R}_N)(Y) = \omega(X)\bar{R}_N(Y),$$

where  $\omega$  denotes an 1-form on  $M$  and any vector fields  $X, Y$  tangent to  $M$  (see Kobayashi and Nomizu [13] page 305). Consequently, we prove the following:

**Theorem 1.1.** *There does not exist any Hopf hypersurface in complex two-plane Grassmannians  $G_2(\mathbb{C}^{m+2})$  with recurrent normal Jacobi operator.*

## 2. Riemannian geometry of $G_2(\mathbb{C}^{m+2})$

In this section we summarize basic material about  $G_2(\mathbb{C}^{m+2})$ , for details refer to [2], [3] and [4].

By  $G_2(\mathbb{C}^{m+2})$  we denote the set of all complex two-dimensional linear subspaces in  $\mathbb{C}^{m+2}$ . The special unitary group  $G = SU(m+2)$  acts transitively on  $G_2(\mathbb{C}^{m+2})$  with stabilizer isomorphic to  $K = S(U(2) \times U(m)) \subset G$ . The space  $G_2(\mathbb{C}^{m+2})$  can be identified with the homogeneous space  $G/K$ , which we equip with the unique analytic structure for which the natural action of  $G$  on  $G_2(\mathbb{C}^{m+2})$  becomes analytic. Denote by  $\mathfrak{g}$  and  $\mathfrak{k}$  the Lie algebra of  $G$  and  $K$ , respectively, and by  $\mathfrak{m}$  the orthogonal complement of  $\mathfrak{k}$  in  $\mathfrak{g}$  with respect to the Cartan-Killing form  $B$  of  $\mathfrak{g}$ . Then  $\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{m}$  is an  $Ad(K)$ -invariant reductive decomposition of  $\mathfrak{g}$ .

We put  $o = eK$  and identify  $T_oG_2(\mathbb{C}^{m+2})$  with  $\mathfrak{m}$  in the usual manner. Since  $B$  is negative definite on  $\mathfrak{g}$ , negative  $B$  restricted to  $\mathfrak{m} \times \mathfrak{m}$  yields a positive definite inner product on  $\mathfrak{m}$ . By  $Ad(K)$ -invariance of  $B$  this inner product can be extended to a  $G$ -invariant Riemannian metric  $g$  on  $G_2(\mathbb{C}^{m+2})$ .

In this way  $G_2(\mathbb{C}^{m+2})$  becomes a Riemannian homogeneous space, even a Riemannian symmetric space. For computational reasons we normalize  $g$  such that the maximum sectional curvature of  $(G_2(\mathbb{C}^{m+2}), g)$  is eight.

When  $m = 1$ ,  $G_2(\mathbb{C}^3)$  is isometric to the two-dimensional complex projective space  $\mathbb{C}P^2$  with constant holomorphic sectional curvature eight. When  $m = 2$ , we note that the isomorphism  $\text{Spin}(6) \simeq SU(4)$  yields an isometry between  $G_2(\mathbb{C}^4)$  and the real Grassmann manifold  $G_2^+(\mathbb{R}^6)$  of oriented two-dimensional linear subspaces of  $\mathbb{R}^6$ . In this paper, we will assume  $m \geq 3$ .

The Lie algebra  $\mathfrak{k}$  has the direct sum decomposition  $\mathfrak{k} = \mathfrak{su}(m) \oplus \mathfrak{su}(2) \oplus \mathfrak{A}$ , where  $\mathfrak{A}$  is the center of  $\mathfrak{k}$ . Viewing  $\mathfrak{k}$  as the holonomy algebra of  $G_2(\mathbb{C}^{m+2})$ , the center  $\mathfrak{A}$  induces a Kähler structure  $J$  and the  $\mathfrak{su}(2)$ -part a quaternionic Kähler structure  $\mathfrak{J}$  on  $G_2(\mathbb{C}^{m+2})$ .

If  $J_1$  is any almost Hermitian structure in  $\mathfrak{J}$ , then  $JJ_1 = J_1J$ , and  $JJ_1$  is a symmetric endomorphism with  $(JJ_1)^2 = I$  and  $\text{tr}(JJ_1) = 0$ .

A canonical local basis  $J_1, J_2, J_3$  of  $\mathfrak{J}$  consists of three local almost Hermitian structures  $J_\nu$  in  $\mathfrak{J}$  such that  $J_\nu J_{\nu+1} = J_{\nu+2} = -J_{\nu+1} J_\nu$ , where the index is taken modulo three. Since  $\mathfrak{J}$  is parallel with respect to the Riemannian connection  $\bar{\nabla}$  of  $(G_2(\mathbb{C}^{m+2}), g)$ , there exist for any canonical local basis  $J_1, J_2, J_3$  of  $\mathfrak{J}$  three local one-forms  $q_1, q_2, q_3$  such that

$$(2.1) \quad \bar{\nabla}_X J_\nu = q_{\nu+2}(X)J_{\nu+1} - q_{\nu+1}(X)J_{\nu+2}$$

for all vector fields  $X$  on  $G_2(\mathbb{C}^{m+2})$ .

The Riemannian curvature tensor  $\bar{R}$  of  $G_2(\mathbb{C}^{m+2})$  is locally given by

$$(2.2) \quad \begin{aligned} \bar{R}(X, Y)Z &= g(Y, Z)X - g(X, Z)Y + g(JY, Z)JX \\ &\quad - g(JX, Z)JY - 2g(JX, Y)JZ \\ &\quad + \sum_{\nu=1}^3 \{g(J_\nu Y, Z)J_\nu X - g(J_\nu X, Z)J_\nu Y \\ &\quad - 2g(J_\nu X, Y)J_\nu Z\} \\ &\quad + \sum_{\nu=1}^3 \{g(J_\nu JY, Z)J_\nu JX - g(J_\nu JX, Z)J_\nu JY\} \end{aligned}$$

for any vector fields  $X, Y$  and  $Z$  on  $G_2(\mathbb{C}^{m+2})$ , where  $J_1, J_2, J_3$  is any canonical local basis of  $\mathfrak{J}$  [2].

### 3. Some fundamental formulas

In this section we derive some basic formulae from the Codazzi equation for a real hypersurface in  $G_2(\mathbb{C}^{m+2})$  (See [2], [3] and [4]).

Let  $M$  be a real hypersurface of  $G_2(\mathbb{C}^{m+2})$ . The induced Riemannian metric on  $M$  will also be denoted by  $g$ , and  $\nabla$  denotes the Riemannian connection of  $(M, g)$ . Let  $N$  be a local unit normal field of  $M$  and  $A$  the shape operator of  $M$  with respect to  $N$ . The Kähler structure  $J$  of  $G_2(\mathbb{C}^{m+2})$  induces on  $M$  an almost contact metric structure  $(\phi, \xi, \eta, g)$ . Furthermore, let  $J_1, J_2, J_3$  be a canonical local basis of  $\mathfrak{J}$ . Then each  $J_\nu$  induces an almost contact metric structure  $(\phi_\nu, \xi_\nu, \eta_\nu, g)$  on  $M$ . Using the above expression for  $\bar{R}$ , the Codazzi equation becomes

$$\begin{aligned} (\nabla_X A)Y - (\nabla_Y A)X &= \eta(X)\phi Y - \eta(Y)\phi X - 2g(\phi X, Y)\xi \\ &\quad + \sum_{\nu=1}^3 \{ \eta_\nu(X)\phi_\nu Y - \eta_\nu(Y)\phi_\nu X - 2g(\phi_\nu X, Y)\xi_\nu \} \\ &\quad + \sum_{\nu=1}^3 \{ \eta_\nu(\phi X)\phi_\nu \phi Y - \eta_\nu(\phi Y)\phi_\nu \phi X \} \\ &\quad + \sum_{\nu=1}^3 \{ \eta(X)\eta_\nu(\phi Y) - \eta(Y)\eta_\nu(\phi X) \} \xi_\nu . \end{aligned}$$

The following identities can be proved in a straightforward method and will be used frequently in subsequent calculations:

$$(3.1) \quad \begin{aligned} \phi_{\nu+1}\xi_\nu &= -\xi_{\nu+2}, & \phi_\nu\xi_{\nu+1} &= \xi_{\nu+2}, \\ \phi\xi_\nu &= \phi_\nu\xi, & \eta_\nu(\phi X) &= \eta(\phi_\nu X), \\ \phi_\nu\phi_{\nu+1}X &= \phi_{\nu+2}X + \eta_{\nu+1}(X)\xi_\nu, \\ \phi_{\nu+1}\phi_\nu X &= -\phi_{\nu+2}X + \eta_\nu(X)\xi_{\nu+1}. \end{aligned}$$

Now let us put

$$(3.2) \quad JX = \phi X + \eta(X)N, \quad J_\nu X = \phi_\nu X + \eta_\nu(X)N$$

for any vector field  $X$  tangent to a real hypersurface  $M$  in  $G_2(\mathbb{C}^{m+2})$ , where  $N$  denotes a normal vector field of  $M$  in  $G_2(\mathbb{C}^{m+2})$ . Then from this and the formulas (2.1) and (3.1) we have that

$$(3.3) \quad (\nabla_X \phi)Y = \eta(Y)AX - g(AX, Y)\xi, \quad \nabla_X \xi = \phi AX,$$

$$(3.4) \quad \nabla_X \xi_\nu = q_{\nu+2}(X)\xi_{\nu+1} - q_{\nu+1}(X)\xi_{\nu+2} + \phi_\nu AX,$$

$$(3.5) \quad \begin{aligned} (\nabla_X \phi_\nu)Y &= -q_{\nu+1}(X)\phi_{\nu+2}Y + q_{\nu+2}(X)\phi_{\nu+1}Y + \eta_\nu(Y)AX \\ &\quad - g(AX, Y)\xi_\nu. \end{aligned}$$

Moreover, from  $JJ_\nu = J_\nu J$ ,  $\nu = 1, 2, 3$ , it follows that

$$(3.6) \quad \phi\phi_\nu X = \phi_\nu\phi X + \eta_\nu(X)\xi - \eta(X)\xi_\nu.$$

#### 4. Recurrent normal Jacobi operator

From (2.2) the normal Jacobi operator  $\bar{R}_N$  of  $M$  is given by

$$\begin{aligned} \bar{R}_N(X) &= \bar{R}(X, N)N \\ &= X + 3\eta(X)\xi + 3\sum_{\nu=1}^3 \eta_\nu(X)\xi_\nu \\ &\quad - \sum_{\nu=1}^3 \left\{ \eta_\nu(\xi)(\phi_\nu\phi X - \eta(X)\xi_\nu) - \eta_\nu(\phi X)\phi_\nu\xi \right\}. \end{aligned}$$

Now let us consider the covariant derivative of the normal Jacobi operator  $\bar{R}_N$  along the direction  $X$  (see [9]). It is given by

$$\begin{aligned} (\nabla_X \bar{R}_N)Y &= 3g(\phi AX, Y)\xi + 3\eta(Y)\phi \\ &\quad + 3\sum_{\nu=1}^3 \left\{ g(\phi_\nu AX, Y)\xi_\nu + \eta_\nu(Y)\phi_\nu AX \right\} \\ &\quad - \sum_{\nu=1}^3 \left[ 2\eta_\nu(\phi AX)(\phi_\nu\phi Y - \eta(Y)\xi_\nu) - g(\phi_\nu AX, \phi Y)\phi_\nu\xi \right. \\ &\quad \left. - \eta(Y)\eta_\nu(AX)\phi_\nu\xi - \eta_\nu(\phi Y)(\phi_\nu\phi AX - g(AX, \xi)\xi_\nu) \right]. \end{aligned}$$

From this, together with formulas given in Section 3, a real hypersurface  $M$  in  $G_2(\mathbb{C}^{m+2})$  with recurrent normal Jacobi operator satisfies the following

$$\begin{aligned}
 & 3g(\phi AX, Y)\xi + 3\eta(Y)\phi AX + 3\sum_{\nu=1}^3 \left\{ g(\phi_\nu AX, Y)\xi_\nu + \eta_\nu(Y)\phi_\nu AX \right\} \\
 & - \sum_{\nu=1}^3 \left[ 2\eta_\nu(\phi AX)(\phi_\nu \phi Y - \eta(Y)\xi_\nu) - g(\phi_\nu AX, \phi Y)\phi_\nu \xi \right. \\
 (4.1) \quad & \left. - \eta(Y)\eta_\nu(AX)\phi_\nu \xi - \eta_\nu(\phi Y)(\phi_\nu \phi AX - g(AX, \xi)\xi_\nu) \right] \\
 & = \omega(X) \left[ Y + 3\eta(Y)\xi + 3\sum_{\nu=1}^3 \eta_\nu(Y)\xi_\nu \right. \\
 & \quad \left. - \sum_{\nu=1}^3 \left\{ \eta_\nu(\xi)(\phi_\nu \phi Y - \eta(Y)\xi_\nu) - \eta_\nu(\phi Y)\phi_\nu \xi \right\} \right].
 \end{aligned}$$

In order to prove our Main Theorem in the introduction, we give the following.

**Lemma 4.1.** *Let  $M$  be a Hopf hypersurface in  $G_2(\mathbb{C}^{m+2})$  with recurrent normal Jacobi operator. Then the Reeb vector field  $\xi$  belongs to either the distribution  $\mathfrak{D}$  or the distribution  $\mathfrak{D}^\perp$ .*

*Proof.* From (4.1), we take  $X = Y = \xi$  and suppose that  $M$  is Hopf, that is,  $A\xi = \alpha\xi$  for a certain function  $\alpha$ . Then this yields

$$(4.2) \quad \alpha \sum_{\nu=1}^3 \eta_\nu(\xi)\phi_\nu \xi = \omega(\xi)(\xi + \sum_{\nu=1}^3 \eta_\nu(\xi)\xi_\nu).$$

Taking its scalar product with  $\xi$  we get  $\omega(\xi) = 0$ . As  $\omega(\xi) = 0$ , (4.2) yields

$$(4.3) \quad \alpha \sum_{\nu=1}^3 \eta_\nu(\xi)\phi_\nu \xi = 0.$$

From this, we consider the case that the function  $\alpha$  is non-vanishing. Now let us put  $\xi = \eta(X_0)X_0 + \eta(\xi_1)\xi_1$  for some unit  $X_0 \in \mathfrak{D}$  and non-vanishing functions  $\eta(X_0)$  and  $\eta(\xi_1)$ .

Then (4.3) yields

$$0 = \eta(\xi_1)\phi_1 \xi = \eta(X_0)\eta(\xi_1)\phi_1 X_0.$$

This gives a contradiction with  $\eta(\xi_1) \neq 0$  and  $\eta(X_0) \neq 0$ . So we get  $\eta(\xi_1) = 0$  or  $\eta(X_0) = 0$ , which means  $\xi \in \mathfrak{D}$  or  $\xi \in \mathfrak{D}^\perp$ .

When the function  $\alpha$  vanishes, we can differentiate  $A\xi = 0$ . Then by a theorem due to Berndt and Suh [4] we know that

$$\sum_{\nu=1}^3 \eta_\nu(\xi)\phi_\nu \xi = 0.$$

This also gives  $\xi \in \mathfrak{D}$  or  $\xi \in \mathfrak{D}^\perp$ . □

### 5. Recurrent normal Jacobi operator with $\xi \in \mathfrak{D}$

In paper [14], Lee and Suh gave a characterization of real hypersurfaces of type B in  $G_2(\mathbb{C}^{m+2})$  in terms of the Reeb vector field  $\xi \in \mathfrak{D}$ . Now we introduce the following:

**Proposition 5.1.** *Let  $M$  be a connected orientable Hopf hypersurface in  $G_2(\mathbb{C}^{m+2})$ . If the Reeb vector field  $\xi$  belongs to the distribution  $\mathfrak{D}$ , then the distribution  $\mathfrak{D}$  is invariant under the shape operator  $A$  of  $M$ , that is,  $g(A\mathfrak{D}, \mathfrak{D}^\perp) = 0$ .*

Then by Proposition 5.1 and Theorem A in the introduction, we know naturally that a Hopf hypersurface  $M$  in  $G_2(\mathbb{C}^{m+2})$  with recurrent normal Jacobi operator and  $\xi \in \mathfrak{D}$  is a tube over a totally geodesic quaternionic projective space  $\mathbb{H}P^n$ ,  $m = 2n$ .

Now let us check if a real hypersurface of type (B) in  $G_2(\mathbb{C}^{m+2})$ , that is, a tube over a totally geodesic  $\mathbb{H}P^n$ , satisfies the notion of recurrent normal Jacobi operator. Corresponding to such a real hypersurface of type (B), we introduce a proposition due to Berndt and Suh [3] as follows:

**Proposition 5.2.** *Let  $M$  be a connected real hypersurface of  $G_2(\mathbb{C}^{m+2})$ . Suppose that  $A\mathfrak{D} \subset \mathfrak{D}$ ,  $A\xi = \alpha\xi$ , and  $\xi$  is tangent to  $\mathfrak{D}$ . Then the quaternionic dimension  $m$  of  $G_2(\mathbb{C}^{m+2})$  is even, say  $m = 2n$ , and  $M$  has five distinct constant principal curvatures*

$$\alpha = -2 \tan(2r), \quad \beta = 2 \cot(2r), \quad \gamma = 0, \quad \lambda = \cot(r), \quad \mu = -\tan(r)$$

with some  $r \in (0, \pi/4)$ . The corresponding multiplicities are

$$m(\alpha) = 1, \quad m(\beta) = 3 = m(\gamma), \quad m(\lambda) = 4n - 4 = m(\mu)$$

and the corresponding eigenspaces are

$$T_\alpha = \mathbb{R}\xi, \quad T_\beta = \mathfrak{J}J\xi, \quad T_\gamma = \mathfrak{J}\xi, \quad T_\lambda, \quad T_\mu,$$

where

$$T_\lambda \oplus T_\mu = (\mathbb{H}\mathbb{C}\xi)^\perp, \quad \mathfrak{J}T_\lambda = T_\lambda, \quad \mathfrak{J}T_\mu = T_\mu, \quad JT_\lambda = T_\mu.$$

Now let us suppose  $M$  is of type (B) with recurrent normal Jacobi operator  $\bar{R}_N$ . From (4.1), by putting  $X = \xi_2$  and  $Y = \xi$  we have

$$\omega(\xi_2)\xi - \beta\phi_2\xi = 0.$$

Then it follows that  $\omega(\xi_2) = 0$  and  $\beta = 0$ . Since  $\beta$  is not zero, this makes a contradiction. Thus we conclude the following:

**Theorem 5.1.** *There does not exist any Hopf hypersurface in  $G_2(\mathbb{C}^{m+2})$ ,  $m \geq 3$ , with recurrent normal Jacobi operator if the Reeb vector  $\xi$  belongs to the distribution  $\mathfrak{D}$ .*

**6. Recurrent normal Jacobi operator with  $\xi \in \mathfrak{D}^\perp$**

In this section, we consider the case that  $\xi \in \mathfrak{D}^\perp$ . Then the unit normal vector field  $N$  is a singular tangent vector of  $G_2(\mathbb{C}^{m+2})$  of type  $JN \in \mathfrak{J}N$ . So there exists an almost Hermitian structure  $J_1 \in \mathfrak{J}$  such that  $JN = J_1N$ . Then we have

$$\xi = \xi_1, \phi\xi_2 = -\xi_3, \phi\xi_3 = \xi_2, \phi\mathfrak{D} \subset \mathfrak{D}.$$

Then, by putting  $X = \xi_\mu$  and  $Y = \xi$  in (4.1), we get

$$3\phi A\xi_\mu + 5\sum_{\nu=1}^3 \eta_\nu(\phi A\xi_\mu)\xi_\nu + 3\phi_1 A\xi_\mu + \sum_{\nu=1}^3 \eta_\nu(A\xi_\mu)\phi_\nu\xi = 8\omega(\xi_\mu)\xi.$$

From this, by taking its scalar product with Reeb vector field  $\xi$  we get  $\omega(\xi_\mu) = 0$ . As  $\omega(\xi_\mu) = 0$ , we have

$$\begin{aligned} 0 &= (\nabla_{\xi_\mu} \bar{R}_N)X \\ &= 3g(\phi A\xi_\mu, X)\xi + 3\eta(X)\phi A\xi_\mu \\ &\quad + 3\sum_{\nu=1}^3 \left\{ g(\phi_\nu A\xi_\mu, X)\xi_\nu + \eta_\nu(X)\phi_\nu A\xi_\mu \right\} \\ &\quad - \sum_{\nu=1}^3 \left[ 2\eta_\nu(\phi A\xi_\mu)(\phi_\nu\phi X - \eta(X)\xi_\nu) - g(\phi_\nu A\xi_\mu, \phi X)\phi_\nu\xi \right. \\ &\quad \left. - \eta(X)\eta_\nu(A\xi_\mu)\phi_\nu\xi - \eta_\nu(\phi X)(\phi_\nu\phi A\xi_\mu - g(A\xi_\mu, \xi)\xi_\nu) \right] \end{aligned}$$

for any  $X \in TM$ . From this, by putting  $X = \xi$  and using  $\xi = \xi_1$ , we have

$$0 = 3\phi A\xi_\mu + 5\sum_{\nu=1}^3 \eta_\nu(\phi A\xi_\mu)\xi_\nu + 3\phi_1 A\xi_\mu + \sum_{\nu=1}^3 \eta_\nu(A\xi_\mu)\phi_\nu\xi.$$

From this, taking its inner product with  $X \in \mathfrak{D}$  and using  $g(\phi_\nu\xi, X) = 0$ , we obtain

$$(6.1) \quad 0 = 3g(\phi A\xi_\mu, X) + 3g(\phi_1 A\xi_\mu, X).$$

On the other hand, by using (3.4) we know that

$$\phi A\xi_\mu = \nabla_{\xi_\mu} \xi = \nabla_{\xi_\mu} \xi_1 = q_3(\xi_\mu)\xi_2 - q_2(\xi_\mu)\xi_3 + \phi_1 A\xi_\mu.$$

From this, taking its inner product with  $X \in \mathfrak{D}$ , we have

$$g(\phi A\xi_\mu, X) = g(\phi_1 A\xi_\mu, X).$$

Substituting this formula into (6.1) gives

$$0 = g(\phi A\xi_\mu, X).$$

From this, let us replace  $X$  by  $\phi X$ . Then it follows that

$$0 = g(\phi A\xi_\mu, \phi X) = -g(A\xi_\mu, \phi^2 X) = g(AX, \xi_\mu)$$

for any vector field  $X \in \mathfrak{D}$ ,  $\mu = 1, 2, 3$ .

Then we obtain the following:

**Lemma 6.1.** *Let  $M$  be a Hopf hypersurface in  $G_2(\mathbb{C}^{m+2})$ ,  $m \geq 3$ , with recurrent normal Jacobi operator and  $\xi \in \mathfrak{D}^\perp$ . Then  $g(A\mathfrak{D}, \mathfrak{D}^\perp) = 0$ .*

From this together with Theorem A in the introduction we know that any real hypersurface in  $G_2(\mathbb{C}^{m+2})$  with recurrent normal Jacobi operator  $\bar{R}_N$  and  $\xi \in \mathfrak{D}^\perp$  is congruent to a tube over a totally geodesic  $G_2(\mathbb{C}^{m+1})$  in  $G_2(\mathbb{C}^{m+2})$ .

Now let us check if real hypersurfaces of type (A) satisfy the condition of recurrent normal Jacobi operator. Then we recall a proposition given by Berndt and Suh [3] as follows:

**Proposition 6.1.** *Let  $M$  be a connected real hypersurface of  $G_2(\mathbb{C}^{m+2})$ . Suppose that  $A\mathfrak{D} \subset \mathfrak{D}$ ,  $A\xi = \alpha\xi$ , and  $\xi$  is tangent to  $\mathfrak{D}^\perp$ . Let  $J_1 \in \mathfrak{J}$  be the almost Hermitian structure such that  $JN = J_1N$ . Then  $M$  has three (if  $r = \pi/2\sqrt{8}$ ) or four (otherwise) distinct constant principal curvatures*

$$\alpha = \sqrt{8} \cot(\sqrt{8}r), \quad \beta = \sqrt{2} \cot(\sqrt{2}r), \quad \lambda = -\sqrt{2} \tan(\sqrt{2}r), \quad \mu = 0$$

with some  $r \in (0, \pi/\sqrt{8})$ . The corresponding multiplicities are

$$m(\alpha) = 1, \quad m(\beta) = 2, \quad m(\lambda) = 2m - 2 = m(\mu),$$

and as corresponding eigenspaces we have

$$\begin{aligned} T_\alpha &= \mathbb{R}\xi = \mathbb{R}JN = \mathbb{R}\xi_1, \\ T_\beta &= \mathbb{C}^\perp\xi = \mathbb{C}^\perp N = \mathbb{R}\xi_2 \oplus \mathbb{R}\xi_3, \\ T_\lambda &= \{X \mid X \perp \mathbb{H}\xi, JX = J_1X\}, \\ T_\mu &= \{X \mid X \perp \mathbb{H}\xi, JX = -J_1X\}, \end{aligned}$$

where  $\mathbb{R}\xi$ ,  $\mathbb{C}\xi$  and  $\mathbb{H}\xi$  respectively denotes real, complex and quaternionic span of the structure vector  $\xi$  and  $\mathbb{C}^\perp\xi$  denotes the orthogonal complement of  $\mathbb{C}\xi$  in  $\mathbb{H}\xi$ .

Now let us suppose  $M$  is of type (A) with recurrent normal Jacobi operator  $\bar{R}_N$  and  $\xi \in \mathfrak{D}^\perp$ . From (4.1), by putting  $X = \xi_2$  and  $Y = \xi$  we have

$$8\omega(\xi_2)\xi - 6\beta\xi_3 = 0.$$

Then it follows that  $\omega(\xi_2) = 0$  and  $\beta = 0$ . Since  $\beta$  is not zero, this gives a contradiction. Thus we conclude the following:

**Theorem 6.1.** *There does not exist any Hopf hypersurface in  $G_2(\mathbb{C}^{m+2})$ ,  $m \geq 3$ , with recurrent normal Jacobi operator if the Reeb vector  $\xi$  belongs to the distribution  $\mathfrak{D}^\perp$ .*

Accordingly, by Lemma 4.1 together with Theorems 5.1 and 6.1 we give a complete proof of our Theorem 1.1 mentioned in the introduction.

## References

- [1] J. Berndt, *Real hypersurfaces in quaternionic space forms*, J. Reine Angew. Math. **419** (1991), 9–26.
- [2] ———, *Riemannian geometry of complex two-plane Grassmannians*, Rend. Sem. Mat. Univ. Politec. Torino **55** (1997), no. 1, 19–83.
- [3] J. Berndt and Y. J. Suh, *Real hypersurfaces in complex two-plane Grassmannians*, Monatsh. Math. **127** (1999), no. 1, 1–14.
- [4] ———, *Real hypersurfaces with isometric Reeb flow in complex two-plane Grassmannians*, Monatsh. Math. **137** (2002), no. 2, 87–98.
- [5] T. Hamada, *On real hypersurfaces of a complex projective space with recurrent second fundamental tensor*, J. Ramanujan Math. Soc. **11** (1996), no. 2, 103–107.
- [6] ———, *On real hypersurfaces of a complex projective space with recurrent Ricci tensor*, Glasg. Math. J. **41** (1999), no. 3, 297–302.
- [7] ———, *Note on real hypersurfaces of complex space forms with recurrent Ricci tensor*, Differ. Geom. Dyn. Syst. **5** (2003), no. 1, 27–30.
- [8] I. Jeong and Y. J. Suh, *Real hypersurfaces in complex two-plane Grassmannians with Lie  $\xi$ -parallel normal Jacobi operator*, J. Korean Math. Soc. **45** (2008), no. 4, 1113–1133.
- [9] I. Jeong, H. J. Kim, and Y. J. Suh, *Real hypersurfaces in complex two-plane Grassmannians with parallel normal Jacobi operator*, Publ. Math. Debrecen **76** (2010), no. 1-2, 203–218.
- [10] I. Jeong, H. Lee, and Y. J. Suh, *Hopf hypersurfaces in complex two-plane Grassmannians with Lie parallel normal Jacobi operator*, Bull. Korean Math. Soc. **48** (2011), no. 2, 427–444.
- [11] I. Jeong, J. D. Pérez, and Y. J. Suh, *Real hypersurfaces in complex two-plane Grassmannians with commuting normal Jacobi operator*, Acta Math. Hungar. **117** (2007), no. 3, 201–217.
- [12] S. Kim, H. Lee, and H. Y. Yang, *Real hypersurfaces in complex two-plane Grassmannians with recurrent shape operator*, Bull. Malays. Math. Sci. Soc. (2) **34** (2011), no. 2, 295–305.
- [13] S. Kobayashi and K. Nomizu, *Foundations of Differential Geometry. Vol. I*, Wiley Classics Library Edition Publ., 1996.
- [14] H. Lee and Y. J. Suh, *Real hypersurfaces of type B in complex two-plane Grassmannians related to the Reeb vector*, Bull. Korean Math. Soc. **47** (2010), no. 3, 551–561.
- [15] J. D. Pérez and F. G. Santos, *Real hypersurfaces in complex projective space with recurrent structure Jacobi operator*, Differential Geom. Appl. **26** (2008), no. 2, 218–223.

IMSOON JEONG  
 DEPARTMENT OF MATHEMATICS  
 KYUNGPOOK NATIONAL UNIVERSITY  
 TAEGU 702-701, KOREA  
*E-mail address:* imsoon.jeong@gmail.com

JUAN DE DIOS PÉREZ  
 DEPARTAMENTO DE GEOMETRÍA Y TOPOLOGÍA  
 FACULTAD DE CIENCIAS  
 UNIVERSIDAD DE GRANADA  
 18071-GRANADA, SPAIN  
*E-mail address:* jdperez@ugr.es

YOUNG JIN SUH  
DEPARTMENT OF MATHEMATICS  
KYUNGPOOK NATIONAL UNIVERSITY  
TAEGU 702-701, KOREA  
*E-mail address:* [yjsuh@knu.ac.kr](mailto:yjsuh@knu.ac.kr)