

Clinical Article

A Comparison of the Clinical Outcomes of Decompression Alone and Fusion in Elderly Patients with Two-Level or More Lumbar Spinal Stenosis

Seong Son, M.D., Woo Kyung Kim, M.D., Ph.D., Sang Gu Lee, M.D., Ph.D., Chan Woo Park, M.D., Ph.D., Keun Lee, M.D.

Department of Neurosurgery, Gachon University Gil Medical Center, Incheon, Korea

Objective : We compared the results of two surgical techniques by retrospective study of 60 elderly patients (65 years or older) who underwent either decompression alone or fusion for the treatment of two-level or more lumbar spinal stenosis.

Methods : During the period of 2003 and 2008, two-level or more decompression alone or fusion was performed for lumbar spinal stenosis by three surgeons at our institution. Patients were allocated to two groups by surgical modality, namely, to a decompression group (31 patients) or a fusion group (29 patients). Overall mean age was 71.1 years (range, 65-84) and mean follow-up was 5.5 years (range, 3-9). A retrospective review of clinical, radiological, and surgical data was conducted.

Results : No significant difference between the two groups was found with respect to age, follow-up period, surgical levels, or preoperative condition. At the last follow-up, correction of lumbar lordotic angle (determined radiologically) was better in the fusion group. However, clinical outcomes including visual analogue scale, Oswestry Disability Index, and the Odom's criteria were not significantly different in the two groups. On the other hand, surgical outcomes, such as, operation time, estimated blood loss, and surgical complications were significantly better in the decompression alone group.

Conclusion : Our findings suggest that decompressive laminectomy alone achieves good outcomes in patients with two-level or more lumbar spinal stenosis, associated with an advanced age, poor general condition, or osteoporosis.

Key Words : Laminectomy · Spinal fusion · Spinal stenosis.

INTRODUCTION

Developments in modern medicine have significantly increased life expectancy, and the resulting population aging means that neurosurgeons are being increasingly confronted with older patients suffering from lumbar spinal stenosis caused by degenerative changes of the lumbar spine^{3,7,19}.

Surgery should be considered when conservative therapy fails to improve the symptoms of lumbar spinal stenosis, and decompressive laminectomy with or without fusion is the standard surgical treatment for patients with two-level or more lumbar spinal stenosis.

Two-level or more decompressive laminectomy with fusion effectively ensures spinal stability, but has significant adverse

events, such as, postoperative complications, instrument failure, and adjacent segment degeneration (ASD). To avoid these adverse events, two-level or more decompressive laminectomy alone has been attempted for lumbar spinal stenosis^{18,20}. However, decompression alone can provoke lumbar instability and aggravate symptoms after surgery.

Some reports have been issued on decompression alone for patients with single-level stenosis^{12,17,22}, and on decompression alone in elderly patients⁹. However, comparative studies of decompression alone and fusion in elderly patients with two-level or more lumbar spinal stenosis were rare.

Accordingly, we retrospectively analyzed clinical, radiologic, and surgical outcomes following decompression alone or fusion in elderly patients with two-level or more lumbar spinal stenosis.

• Received : August 20, 2012 • Revised : November 14, 2012 • Accepted : January 15, 2013

• Address for reprints : Woo Kyung Kim, M.D., Ph.D.

Department of Neurosurgery, Gachon University, Gil Hospital, 21 Namdong-daero, Namdong-gu, Incheon 405-760, Korea

Tel : +82-32-460-3304, Fax : +82-32-460-3899, E-mail : ns@gilhospital.com

• This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

MATERIALS AND METHODS

Patients

Sixty patients, aged 65 years or older, who received multi-level (two-level or more) surgery at our Spine Center from June 2003 to December 2008 were selected for this study. Three spine surgeons participated in this study.

The clinical indications for surgery were radiating leg pain and/or neurogenic intermittent claudication (NIC) with or without low back pain resistant to conservative treatment. Magnetic resonance imaging (MRI) was used to confirm the diagnosis of central stenosis or lateral stenosis. The selection of surgical methods depends on each surgeon's clinical impression that which would elicit a better outcome, based on symptom, general condition, and the degree of osteoporosis. The degree of osteoporosis was evaluated by subjective judgment based on imaging study or by

Table 1. American Society of Anesthesiologists classification

Class	Definition
I	No systemic disease
II	Mild to moderate systemic disease
III	Severe systemic disease
IV	Severe systemic disease that is life threatening
V	Moribund patient with little chance of survival



Fig. 1. Lateral plain radiograph of the lumbar spine showing Cobb's method. Lumbar lordotic angle was determined at the intersection of lines drawn at the level of the inferior plateau of T12 and the superior plateau of S1.

dual-energy X-ray absorptiometry (DEXA) if needed.

The general indications of decompression alone were patients with overwhelming main symptom of radiating pain or NIC rather than low back pain, patients who successful fusion was not expected due to severe osteoporosis, and patients with intolerable general condition to fusion surgery. Whereas, the general indications of fusion were patients with relatively severe low back pain suspected mechanical back pain due to degeneration without overt segmental instability, patients who successful fusion was expected due to tolerable bone marrow density, and patients with tolerable general condition to fusion surgery.

Patients with no choice but fusion surgery, for example, patients with overt segmental instability (defined by White and Panjabi²³) or severe spondylolisthesis (grade II or more), were excluded. Other exclusion criteria included infectious disease, traumatic lumbar disease, developmental spinal deformities, metabolic bone disease, and tumors.

There were 27 men and 33 women, and overall mean age for surgery was 71.1 years (range, 65-84). Mean follow-up was 5.5 years (range, 3.0-9.0). Patients followed up for less than 3 years were excluded. Fifty patients underwent surgery at 2 levels, 8 at 3 levels, and 2 at 4 levels.

Outcome parameters

Preoperative conditions were assessed using the American Society of Anesthesiologists (ASA) classification of physical status (Table 1)⁸, and clinical outcomes were assessed using a visual analogue scale (VAS) for low back pain, a VAS for leg pain, the Oswestry Disability Index (ODI), and the Odom's criteria. VAS scores were determined using 0 to 10 point scales, where a score of 0 means symptom-free, and a score of 10 means the most serious symptom. VAS and ODI were scored preoperatively and at 6 weeks, 6 months, 1 year, and 3 years postoperatively. Odom's criteria were used to evaluate patient satisfaction at final follow-up.

Standard anterior-posterior and lateral radiographs of the lumbar spine were taken preoperatively and during follow-up for all patients. Lumbar lordotic angle was measured on a lateral radiograph in the neutral position using Cobb's method (Fig. 1). Preoperative and postoperative lumbar lordotic angles were compared at 6 weeks, 6 months, 1 year, and 3 years after surgery.

Surgical methods were compared with respect to estimated blood losses (EBL), operation times, and hospital stays. The occurrences of perioperative morbidities [e.g. neurologic deterioration, cerebrospinal fluid (CSF) leakage, wound infection, pneumonia, heart problem, urinary difficulty, epidural hematoma, and deep vein thrombosis] were checked. In addition, reoperation and developments of late postoperative complications (e.g. recurrence, instability, subsidence, screw loosening, non-union, and ASD) were documented.

Surgical technique

All operations were performed in the prone position. In cases of

decompression alone, after a traditional median incision, partial laminectomy and decompressive ligamentectomy were performed at each symptomatic stenotic level. Unilateral foraminotomy, bilateral foraminotomy, or unilateral laminotomy and bilateral decompression were performed according to symptoms and MRI findings.

In cases of fusion, after decompression and discectomy, posterior lumbar interbody fusion (PLIF) or transforaminal lumbar interbody fusion (TLIF) was performed, and followed by two-level or more transpedicular screw fixation (percutaneous or open).

Statistical methods

SPSS version 16.0 (SPSS Inc., Chicago, IL, USA) was used to analyze all data. The chi square test, the independent 2-sample t-test, and the one-way analysis of variance were used depending on the characteristics of the variables being compared. Statistical significance was accepted for p values of <0.05 .

RESULTS

Demographic and preoperative data

The patients were divided into two groups according to surgical technique. Of the 60 patients, 31 (51.7%) were allocated to the decompression group, and 29 (48.3%) to the fusion group.

No significant intergroup difference was found with respect to age, sex ratio, follow-up period, or surgical levels (Table 2). There was no significant difference with respect to ASA classification, preoperative ODI, or preoperative lumbar lordotic angle. Although there was no statistical significance, ASA classification tended to be worse in the decompression group, whereas preoperative ODI and lumbar lordotic angle tended to be worse in the fusion group. Also, unlike preoperative VAS for leg pain, preoperative VAS for low back pain of fusion group was higher than that of decompression group ($p<0.05$) (Table 3).

Unfortunately, the objective comparison of degree of osteoporosis was impossible because DEXAs were not evenly performed in the two groups.

Clinical outcome

In both groups, VAS for low back pain and leg pain was decreased during follow-up, sequentially.

In decompression group, VAS for low back pain was im-

Table 2. Summary of demographic data of both groups

Variable	Decompression group	Fusion group	p -value
Number of patients	31	29	
Mean age (years)	72.8±6.8	69.4±3.8	0.497
Female sex ratio (%)	48.4	62.1	0.200
Mean follow-up period (years)	5.8±1.7	5.2±3.1	0.866
Number of levels of operation			0.753
2 levels	28	22	
3 levels	3	5	
4 levels	0	2	
Levels of operation			0.599
L2-3	0	7	
L3-4	16	18	
L4-5	31	29	
L5-S1	18	11	

Table 3. Summary of preoperative data of both groups

Variable	Decompression group (n=31)	Fusion group (n=29)	p -value
ASA classification			0.570
I	6 (19.4%)	7 (24.1%)	
II	21 (67.7%)	20 (69.0%)	
III	4 (12.9%)	2 (6.9%)	
IV	-	-	
V	-	-	
Pre-OP VAS for low back pain	5.9±1.3	7.1±1.1	0.017
Pre-OP VAS for leg pain	7.4±1.2	7.5±1.3	0.857
Pre-OP ODI	68.6±6.8	63.1±12.3	0.246
Pre-OP lumbar lordotic angle	35.3±11.5	32.3±13.4	0.177

ASA : American Society of Anesthesiologists, pre-OP : preoperative, VAS : visual analogue scale, ODI : Oswestry Disability Index

proved from 5.9 to 3.1 ($p<0.05$). Also, in fusion group, VAS for low back pain was improved from 7.1 to 3.2 ($p<0.05$) (Fig. 2).

In decompression group, the mean preoperative VAS for leg pain of 7.4 decreased to 2.9 at last follow-up ($p<0.05$). Also, in fusion group, the mean preoperative VAS for leg pain of 7.5 decreased to 3.1 at last follow-up ($p<0.05$) (Fig. 3).

Follow-up VAS scores were not significantly different in two groups, but decompression group showed a better improvement in low back pain VAS scores at 6 weeks after surgery.

The functional aspects were evaluated using ODI scores. In both groups, ODI decreased during follow-up, sequentially ($p<0.05$). ODI scores were not significantly different in two groups (Fig. 4).

Odom's criteria failed to reveal a significant intergroup difference at last follow-up (Table 4).

Radiological outcomes

In decompression group, the mean preoperative lumbar lordotic angle of 35.3° decreased to 34.8° at last follow-up. However, the changes of angles were not significant.

On the other hand, in fusion group, the mean preoperative




Fig. 2. Sequential changes in mean low back pain VAS scores. Mean preoperative VAS for low back pain was 5.9 in decompression group, and 7.1 in fusion group. In decompression group, VAS for low back pain decreased to 3.2 at 6 weeks, 3.2 at 6 months, and to 3.1 at 1 year. In fusion group, VAS for low back pain decreased to 6.1 at 6 weeks, 3.8 at 6 months, 3.5 at 1 year, and to 3.2 at 3 years. f/u : follow-up, pre OP : preoperation, VAS : visual analogue scale.




Fig. 3. Sequential changes in mean leg pain VAS scores. The mean preoperative VAS for leg back pain was 7.4 in decompression group, and 7.5 in fusion group. In decompression group, VAS for leg pain decreased to 3.4 at 6 weeks, 3.1 at 6 months, 3.0 at 1 year, and to 2.9 at 3 years. In fusion group, VAS for leg pain decreased to 3.3 at 6 weeks, 3.2 at 6 months, and to 3.1 at 3 years. f/u : follow-up, pre OP : preoperation, VAS : visual analogue scale.

lumbar lordotic angle of 32.3° increased to 37.5° at last follow-up ($p < 0.05$) (Fig. 5).

Comparison of surgical methods and complications (Table 5)

The EBL, operation time, and length of hospital stay were significantly better in the decompression group ($p < 0.05$).

There was no postoperative mortality and no neurologic deterioration after surgery. Perioperative morbidity was noted in 3 patients (9.7%) in the decompression group, and in 7 patients (24.1%) in the fusion group. In decompression group, CSF leakage in 1 patient, wound infection in 1 patient, and facet fracture in 1 patient were occurred. In fusion group, CSF leakage in 2 patients, wound infection in 2 patients, epidural hematoma in 1 patient, pneumonia in 1 patient, urinary difficulty in




Fig. 4. Sequential changes in mean ODI scores. Mean preoperative ODI was 63.1 in decompression group, and 68.6 in fusion group. In decompression group, ODI decreased to 32.0 at 6 weeks, and to 25.4 at last follow-up. In fusion group, ODI decreased to 45.3 at 6 weeks, and to 25.6 at last follow-up. f/u : follow-up, ODI : Oswestry Disability Index, pre OP : preoperation.




Fig. 5. Sequential changes in mean lumbar lordotic angles before to after surgery. The mean preoperative lumbar lordotic angle was 35.3° in decompression group, and 32.3° in fusion group. In decompression group, the mean lumbar lordotic angle decreased to 31.8° at last follow-up, but without a significance. In fusion group, the mean lumbar lordotic angle increased to 37.8° at 6 weeks, and since there was no significant change. f/u : follow-up, pre OP : preoperation.

1 patient were found. Although there was no statistical significance, perioperative morbidities were more frequent in the fusion group.

In decompression group, 3 patients (9.7%) developed late complications (1 recurrence, 2 instability), and 2 patients (6.5%) underwent reoperation due to spinal instability and recurrence of symptoms with disc herniation. In fusion group, 8 patients (27.6%) developed late complications (1 subsidence, 3 screw loosening, 2 non-union, and 2 ASD) and fusion rate was 93.1%. Also, 3 patients (10.3%) underwent reoperation due to postoperative epidural hematoma, development of ASD, and fusion failure (Table 4). Although late complications were more frequent in the fusion group, it is not reasonable to compare the late complication rate between two groups because the items of complications are different.

DISCUSSION

The indications for decompressive surgery for two-level or more lumbar spinal stenosis are radiating leg pain or NIC resistant to conservative treatment. However, the appropriate surgical technique for treating two-level or more lumbar spinal stenosis remains controversial because the different surgical techniques (i.e., decompression alone or fusion) have their own unique advantages and disadvantages.

Spinal fusion has been performed on patients with severe spinal stenosis and instability to reduce the movement of the injured segment or to increase segment weight bearing ability^{2,4,10,13,16}. Spinal fusion is known to relieve motion-induced discogenic pain and is biomechanically more stable than other procedures.

However, fusion also has its problems, such as, hardware failure, non-union, donor site pain, a protracted operation time, and invasiveness causing injury to paravertebral soft tissues which results in high perioperative morbidity¹⁵. In addition, fusion causes restricts motion by eliminating the functional mobilities of involved segments, and may increase biomechanical stresses on adjacent segments and lead to ASD^{14,21}. Furthermore, two-level or more fusion is likely to cause more ASD than single-level fusion, therefore, selection of fusion should be carefully considered in cases of two-level or more lumbar spinal stenosis⁶.

To prevent these problems associated with fusion, decompressive laminectomy alone has been adopted by many surgeons. Decompressive laminectomy alone can minimize tissue injuries, shorten operation times, reduce perioperative morbidity, and prevent instrument-related complications. Some surgeons have reported that decompressive surgery alone in single-level spinal stenosis produces good results^{5,12,17,22}, and other have reported that old age does not increase morbidity associated with decompressive surgery alone in lumbar spinal stenosis^{9,11,19}.

However, the stability, safety, and efficacy of decompression without fusion are somewhat controversial. Two-level or more decompression alone has a risk of lumbar instability after surgery. In a meta-analysis, some authors found that 69% of patients treated with decompression alone for lumbar spinal stenosis experienced a favorable outcome, as compared with 90%

of those treated with fusion¹.

The present study demonstrates that the postoperative total lumbar lordotic angles increased significantly in the fusion group, which indicates in terms of the correction of lumbar lordosis, fusion is better than decompression alone. However, in our decompression group, decreases in lumbar lordotic angles were not significant, and aggravations of instability were uncommon (only 2 of 31 patients). These results suggest two-level or more laminectomy does not normally cause instability, and that preoperative mild instability is not normally aggravated after surgery. In old age groups, these findings are probably due to the spinal stability conferred by age-related changes of discs and facet joints.

In fact, many authors have reported that the corrections of lumbar lordosis have not been well correlated with clinical outcomes^{15,17,24}. In this study, irrespective of radiological outcome, clinical outcomes, as determined using VAS and ODI scores

Table 4. Summary of clinical outcome according to Odom's criteria

Variable	Decompression group (n=31)	Fusion group (n=29)	p-value
Odom's criteria (number of patients)	-	-	0.407
Excellent	7 (22.5%)	5 (17.3%)	
Good	19 (61.3%)	18 (62.1%)	
Fair	3 (9.7%)	3 (10.3%)	
Poor	2 (6.5%)	3 (10.3%)	

Table 5. Comparison of both surgical methods with respect to perioperative morbidities and late complications

Variable	Decompression group (n=31)	Fusion group (n=29)	p-value
Estimated blood loss (mL)	100±20	560±210	0.001
Operation time (hours)	2.3±0.6	5.2±1.8	0.002
Length of hospital day (days)	7.1±1.0	11.4±5.8	0.008
Perioperative morbidity (number of patients)	3 (9.7%)	7 (24.1%)	0.133
CSF leakage	1	2	
Wound infection	1	2	
Epidural hematoma		1	
Facet fracture	1		
Pneumonia		1	
Urinary difficulty		1	
Late complications (number of patients)	3 (9.7%)	8 (27.6%)	0.073
Recurrence	1		
Instability	2		
Subsidence		1	
Screw loosening		3	
ASD		2	
Non-union		2	
Reoperation	2 (6.5%)	3 (10.3%)	0.586

ASD : adjacent segment degeneration, CSF : cerebrospinal fluid

and Odom's criteria, were not significantly different between our two groups, except low back pain VAS scores at 6 weeks after surgery (i.e., in the short-term). This slow improvement of low back pain in fusion group was probably due to the greater retraction and more severe injury of soft tissues.

In the viewpoint of comparison of each surgical method, EBL, operation time, and hospital stay of decompression group was obviously better than that of fusion group. As a result, although no significant difference in preoperative general conditions was found between the two groups, perioperative morbidities were more frequent in the fusion group. We attribute these differences to more blood loss, longer anesthesia time, more transfusion, and more fluid infusion in fusion group.

The selection of decompression alone or fusion depends on numerous factors, such as, symptoms, age, general condition, the presence of osteoporosis, the number of segments involved, the presence of instability, and surgeon's preference. The most important factor, during the selection process for two-level or more lumbar spinal stenosis, is radiographic instability of the lumbar spine, because fusion is the treatment of choice in overt instability. However, in mild or equivocal instability, the selection of fusion is sometimes worrisome. For example, in elderly patients or those with a poor general condition who is vulnerable to major surgery, protracted surgery, heavy bleeding, and extensive soft tissue injury can cause severe perioperative morbidities, such as, cardiopulmonary complications, wound infections, or even death. Also, in patients with osteoporosis, transpedicular screw fixation or interbody fusion can cause instrument-related complications, such as, subsidence, screw failure, or non-union. Accordingly, decompression alone can be the better choice for patients with two-level or more lumbar spinal stenosis with an advanced age, a poor preoperative condition, or osteoporosis, if there is no severe instability.

Our study has some limitations however. First, since it limited by its retrospective nature, surgical indications in the two study groups were not precisely the same. In particular, patients in fusion group had worse preoperative symptom including low back pain and lumbar lordotic angle, and better preoperative general condition. Second, surgical techniques were not separated in each group (i.e., unilateral or bilateral laminectomy in decompression group, PLIF or TLIF, and percutaneous or open screw fixation in fusion group). Surgical techniques were applied in varying depending on the type of stenosis, and these differences could affect the results. Third, three spinal surgeons were involved in this study, and difference between surgical techniques could have influence outcomes. Finally, the lack of long-term clinical follow-up is a concern, because the risk of symptoms recurrence is generally assessed over 5 years after surgery²⁴).

CONCLUSION

Although the correction of lumbar lordotic angle was better in fusion group, clinical outcomes were not significantly different

in the two groups. Moreover, perioperative morbidities and late complications were better in the decompression group. We suggest that decompressive laminectomy alone to be considered rather than fusion, if there is no overt instability, for patients with two-level or more lumbar spinal stenosis, and especially for elderly patients with a poor general condition or osteoporosis.

References

1. Aiki H, Ohwada O, Kobayashi H, Hayakawa M, Kawaguchi S, Takebayashi T, et al. : Adjacent segment stenosis after lumbar fusion requiring second operation. *J Orthop Sci* 10 : 490-495, 2005
2. Aota Y, Kumano K, Hirabayashi S : Postfusion instability at the adjacent segments after rigid pedicle screw fixation for degenerative lumbar spinal disorders. *J Spinal Disord* 8 : 464-473, 1995
3. Arnoldi CC, Brodsky AE, Cauchoix J, Crock HV, Domisse GF, Edgar MA, et al. : Lumbar spinal stenosis and nerve root entrapment syndromes. Definition and classification. *Clin Orthop Relat Res* : 4-5, 1976
4. Bastian L, Lange U, Knop C, Tusch G, Blauth M : Evaluation of the mobility of adjacent segments after posterior thoracolumbar fixation : a biomechanical study. *Eur Spine J* 10 : 295-300, 2001
5. Fredman B, Arinon Z, Zohar E, Shabat S, Jedeikin R, Fidelman ZG, et al. : Observations on the safety and efficacy of surgical decompression for lumbar spinal stenosis in geriatric patients. *Eur Spine J* 11 : 571-574, 2002
6. Ghiselli G, Wang JC, Bhatia NN, Hsu WK, Dawson EG : Adjacent segment degeneration in the lumbar spine. *J Bone Joint Surg Am* 86A : 1497-1503, 2004
7. Gunzburg R, Szpalski M : The conservative surgical treatment of lumbar spinal stenosis in the elderly. *Eur Spine J* 12 Suppl 2 : S176-S180, 2003
8. Haynes SR, Lawler PG : An assessment of the consistency of ASA physical status classification allocation. *Anaesthesia* 50 : 195-199, 1995
9. Jang IT, Lee SW, Atienza PM, You JS : Decompressive surgery alone for lumbar spinal stenosis in elderly patients. *Korean J Spine* 5 : 83-88, 2008
10. Kanayama M, Hashimoto T, Shigenobu K, Harada M, Oha F, Ohkoshi Y, et al. : Adjacent-segment morbidity after Graf ligamentoplasty compared with posterolateral lumbar fusion. *J Neurosurg* 95 : 5-10, 2001
11. Kim DW, Kim SB, Kim YS, Ko Y, Oh SH, Oh SJ : Surgical treatment of lumbar spinal stenosis in geriatric population : is it risky? *J Korean Neurosurg Soc* 38 : 107-110, 2005
12. Kim SW, Lee SM, Shin H : Surgical outcomes after simple ligamentectomy without fusion for lumbar spinal stenosis. *Korean J Spine* 2 : 350-357, 2005
13. Kumar MN, Jacquot F, Hall H : Long-term follow-up of functional outcomes and radiographic changes at adjacent levels following lumbar spine fusion for degenerative disc disease. *Eur Spine J* 10 : 309-313, 2001
14. Lee CK : Accelerated degeneration of the segment adjacent to a lumbar fusion. *Spine (Phila Pa 1976)* 13 : 375-377, 1988
15. Lehmann TR, Spratt KE, Tozzi JE, Weinstein JN, Reinartz SJ, el-Khoury GY, et al. : Long-term follow-up of lower lumbar fusion patients. *Spine (Phila Pa 1976)* 12 : 97-104, 1987
16. Loupasis GA, Stamos K, Katonis PG, Sapkas G, Korres DS, Hartofilakidis G : Seven- to 20-year outcome of lumbar discectomy. *Spine (Phila Pa 1976)* 24 : 2313-2317, 1999
17. Paik JY, Yoon SM, Yoo CJ : A comparison of clinical outcomes between decompressive laminectomy alone and with arthrodesis in lumbar single level spinal stenosis. *Korean J Spine* 7 : 17-23, 2010
18. Palmer S, Turner R, Palmer R : Bilateral decompressive surgery in lumbar spinal stenosis associated with spondylolisthesis : unilateral approach and use of a microscope and tubular retractor system. *Neuro-*

- surg Focus 13** : E4, 2002
19. Ragab AA, Fye MA, Bohlman HH : Surgery of the lumbar spine for spinal stenosis in 118 patients 70 years of age or older. **Spine (Phila Pa 1976)** **28** : 348-353, 2003
 20. Rosen DS, O'Toole JE, Eichholz KM, Hrubes M, Huo D, Sandhu FA, et al. : Minimally invasive lumbar spinal decompression in the elderly : outcomes of 50 patients aged 75 years and older. **Neurosurgery** **60** : 503-509; discussion 509-510, 2007
 21. Schlegel JD, Smith JA, Schleusener RL : Lumbar motion segment pathology adjacent to thoracolumbar, lumbar, and lumbosacral fusions. **Spine (Phila Pa 1976)** **21** : 970-981, 1996
 22. Son BG, Choi ES, Jung ES, Shin JH, Kim MJ, Chi YC : Clinical comparison between decompression and lumbar interbody fusion with instrumentation for lumbar lateral zone stenosis. **Korean J Spine** **1** : 88-93, 2004
 23. White AA III, Panjabi MM : Physical properties and functional mechanics of the spine in White AA III, Panjabi MM (eds) : **Clinical biomechanics of the spine**. Philadelphia : JB Lippincott, 1978, pp1-60
 24. Yamashita K, Ohzono K, Hiroshima K : Five-year outcomes of surgical treatment for degenerative lumbar spinal stenosis : a prospective observational study of symptom severity at standard intervals after surgery. **Spine (Phila Pa 1976)** **31** : 1484-1490, 2006