

Development of Loop-Mediated Isothermal Amplification Targeting 18S Ribosomal DNA for Rapid Detection of *Azumiobodo hoyamushi* (Kinetoplastea)

Su-Min Song^{1,†}, Dinzouna-Boutamba Sylvatrie-Danne^{1,†}, So-Young Joo¹, Yun Kyung Shin², Hak Sun Yu³, Yong-Seok Lee⁴, Ji-Eon Jung⁴, Noboru Inoue⁵, Won Kee Lee⁶, Youn-Kyoung Goo¹, Dong-Il Chung¹, Yeonchul Hong^{1,*}

¹Department of Parasitology and Tropical Medicine, Kyungpook National University School of Medicine, Daegu 700-422, Korea;

²Aquaculture Management Division, National Fisheries Research & Development Institute, Busan 619-705, Korea; ³Department of Parasitology, School of Medicine, Pusan National University, Yangsan 626-870, Korea; ⁴Department of Life Science and Biotechnology, College of Natural Sciences, Soonchunhyang University, Asan 336-745, Korea; ⁵National Research Center for Protozoan Diseases, Obihiro University of Agriculture and Veterinary Medicine, Obihiro 080-8555, Japan; ⁶Department of Preventive Medicine, Kyungpook National University School of Medicine, Daegu 700-422, Korea

Abstract: Ascidian soft tunic syndrome (AsSTS) caused by *Azumiobodo hoyamushi* (*A. hoyamushi*) is a serious aquaculture problem that results in mass mortality of ascidians. Accordingly, the early and accurate detection of *A. hoyamushi* would contribute substantially to disease management and prevention of transmission. Recently, the loop-mediated isothermal amplification (LAMP) method was adopted for clinical diagnosis of a range of infectious diseases. Here, the authors describe a rapid and efficient LAMP-based method targeting the 18S rDNA gene for detection of *A. hoyamushi* using ascidian DNA for the diagnosis of AsSTS. *A. hoyamushi* LAMP assay amplified the DNA of 0.01 parasites per reaction and detected *A. hoyamushi* in 10 ng of ascidian DNA. To validate *A. hoyamushi* 18S rDNA LAMP assays, AsSTS-suspected and non-diseased ascidians were examined by microscopy, PCR, and by using the LAMP assay. When PCR was used as a gold standard, the LAMP assay showed good agreement in terms of sensitivity, positive predictive value (PPV), and negative predictive value (NPV). In the present study, a LAMP assay based on directly heat-treated samples was found to be as efficient as DNA extraction using a commercial kit for detecting *A. hoyamushi*. Taken together, this study shows the devised *A. hoyamushi* LAMP assay could be used to diagnose AsSTS in a straightforward, sensitive, and specific manner, that it could be used for forecasting, surveillance, and quarantine of AsSTS.

Key words: *Azumiobodo hoyamushi*, LAMP, diagnostic method, soft tunic syndrome, ascidian aquaculture

As its name suggests, soft tunic syndrome (AsSTS) affects the tunics of affected ascidians, such as *Halocynthia roretzi* (Dra-sche), which become thin and soft and are easily torn or ruptured and may result in mass mortality [1]. The repeated occurrence of AsSTS presents serious problems to those involved in the aquaculture of ascidians with respect to the economic burden imposed by epizootic outbreaks and its incidence is gradually increasing in-line with massive increases in aquaculture and devastation of the marine environment. However, early diagnosis and management of AsSTS are difficult, and

thus a rapid, sensitive, and easily performed diagnostic method that produces readily interpretable results is required to forecast outbreaks and to devise effective control programs [2].

Azumiobodo hoyamushi, a flagellated parasite belonging to the order Neobodonida, was recently identified as a causative agent of AsSTS. The diagnosis of AsSTS is established by histopathologic examination of the ascidian tunic by microscopy, but it has limited sensitivity in the case of low parasite density in early infection [2]. Recently, a highly sensitive PCR method based on the 18S rRNA of *A. hoyamushi* was developed for the diagnosis of AsSTS [3], but it requires relatively expensive laboratory equipments, a trained specialist, and has a long turnaround time, which shows some difficulties of direct application of the massive surveillance in the field.

Recently, the loop-mediated isothermal amplification (LAMP) method, which is a relatively straightforward and sen-

• Received 28 February 2014, revised 3 April 2014, accepted 4 April 2014.

† These authors contributed equally to this work.

* Corresponding author (ychong@knu.ac.kr)

© 2014, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

sitive technique based on rapid DNA amplification, was developed and adopted for the clinical diagnosis of a range of infectious diseases [4]. In the LAMP assay, the target DNA is specifically amplified by *Bst* DNA polymerase, which performs strand displacement DNA-synthesis under isothermal conditions and uses a set of 6 oligonucleotides that recognize independent regions of the target gene. The use of 6 oligonucleotides improves the specificity and speed of amplification and forms a loop-structured amplicon, which produces a typical ladder-pattern of multiple bands [5]. A positive reaction is easily determined by eye as turbidity [6] or by adding fluorescent dyes [7,8]. Until now, LAMP assays have been applied to detections of pathogens, such as, viruses, bacteria, fungi, and parasites [5]. Accordingly, the present study was undertaken to develop a simple, rapid, and sensitive diagnostic assay for AsSTS based on LAMP that can be used in the field.

First, we designed a set of 6 LAMP primers targeting the kinetoplast gene for 18S ribosomal RNA (GenBank accession no. AB636162) [9] using the Primer Explorer program (<http://primerexplorer.jp/elamp4.0.0/index.html>; Eiken Chemical Co., Japan) [4] (Table 1). *A. hoyamushi* LAMP was performed for 90 min at 64°C in a 25 µl mixture containing 40 pmol each of FIP and BIP, 5 pmol each of F3 and B3, 20 pmol each of LF and LB, 1.4 mM of deoxynucleoside triphosphates, 0.8 M betaine, and 1 µl of *Bst* DNA polymerase (NEB, Beverly, Massachusetts, USA) in 2.5 µl of buffer [20 mM Tris-HCl pH 8.8, 10 mM KCl, 10 mM (NH₄)₂SO₄, 8 mM MgSO₄, and 0.1% Tween 20], and 1 µl of template DNA in a Loopamp real-time turbidimeter (Realoop-30; Eiken Chemical Co., Tokyo, Japan). Reactions were inactivated by heating at 80°C for 2 min. Optimal temperature and time for LAMP reaction were determined using cloned 18S rDNA (10⁶ copies per reaction) under isother-

mal conditions at temperatures of 60-65°C for 120 min, by monitoring turbidity. Although amplification targeting the 18S rDNA gene was detected at all temperatures tested, the LAMP assay reached a threshold value (0.1) quickest at 64°C (data not shown). No non-specific amplification was detected in the negative control (plasmid containing no insert) after at least 120 min of incubation. Thus, subsequent LAMP reactions were conducted at 64°C for 90 min.

The sensitivity of the LAMP amplification was assessed in 2 ways. First, the LAMP assay was carried out using serially diluted *A. hoyamushi* genomic DNA extracted from *in vitro* cultured flagellates. The LAMP procedure amplified the targeted region at each dilution from the highest DNA concentration examined (1 ng per reaction) to as little as 1 fg per reaction, which reached the threshold value of absorbance (0.1) within 40 min (Fig. 1A). As shown in Fig. 1B, a typical ladder pattern of amplified LAMP products was observed by agarose gel (2.0%) electrophoresis in lanes containing 1 ng to 100 fg per reaction (Fig. 1B, lanes 1-5). Amplified products of positive reactions were also visually detectable using the Loopamp fluorescent detection reagent (FD; Eiken Chemical Co.) under UV light (Fig. 1C). Alternatively, DNA was extracted from ca 1 × 10⁶ parasites, serially diluted to the equivalent of 1,000 to 0.01 parasites per reaction, and used as the template for the PCR and LAMP assays. F3 and B3 primers (Table 1) were used for PCR and generated a product 205 bp long. The PCR conditions of 18S rDNA used were as follows: an initiation step of 94°C for 10 min followed by 35 amplification cycles (94°C for 30 sec, 64°C for 30 sec, and 72°C for 60 sec) and a final extension step at 72°C for 5 min. The LAMP procedure amplified the DNA of parasites diluted to a concentration equivalent to 0.01 parasites per reaction (Fig. 1D). In contrast, PCR amplified the DNA of parasites diluted to 0.1 parasite per reaction (Fig. 1E). Thus, the detection limit of the LAMP assay was 10 times higher than that of PCR. To determine the specificity of the LAMP assay amplification, amplified LAMP products were digested with *Mbo*I, of which cleavage site was represented in the target region of the 18S rDNA gene, and that generated the expected 144 bp fragment (Fig. 1B, lane 9). No amplification was detected in LAMP assays conducted on the genomic DNAs of *Pseudoalteromonas* sp. or *Haslibacter halocynthiae* found in *H. roretzi* [10,11] or in those conducted on the 2 pathogenic *H. roretzi* unrelated flagellates (*Trichomonas vaginalis* and *Trypanosoma brucei*) (data not shown).

Then, template DNAs from non-diseased and AsSTS-posi-

Table 1. Names and sequences of the primers used for *A. hoyamushi* 18S rDNA LAMP reaction

Primer	Sequence (5'-3')
F3	GAATGGTGGTGCATGGCC
B3	GCCCCAAATCTCACCTCGTT
FIP (F1c-F2)	CTACTGGGCGGCTTGGATCTCGCTTTTGGTCGGTG-GAGT
BIP (B1c-B2)	AGCAATCCTCTTGCTCGGCTTAGGAATCCCGCAGA-GAAGG
LF	GTTGACGGAATCAACCAAACAATC
LB	CGGCTGACTGAGGCAACCT

F3, forward outer primer; B3, backward outer primer; FIP, forward inner primer; BIP, backward inner primer; LF, loop forward primer; and LB, loop backward primer.




Fig. 1. Detection limit of *Azumiobodo hoyamushi* 18S rDNA LAMP assays (A). LAMP assays were performed using serial dilutions of *A. hoyamushi* genomic DNA (1 ng to 1 fg per reaction). Distilled water was used as a negative control. LAMP products were visualized by gel electrophoresis (B) and using Loopamp® fluorescent detection reagent (FD) (C). (B, C) Lane M, 100-bp DNA marker; lane 1, 1 ng; lane 2, 100 pg; lane 3, 10 pg; lane 4, 1 pg; lane 5, 100 fg; lane 6, 10 fg; lane 7, 1 fg of *A. hoyamushi* genomic DNA; lane 8, distilled water; and lane 9, LAMP product after *Mbol* digestion. (D-E) *A. hoyamushi* at a density of 1×10^3 parasites/ μ l was serially diluted and tested (D) using the LAMP assay (D) and by PCR (E) using F3 and B3 primers. Lane M, 100-bp DNA marker; lane 1, 1,000; lane 2, 100; lane 3, 10; lane 4, 1; lane 5, 0.1; lane 6, 0.01 of parasites per reaction; lane 7, distilled water. *A. hoyamushi* genomic DNA was prepared using DNeasy tissue kits (Qiagen) from in vitro cultured *A. hoyamushi* species [9] which were kindly provided by Dr. Kyung Il Park (Kunsan National University, Gunsan, Korea).

tive ascidians (determined the presence of *A. hoyamushi* flagellates by microscopic examinations) were tested for the detection of *A. hoyamushi* DNA using the PCR and LAMP assays. Ascidians with or without AsSTS were collected from aquaculture sites in Tongyeong-gun (County) on the southeastern coast of the Republic of Korea in March and April 2013. Animals were classified as Grade 0 to Grade 4, as previously described (Grade 0=healthy ascidian in a non-infected population; Grade 1=apparently non-diseased ascidian in a diseased population; Grade 2=slightly soft tunicate; Grade 3=considerably soft tunicate; Grade 4=severe soft tunicate) [12]. Grade 0

specimens were not available in Korea and thus Grade 1 specimens (non-diseased) were used as a control following the previous criteria. The collected ascidians were washed several times with 0.2 μ m-filtered artificial seawater, tunics were dissected adjacent to the 2 siphons, chopped into small pieces ($\sim 0.2 \times 0.2$ cm) and placed in tubes containing 20 ml of sterilized artificial seawater, incubated at room temperature overnight, and directly examined under microscope the presence of flagellates as previously described [13]. To prepare ascidians genomic DNA for PCR and LAMP, the dissected ascidians prepared above were incubated with 10 ml of PBS at room tem-




Fig. 2. Results of the LAMP assay for detection of *Azumiobodo hoyamushi* from AsSTS (Grades 2, 3) and non-diseased (Grade 1) ascidians. (A) The amplified products of LAMP reactions were examined by real-time turbidity: *A. hoyamushi* genomic DNA (1 pg), non-diseased ascidian DNA (100 ng), AsSTS ascidian DNA (Grade 3, 100 ng; Grade 2, 100 ng; [please confirm here] Grade 2, 10 ng), and distilled water. (B-C) Detection of *A. hoyamushi* in non-diseased (Grade 1) and in AsSTS infected ascidians (Grades 2, 3) using DNAs (B) prepared by a commercial kit (DNeasy tissue kit, Qiagen) or heat-treated lysates (C).

Table 2. Sensitivity, specificity, PPV, NPV, and agreements of microscopy and LAMP versus PCR (the gold standard)

Assay	Microscopy	PCR	LAMP
AsSTS-suspected ascidians (n=36)	Positive: 27 Negative: 9	Positive: 27 Positive: 5 Negative: 4	Positive: 27 Positive: 5 Positive: 1 Negative: 3
Non-diseased ascidians (n=50)	Positive: 0 Negative: 50	Positive: 1 Negative: 49	Positive: 3 Negative: 47
Sensitivity (%) (95% CI)	79.4 (61.6-90.7)		100 (87.4-100)
Specificity (%) (95% CI)	100 (91.6-100)		94.3 (83.4-98.5)
PPV (%) (95% CI)	100 (84.5-100)		91.9 (77.0-97.9)
NPV (%) (95% CI)	88.3 (76.8-94.8)		100 (91.1-100)
Kappa ^a (95% CI)	0.825 (0.702-0.947)		0.926 (0.850-1.000)

CI, confidence interval; PPV, positive predictive value; NPV, negative predictive value.

^aκ values range from a minimum of 0 (complete disagreement) to 1 (perfect agreement), and κ values of 0.9 to 1.0 are generally accepted to mean excellent agreement.

perature for 30 min. After brief centrifugation, 1 ml of each supernatant was transferred to a new tube, centrifuged, and total DNA was extracted using a DNeasy tissue kit (Qiagen, Valencia, California, USA). One microliter of extracted DNA dissolved in 20 µl of doubly distilled water was used as the template for the PCR and LAMP assays.

Results of the amplification of *A. hoyamushi* DNA by PCR or LAMP were positive only for AsSTS-positive ascidians (Grade 2 to Grade 3) by microscopy. No amplification was detected in non-diseased (Grade 1) ascidians. PCR using F3 and B3 primers amplified the DNA from 100 ng of template DNA of AsSTS-positive ascidians (Grade 2 to Grade 3) (data not shown), whereas the *A. hoyamushi* LAMP assay amplified DNA from 10 ng of template DNA (Grade 2) within 40 min from reaction initiation (Fig. 2A). Thus, the detection limit of the LAMP assay using the DNA extracted from ascidians was 10 times higher than that of PCR, which was consistent with the results above using the pure DNA from in vitro cultured flagellates (Fig. 1D, E).

Because the straightforward sample preparation for molecular diagnosis reduces the risk of cross-contamination and the turnaround time for diagnosis, we prepared the crude cell lysates of ascidians by direct heat-treatment. Briefly, 1 ml of the PBS supernatants above were collected, heated at 100°C for 5 min to disrupt *A. hoyamushi* flagellates, and centrifuged at 12,000 g for 2 min at room temperature. One microliter aliquots of supernatants were collected and used as templates for PCR and LAMP. LAMP assays conducted using heat-treated ly-

sates (Fig. 2C) produced the same results as those obtained using DNA extracted using a DNeasy tissue kit (Fig. 2B), although the incubation time required for LAMP assay to reach a threshold absorbance value of 0.1 was slightly delayed (Fig. 2C).

Next, AsSTS-suspected ascidians (n=36) (Grades 2 or 3) were then subjected to microscopic analysis, PCR, and to the LAMP assay (Table 2). Of the 36 AsSTS-suspected ascidians, positive results by microscopy, PCR, and LAMP were 27 (75%), 32 (89%), and 33 (92%), respectively. All the 27 positive samples by microscopy were positive by PCR and LAMP. Of the 9 negatives by microscopy, 5 samples (56%) were positive by PCR and LAMP and thus considered false negatives. We attribute these 5 false negatives by microscopy to have been caused by a low parasite density in early infection. Of the 4 negatives by microscopy and PCR, 1 was positive by LAMP. Of the 50 non-diseased ascidians (Grade 1) subjected to the same 3 tests, all were negative by microscopy, and of these 50 negatives by microscopy, 1 was positive by PCR and LAMP and 2 were only positive by LAMP.

To determine the accuracy of the 3 tests for diagnosis of AsSTS, PCR was used as a gold standard and sensitivity, specificity, positive predictive value (PPV), and negative predictive value (NPV) were calculated (Table 2). The sensitivity and specificity of microscopic analysis were 79.4% with 95% confidence interval (CI) from 61.6 to 90.7% and 100% (95% CI, 91.6-100%), respectively, whereas those of the LAMP assay were 100% (95% CI, 87.4-100%) and 94.3% (95% CI, 83.4-98.5%), respectively. These results indicate that PCR and LAMP had higher sensitivity compared to that of optical microscopy and thus can be used in the diagnosis of early infections of AsSTS. When PCR was used as the gold standard, LAMP showed excellent PPV (91.9%, 95% CI: 77.0-97.9%) and NPV (100%, 95% CI: 91.1-100%) and better agreement with PCR (Kappa=0.929) (95% CI, 0.850-1.000) ($P<0.001$) than microscopy (Kappa=0.825) (95% CI, 0.702-0.947) ($P<0.001$).

Since AsSTS was first reported, efforts have been made to identify the causative agent, but recently, *A. hoyamushi* was found to be responsible [9,13]. In agreement with previous studies, the LAMP assay detected *A. hoyamushi* only in ascidians with AsSTS (Fig. 2A). Furthermore, the *A. hoyamushi* LAMP assay showed no cross-reactivity with the ascidian tunic, with other organisms previously suggested to be associated with AsSTS, or with AsSTS unrelated flagellates (data not shown). In addition, the devised LAMP assay showed good agreement

with PCR and high PPV and NPV values.

Previous authors have reported that the clinical signs of AsSTS are evident at siphons during the early stage of infection and subsequently extend into adjacent regions [2]. In the present study, to enhance the sensitivity of diagnostic methods, tunics adjacent to the siphons were chopped, incubated with PBS for only 30 min, heat-treated, and directly used as the template for the LAMP assay instead of using the extracted DNA by conventional methods from the overnight enrichment of the sliced tunics with seawater [13]. The LAMP assay conducted on heat-treated samples was found to be as efficient at detecting *A. hoyamushi* as DNA extracted using a commercial kit probably due to the ability of LAMP to better tolerate biological contaminants than PCR [14,15]. In this study, the LAMP assay conducted using heat-treated lysates was completed within 50 min (Fig. 2C), which is substantially faster than PCR using DNA prepared by conventional methods or commercially available kits. Thus, the devised *A. hoyamushi* LAMP assay could eliminate the need for DNA extraction without compromising sensitivity and reduce the time required to reach a diagnosis. Furthermore, it could be applied for AsSTS surveillance and quarantine for importation and exportation purposes. In addition, simplicity, rapid sample preparation, and straightforward detection suggest that the devised assay has potential forecasting applications in seawater regions where AsSTS is prevalent [16,17].

ACKNOWLEDGMENT

This study was supported by the Aquaculture Management Division, National Fisheries Research & Development Institute (Grant no. RP-2012-AQ-32).

CONFLICT OF INTEREST

We have no conflict of interest related to this study.

REFERENCES

1. Hirose E, Ohtake SI, Azumi K. Morphological characterization of the tunic in the edible ascidian, *Halocynthia roretzi* (Drasche), with remarks on 'soft tunic syndrome' in aquaculture. J Fish Dis 2009; 32: 433-445.
2. Kumagai A, Suto A, Ito H, Tanabe T, Takahashi K, Kamaishi T, Miwa S. Mass mortality of cultured ascidians *Halocynthia roretzi* associated with softening of the tunic and flagellate-like cells. Dis Aquat Organisms 2010; 90: 223-234.

3. Kumagai A, Kamaishi T. Development of polymerase chain reaction assays for detection of the kinetoplastid *Azumiobodo hoyamushi*, the causative agent for soft tunic syndrome in the ascidian *Halocynthia roretzi*. *Fish Pathol* 2013; 48: 42-47.
4. Notomi T, Okayama H, Masubuchi H, Yonekawa T, Watanabe K, Amino N, Hase T. Loop-mediated isothermal amplification of DNA. *Nucl Acids Res* 2000; 28: E63.
5. Mori Y, Notomi T. Loop-mediated isothermal amplification (LAMP): a rapid, accurate, and cost-effective diagnostic method for infectious diseases. *J Infect Chemother* 2009; 15: 62-69.
6. Mori Y, Nagamine K, Tomita N, Notomi T. Detection of loop-mediated isothermal amplification reaction by turbidity derived from magnesium pyrophosphate formation. *Biochem Biophys Res Commun* 2001; 289: 150-154.
7. Goto M, Honda E, Ogura A, Nomoto A, Hanaki K. Colorimetric detection of loop-mediated isothermal amplification reaction by using hydroxy naphthol blue. *Biotechniques* 2009; 46: 167-172.
8. Tomita N, Mori Y, Kanda H, Notomi T. Loop-mediated isothermal amplification (LAMP) of gene sequences and simple visual detection of products. *Nat Protoc* 2008; 3: 877-882.
9. Hirose E, Nozawa A, Kumagai A, Kitamura S. *Azumiobodo hoyamushi* gen. nov. et sp. nov. (Euglenozoa, Kinetoplastea, Neobodonida): a pathogenic kinetoplastid causing the soft tunic syndrome in ascidian aquaculture. *Dis Aquat Organisms* 2012; 97: 227-235.
10. Kim D, Baik KS, Park SC, Kim SJ, Shin TS, Jung SJ, Oh MJ, Seong CN. Cellulase production from *Pseudoalteromonas* sp. NO3 isolated from the sea squirt *Halocynthia roretzi*. *J Indust Microbiol Biotechnol* 2009; 36: 1375-1382.
11. Kim SH, Yang HO, Shin YK, Kwon HC. *Hasllibacter halocynthiae* gen. nov., sp. nov., a nutriacholic acid-producing bacterium isolated from the marine ascidian *Halocynthia roretzi*. *Int J Syst Evol Microbiol* 2012; 62: 624-631.
12. Kitamura SI, Ohtake SI, Song JY, Jung SJ, Oh MJ, Choi BD, Azumi K, Hirose E. Tunic morphology and viral surveillance in diseased Korean ascidians: soft tunic syndrome in the edible ascidian, *Halocynthia roretzi* (Drasche), in aquaculture. *J Fish Dis* 2010; 33: 153-160.
13. Kumagai A, Ito H, Sasaki R. Detection of the kinetoplastid *Azumiobodo hoyamushi*, the causative agent of soft tunic syndrome, in wild ascidians *Halocynthia roretzi*. *Dis Aquat Organisms* 2013; 106: 267-271.
14. Enomoto Y, Yoshikawa T, Ihira M, Akimoto S, Miyake F, Usui C, Suga S, Suzuki K, Kawana T, Nishiyama Y, Asano Y. Rapid diagnosis of herpes simplex virus infection by a loop-mediated isothermal amplification method. *J Clin Microbiol* 2005; 43: 951-955.
15. Kaneko H, Kawana T, Fukushima E, Suzutani T. Tolerance of loop-mediated isothermal amplification to a culture medium and biological substances. *J Biochem Biophys Methods* 2007; 70: 499-501.
16. Sharma S, Srinivasan M, George C. *Acanthamoeba* keratitis in non-contact lens wearers. *Arch Ophthalmol* 1990; 108: 676-678.
17. Srinivasan M, Burman S, George C, Nirmalan PK. Non-contact lens related *Acanthamoeba* keratitis at a tertiary eye care center in south India: implications for eye care programs in the region. *Med Sci Monit* 2003; 9: CR125-CR129.