

SOME PROOFS OF THE CLASSICAL INTEGRAL HARDY INEQUALITY

IDDRISU MOHAMMED MUNIRU*, OKPOTI CHRISTOPHER ADJEI,
AND GBOLAGADE KAZEEM ALAGBE

ABSTRACT. We present some proofs of the classical integral Hardy inequality. Our approach makes use of continuous functions with compact support in $(0, \infty)$, homogeneity of the norm and Schur's criterion for integral operators.

1. Introduction

The classical integral inequality announced by G. H. Hardy in 1920 is given by

$$(1) \quad \int_0^\infty \left(\frac{1}{x} \int_0^x f(t) dt \right)^p dx \leq \left(\frac{p}{p-1} \right)^p \int_0^\infty f^p(x) dx,$$

where $p > 1$, $x > 0$, f is a nonnegative measurable function on $(0, \infty)$ and the constant $\left(\frac{p}{p-1} \right)^p$ is the best possible [4]. This interesting result (1) was later proved by Hardy himself in 1925 (see [1], [5], [7], [8], [9] and the references therein.) Inequality (1) can also be written as

$$(2) \quad \int_0^\infty F^p(x) dx \leq \left(\frac{p}{p-1} \right)^p \int_0^\infty f^p(x) dx,$$

Received May 5, 2014. Revised July 25, 2014. Accepted July 25, 2014.

2010 Mathematics Subject Classification: 26D10, 26D15.

Key words and phrases: Hardy inequality, Proofs, homogeneity, integral operators.

*Corresponding author.

© The Kangwon-Kyungki Mathematical Society, 2014.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

where $0 < F(x) = \frac{1}{x} \int_0^x f(t)dt < \infty$, $f > 0$.

The inequalities (1) and (2) are very popular in the research environment. See also [6].

Our task in this paper is mainly to deepen understanding of the Hardy inequality (2) by providing elaborate proofs.

2. Preliminary Notes

We define continuous functions and present some auxiliary results.

Definition (Continuous functions) [11]. Let X be a subset of the set of real numbers \mathfrak{R} , and let $f : X \rightarrow \mathfrak{R}$ be a function. Let $x_0 \in X$. We say that f is continuous at x_0 if and only if we have $\lim_{x \rightarrow x_0} f(x) = f(x_0)$ for every $x \in X$. In other words, the limit of $f(x)$ as x converges to x_0 in X exists and is equal to $f(x_0)$.

Support of a function. Let I be a nonempty open set in \mathfrak{R}^n , and let f be a continuous function on I . The support of f , denoted by $\text{supp}(f)$, is defined to be the complement of the largest open set on which f is zero. That is

$$\text{supp}(f) = \overline{\{x \in I : f(x) \neq 0\}},$$

the closure of the set $x \in I$ where $f(x) \neq 0$. (See [3], p. 134).

Fatou's Lemma ([2], p. 52). Let X be a measure space with measure μ . Let $\{f_n\}$ be any sequence of measurable functions on X with range in $[0, \infty]$. For each positive integer n ,

$$(3) \quad \int_X (\liminf_{n \rightarrow \infty} f_n) d\mu \leq \liminf_{n \rightarrow \infty} \int_X f_n d\mu.$$

Hölder's inequality ([2], p.182). Suppose $1 < p < \infty$ and $\frac{1}{p} + \frac{1}{q} = 1$ (that is $p + q = pq$). Let X be a measure space with measure μ . If f and g are measurable functions on X with range in $[0, \infty]$, then

$$(4) \quad \int_X fg d\mu \leq \left\{ \int_X f^p d\mu \right\}^{\frac{1}{p}} \left\{ \int_X g^q d\mu \right\}^{\frac{1}{q}},$$

In particular, if $f \in L^p$ and $g \in L^q$, then $fg \in L^1$, and in this case equality holds in (4) if and only if $\alpha|f|^p = \beta|g|^q$ almost everywhere for some constants α, β with $\alpha\beta \neq 0$. See [2] and also [10] for proofs of inequalities (3) and (4).

PROPOSITION 1. ([2], p.195). For $f \in L^p$ and $g \in L^q$, let

$$Tf(x) = x^{-1} \int_0^x f(y)dy, \quad Sg(y) = \int_y^{+\infty} x^{-1}g(x)dx.$$

Then for $1 < p \leq \infty$ and $1 \leq q < \infty$,

$$\|Tf\|_p \leq \frac{p}{p-1} \|f\|_p, \quad \|Sg\|_q \leq q \|g\|_q.$$

Proof. Let

$$K(x, y) = \begin{cases} \frac{1}{x} & \text{if } 0 < y < x \\ 0 & \text{otherwise} \end{cases}.$$

Then

$$\int_0^{+\infty} |K(1, y)|y^{-\frac{1}{p}}dy = \int_0^1 y^{-\frac{1}{p}}dy = \frac{p}{p-1} = q,$$

where $\frac{1}{p} + \frac{1}{q} = 1$, yielding the result. \square

We now present our main results which are basically the different approaches to the proof of inequality (2). We denote by $C_c(0, +\infty)$, the set of all continuous functions with compact support in $(0, +\infty)$.

3. First Proof

Integration by parts and Hölder's inequality are essentially applied here.

3.1. Case 1: Let $p > 1$, f is positive and continuous with compact support in $(0, +\infty)$ and F is positive and differentiable on $[0, +\infty)$.

Setting $u = F^p$ and $dv = dx$ implies $du = pF^{p-1}F'dx$ and $v = x$.

Consider (a, A_0) with $0 < a < A_0 < \infty$ so that the $\text{supp} f \subset [a, A_0]$. Integration by parts of $\int_a^{A_0} F^p dx$ gives

$$(5) \quad \int_a^{A_0} F^p dx = [xF^p]_a^{A_0} - p \int_a^{A_0} xF^{p-1}F' dx.$$

But $\int_0^A F^p dx = \int_0^a F^p dx + \int_a^{A_0} F^p dx + \int_{A_0}^A F^p dx$ for $0 < A_0 < A < \infty$. Since $f \in C_c(0, +\infty)$ with $\text{supp} f \subset [a, A_0]$ then $F(x) = 0$ for $0 \leq x < a$ and $A_0 < x \leq A$. Thus

$$\int_0^{A_0} F^p(x) dx = \int_0^A F^p(x) dx$$

and (5) becomes

$$(6) \quad \int_0^A F^p(x) dx = AF^p(A) - p \int_0^A xF^{p-1}(x)F'(x) dx.$$

But $F(x) = \frac{1}{x} \int_0^x f(t) dt$ implies $xF(x) = \int_0^x f(t) dt$. Differentiating gives $xF'(x) + F(x) = f(x)$

Thus (6) becomes

$$\begin{aligned} \int_0^A F^p(x) dx &= AF^p(A) - p \int_0^A F^{p-1}(x) \{f(x) - F(x)\} dx \\ &= AF^p(A) - p \int_0^A F^{p-1}(x) f(x) dx + p \int_0^A F^p(x) dx, \end{aligned}$$

$$(7) \quad \int_0^A F^p(x) dx = \frac{AF^p(A)}{1-p} + \frac{p}{p-1} \int_0^A F^{p-1}(x) f(x) dx.$$

By Hölder's inequality,

$$(8) \quad \int_0^A F^{p-1}(x) f(x) dx \leq \left(\int_0^A F^{(p-1)q}(x) dx \right)^{\frac{1}{q}} \left(\int_0^A f^p(x) dx \right)^{\frac{1}{p}}$$

where $\frac{1}{p} + \frac{1}{q} = 1$ or $q = \frac{p}{p-1}$.

Putting (8) into (7), then

$$(9) \quad \int_0^A F^p(x) dx \leq \frac{AF^p(A)}{1-p} + \frac{p}{p-1} \left(\int_0^A F^{(p-1)q}(x) dx \right)^{\frac{1}{q}} \left(\int_0^A f^p(x) dx \right)^{\frac{1}{p}}.$$

Let $I = \int_0^A f(t)dt$ so that $F(A) = \frac{I}{A}$. Then $AF^p(A) = \frac{I^p}{A^{p-1}}$ turns to 0 as $A \rightarrow \infty$. Thus (9) simplifies to

$$(10) \quad \int_0^\infty F^p(x)dx \leq \frac{p}{p-1} \left(\int_0^\infty F^p(x)dx \right)^{(1-\frac{1}{p})} \left(\int_0^\infty f^p(x)dx \right)^{\frac{1}{p}}.$$

Since $(\int_0^\infty F^p(x)dx)^{(1-\frac{1}{p})} > 0$, then

$$\left(\int_0^\infty F^p(x)dx \right)^{\frac{1}{p}} \leq \frac{p}{p-1} \left(\int_0^\infty f^p(x)dx \right)^{\frac{1}{p}},$$

Thus

$$\int_0^\infty F^p(x)dx \leq \left(\frac{p}{p-1} \right)^p \int_0^\infty f^p(x)dx. \quad \square$$

3.2. Case 2: We consider f not necessarily positive. Let $f \in L^p(0, +\infty)$ and set $f_+ = \max(f, 0)$ and $f_- = -\min(0, f)$. Consider sequence $(f_n)_{n \in \mathbb{N}}$ of functions in $C_c(0, +\infty)$ such that $f_n \rightarrow f$ in $L^p(0, +\infty)$. For a fixed $x > 0$, $F(x) = \frac{1}{x} \int_0^x f(t)dt$ is well defined for $f \geq 0$. By Hölder's inequality,

$$\begin{aligned} \left| \frac{1}{x} \int_0^x f(t)dt \right| &\leq \frac{1}{x} \int_0^x |f(t)|dt \\ &\leq \frac{1}{x} \left(\int_0^x |f(t)|^p dt \right)^{\frac{1}{p}} \left(\int_0^x 1^q dt \right)^{\frac{1}{q}} \\ &\leq x^{-\frac{1}{p}} \left(\int_0^x |f(t)|^p dt \right)^{\frac{1}{p}}. \end{aligned}$$

Hence, by Fubini, we get

$$\int_0^\infty |F(x)|^p dx \leq \left(\frac{p}{p-1} \right)^p \int_0^\infty |f(x)|^p dx,$$

where the constant $\left(\frac{p}{p-1} \right)^p$ is the best possible.

Also set $f = f_+ - f_-$, $|f| = f_+ + f_-$ and $F = F_+ - F_-$, $|F| = F_+ + F_-$

and $\int |F| < \infty$. Let $\|f_n - f\| < \frac{1}{n}$. Then

$$\begin{aligned} |F_n(x) - F(x)| &= \left| \frac{1}{x} \int_0^x f_n(t) dt - \frac{1}{x} \int_0^x f(t) dt \right| \\ &\leq \frac{1}{x} \int_0^x |f_n(t) - f(t)| dt \\ &\leq \frac{1}{x} \left(\int_0^x |f_n(t) - f(t)|^p dt \right)^{\frac{1}{p}} \left(\int_0^x 1^q dt \right)^{\frac{1}{q}} \quad \text{by (4)} \\ &\leq x^{-\frac{1}{p}} \|f_n - f\|_p \\ &< \frac{x^{-\frac{1}{p}}}{n} \rightarrow 0 \quad \text{as } n \rightarrow \infty. \end{aligned}$$

This shows pointwise convergence: $|F_n(x)| \rightarrow |F(x)|$ as $x \rightarrow \infty$. Therefore

$$(11) \quad \int_0^\infty |F_n(x)|^p dx \leq \left(\frac{p}{p-1} \right)^p \int_0^\infty |f_n(x)|^p dx.$$

Suppose that f is positive and so is F . Suppose also that f_n is positive which implies F_n is positive. Then

$$\begin{aligned} \int_0^{+\infty} |F(x)|^p dx &= \int_0^{+\infty} \lim_{n \rightarrow +\infty} |F_n(x)|^p dx \\ &\leq \lim_{n \rightarrow +\infty} \int_0^{+\infty} |F_n(x)|^p dx \quad (\text{Fatou's lemma}) \\ &\leq \lim_{n \rightarrow +\infty} \left[\left(\frac{p}{p-1} \right)^p \int_0^{+\infty} |f_n(x)|^p dx \right] \\ &\leq \left(\frac{p}{p-1} \right)^p \int_0^{+\infty} |f(x)|^p dx. \quad \square \end{aligned}$$

4. Second Proof

The approach here makes use of homogeneity of a norm and the use of a kernel.

4.1. Case 1: Homogeneity of a norm.

Let

$$F(x) = \frac{1}{x} \int_0^x f(t) dt = \int_0^1 f(tx) dt.$$

Set $f_t(x) = f(tx)$. By Minkowski inequality for integrals,

$$\begin{aligned}\|F(x)\|_p &\leq \int_0^1 \|f_t(x)\|_p dt \\ &\leq \int_0^1 \left(\int_0^1 |f_t(x)|^p dx \right)^{\frac{1}{p}} dt\end{aligned}$$

By change of variables with $s = tx$, $ds = tdx$, and also by Fubini's theorem we have

$$\begin{aligned}\|F(x)\|_p &\leq \int_0^1 \left(\int_0^1 |f(s)|^p \frac{ds}{t} \right)^{\frac{1}{p}} dt \\ &\leq \int_0^1 t^{-\frac{1}{p}} dt \left(\int_0^1 |f(s)|^p ds \right)^{\frac{1}{p}} \\ \|F(x)\|_p &\leq \frac{p}{p-1} \|f(s)\|_p\end{aligned}$$

Hence

$$\|F(x)\|_p^p \leq \left(\frac{p}{p-1} \right)^p \|f\|_p^p. \quad \square$$

4.2. Case 2: Kernel's approach.

Let $K > 0$ be a Lebesgue measurable function on $(0, \infty) \times (0, \infty)$ and define

$$K(x, y) = \begin{cases} \frac{1}{x} & \text{if } 0 < y < x \\ 0 & \text{otherwise} \end{cases}.$$

Let $f \in L^p$ and consider the integral operator

$$Tf(x) = \int_0^\infty |K(x, y)f(y)| dy.$$

By change of variables with $y = xz$, $dy = xdz$, we have

$$Tf(x) = \int_0^\infty |K(x, xz)|f(xz)x dz.$$

Set $f_z(x) = f(xz)$ and by Minkowski inequality for integrals, we have

$$\begin{aligned} \|Tf(x)\|_p &\leq \int_0^\infty |K(1, z)| \|f_z(x)\|_p dz \\ &\leq \int_0^1 \left(\int_0^\infty |f_z(x)|^p dx \right)^{\frac{1}{p}} dz \end{aligned}$$

Again by change of variables with $xz = y$, $zdx = dy$ and by Fubini's theorem, we have

$$\begin{aligned} \|Tf(x)\|_p &\leq \int_0^1 \left(\int_0^\infty |f(y)|^p \frac{dy}{z} \right)^{\frac{1}{p}} dz \\ &\leq \int_0^1 z^{-\frac{1}{p}} dz \left(\int_0^\infty |f(y)|^p dy \right)^{\frac{1}{p}} \\ &\leq \left(\frac{p}{p-1} \right) \|f(y)\|_p \end{aligned}$$

Hence

$$\|Tf\|_p^p \leq \left(\frac{p}{p-1} \right)^p \|f\|_p^p. \quad \square$$

REMARK 1. Let us remark that the proofs for the cases 1 and 2 here can be described as almost the same, except that the mention of a kernel and its application is demonstrated in case 2 .

5. Third Proof

Let K be a Lebesgue measurable function on $(0, \infty) \times (0, \infty)$. For $1 < p < \infty$, and $\frac{1}{p} + \frac{1}{q} = 1$, define a nonnegative kernel

$$K(x, y) = \begin{cases} \frac{1}{x} & \text{if } 0 < y < x \\ 0 & \text{otherwise} \end{cases} .$$

and consider the integral operator

$$(12) \quad Tf(x) = \int_0^{+\infty} K(x, y)f(y)dy = \frac{1}{x} \int_0^x f(y)dy = F(x)$$

Let $h(t) = t^\alpha$ for $-1 < \alpha < 0$. By Proposition 1, we compute

$$(13) \quad \int_0^{+\infty} K(x, y)h(y)dy = \int_0^x \frac{1}{x}y^\alpha dy = \frac{1}{x} \frac{x^{(\alpha+1)}}{\alpha+1} = \frac{x^\alpha}{\alpha+1}$$

and

$$(14) \quad \int_0^{+\infty} K(x, y)h(x)dx = \int_y^{+\infty} \frac{1}{x}x^\alpha dx = \int_y^{+\infty} x^{(\alpha-1)}dx = -\frac{1}{\alpha}y^\alpha.$$

Introduce y^α into (12) and applying Hölders inequality, we have

$$\begin{aligned} Tf(x) &= \int_0^{+\infty} K(x, y)y^\alpha \frac{f(y)}{y^\alpha} dy \\ &\leq \left(\int_0^{+\infty} K(x, y)y^{q\alpha} dy \right)^{\frac{1}{q}} \left(\int_0^{+\infty} K(x, y) \left(\frac{|f(y)|}{y^\alpha} \right)^p dy \right)^{\frac{1}{p}} \\ &\leq \left(\frac{x^{q\alpha}}{q\alpha+1} \right)^{\frac{1}{q}} \left(\int_0^{+\infty} K(x, y) \left(\frac{|f(y)|}{y^\alpha} \right)^p dy \right)^{\frac{1}{p}} \quad \{\text{by (13)}\} \\ (15) \quad &\leq \left(\frac{1}{q\alpha+1} \right)^{\frac{1}{q}} x^\alpha \left(\int_0^{+\infty} K(x, y) \left(\frac{|f(y)|}{y^\alpha} \right)^p dy \right)^{\frac{1}{p}} \end{aligned}$$

Let $\lambda = \left(\frac{1}{q\alpha+1} \right)^{\frac{1}{q}}$. Thus (15) becomes

$$(16) \quad |Tf(x)|^p \leq \lambda^p x^{p\alpha} \left(\int_0^{+\infty} K(x, y) \left(\frac{|f(y)|}{y^\alpha} \right)^p dy \right)$$

By Fubini's theorem, we get

$$\begin{aligned} \int_0^{+\infty} |Tf(x)|^p dx &\leq \lambda^p \int_0^{+\infty} \left(x^{p\alpha} \left(\int_0^{+\infty} K(x, y) \left(\frac{|f(y)|}{y^\alpha} \right)^p dy \right) \right) dx \\ (17) \quad &\leq \lambda^p \int_0^{+\infty} \left(\int_0^{+\infty} K(x, y)x^{p\alpha} dx \right) \left(\frac{|f(y)|}{y^\alpha} \right)^p dy. \end{aligned}$$

From (14), $Th(y) = \int_0^{+\infty} K(x, y)x^{p\alpha}dx = -\frac{y^{p\alpha}}{p\alpha}$. Thus (17) becomes

$$\begin{aligned} \int_0^{+\infty} |Tf(x)|^p dx &\leq \lambda^p \int_0^{+\infty} -\frac{y^{p\alpha}}{p\alpha} \frac{|f(y)|^p}{y^{p\alpha}} dy \\ &\leq -\frac{\lambda^p}{p\alpha} \int_0^{+\infty} |f(y)|^p dy \\ &\leq C \int_0^{+\infty} |f(y)|^p dy \end{aligned}$$

where $C = -\frac{\lambda^p}{p\alpha}$ for $-1 < \alpha < 0$. Equivalently

$$\|Tf(x)\|_p^p \leq C \|f(y)\|_p^p. \quad \square$$

REMARK 2. The Schur criterion discussed above shows that the operator T is bounded on $L^p(0, +\infty)$ with $\|T\| \leq C$. See ([12], p. 45) for discussions on Schur test.

6. Conclusion

Precise proofs for the classical integral Hardy inequality were presented. A number of useful applications of some important theorems such as Fatou's lemma and Fubini's theorem as well as Hölder and Minkowski inequalities were provided.

Acknowledgement

The first author wish to thank Emeritus Prof. Aline BONAMI and Prof. Frederic SYMESAK for their care and attention during his research period at the University of Angers, France. Also, his deepest and sincere appreciation goes to the French Government for the financial support during the research period.

References

- [1] S. Abramovich, K. Krulic, J. Pecaric and L.-E. Persson, *Some new refined Hardy type inequalities with general kernels and measures*, Aequat. Math. **79** (2010), 157–172.
- [2] G. B. Folland, *Real Analysis: Modern Techniques and Their Applications*, 2nd ed., John Wiley and Sons, Inc., New York, 1999.

- [3] C. Gasquet and P. Witomski, *Fourier Analysis and Applications: Filtering, Numerical Computation, Wavelets*, Springer-Verlag New York, Inc., 1999.
- [4] G. H. Hardy, *Note on a theorem of Hilbert*, Math. Z. **6** (1920), 314–317.
- [5] G. H. Hardy, J. E. Littlewood and G. Polya, *Inequalities*, 2nd ed., Cambridge University Press, Cambridge, 1952.
- [6] S. Kaijser, L. Nikolova, L.- E. Persson and A. Wedestig, *Hardy-Type Inequalities Via Convexity*, J. Math. Ineq. Appl., **8** (3) (2005), 403–417.
- [7] A. Kufner, L. Maligranda and L.- E. Persson, *The Hardy inequality: About its History and Some Related Results*, Vydavatelsky Sevis Publishing House, Pilsen, 2007.
- [8] A. Kufner, L. Maligranda and L.-E. Persson, *The Prehistory of the Hardy Inequality*, Amer. Math. Monthly, **113** (8) (2006), 715–732.
- [9] J. A. Oguntuase and L.- E. Persson, *Refinement of Hardy inequalities via superquadratic and subquadratic functions*, J. Math. Anal. Appl., **339** (2) (2008), 1305–1312.
- [10] W. Rudin, *Real and Complex Analysis*, 2nd edition, McGraw-Hill Inc, 1974.
- [11] T. Tao, *Analysis I*, Hindustan Book Agency, Volume I, 2006.
- [12] K. Zhu, *Spaces of Holomorphic Functions in the Unit Ball*, Springer, New York, 2005.

Iddrisu Mohammed Muniru
Department of Mathematics
University for Development Studies
Navrongo Campus
P. O. Box 24, Navrongo, UE/R, Ghana
E-mail: immuniru@gmail.com

Okpoti Christopher Adjei
Department of Mathematics
University of Education
Winneba, Ghana
E-mail: caokpoti@yahoo.com

Gbolagade Kazeem Alagbe
Department of Computer Science
University for Development Studies
Navrongo Campus
P. O. Box 24, Navrongo, UE/R, Ghana
E-mail: gkazy1@yahoo.com