

Outcomes of nonsurgical periodontal therapy in severe generalized aggressive periodontitis

Amal Bouziane^{1,*}, Latifa Benrachadi², Redouane Abouqal³, Oumkeltoum Ennibi⁴

¹Department of Periodontology, Faculty of Dental Medicine, Biostatistical, Clinical, and Epidemiological Research Laboratory, Faculty of Medicine and Pharmacy, Mohammed V Souissi University, Rabat, Morocco

²Department of Periodontology, Faculty of Dental Medicine, Mohammed V Souissi University, Rabat, Morocco

³Biostatistical, Clinical, and Epidemiological Research Laboratory, Faculty of Medicine and Pharmacy, Mohammed V Souissi University, Rabat, Morocco

⁴Department of Periodontology, Faculty of Dental Medicine, Mohammed V Souissi University, Rabat, Morocco

Purpose: Aggressive periodontitis, especially in its severe form, was traditionally considered to have an unfavourable prognosis. It required a complex treatment and its stabilization was often achieved by surgical therapy. The aim of this study was to investigate the results of nonsurgical periodontal treatment in severe generalized forms of aggressive periodontitis.

Methods: Patients with advanced generalized aggressive periodontitis were included in the study. Probing depth (PD) of pockets ≥ 7 mm and clinical attachment level (CAL) of sites with attachment loss ≥ 5 mm were measured at baseline before nonsurgical periodontal treatment, at re-evaluation, and after treatment. The following other parameters were recorded: resolution of inflammation and bone fill. We compared the baseline values with re-evaluation and posttreatment values using the Friedman test. The Wilcoxon test with the Bonferroni correction was used for both re-evaluation and posttreatment values.

Results: Seven patients with 266 periodontal sites were examined. A significant difference was found between values, reported as medians with interquartile ranges, for PD at baseline (7.94 [7.33–8.19] mm) and both re-evaluation (4.33 [3.63–5.08] mm) and posttreatment (3.54 [3.33–4.11] mm) values ($P=0.002$). A significant difference was also found between values for CAL at baseline (9.02 [7.5–9.2] mm) and both re-evaluation (6.55 [6.30–6.87] mm) and posttreatment (6.45 [5.70–6.61] mm) ($P=0.002$). Inflammation was resolved and angular bone defects were repaired in all cases.

Conclusions: These therapeutic results suggest that this form of periodontitis could have positive outcomes after nonsurgical periodontal treatment. The reparative potential of tissue affected by severe aggressive periodontitis should encourage clinicians to save apparently hopeless teeth in cases of this form of periodontitis.

Keywords: Aggressive periodontitis, Periodontal debridement, Treatment outcome.

INTRODUCTION

Currently, aggressive periodontitis is considered to be a well-defined inflammatory periodontal disease. It usually affects people who, in most cases, otherwise appear healthy. It tends to have a familial aggregation and there is a rapid rate of disease progression. Aggressive periodontitis occurs in localized and generalized forms. Localized aggressive periodontitis is frequently associated with *Aggregatibacter actinomycetemcomitans* while generalized aggressive periodontitis is frequently associated with *A. actinomycetemcomitans* and *Porphyromonas gingivalis* [1].

The specific attributes of the disease process are due to the virulence of the pathogens and the host susceptibility of patients. Since it is difficult to modify underlying immune de-

pISSN 2093-2278
eISSN 2093-2286



JPIS >
Journal of Periodontal
& Implant Science

Research Article

J Periodontal Implant Sci 2014;44:201-206
<http://dx.doi.org/10.5051/jpis.2014.44.4.201>

Received: Apr. 21, 2014

Accepted: Aug. 10, 2014

*Correspondence:

Amal Bouziane

Department of Periodontology, Faculty of Dental
Medicine, Mohammed V Souissi University, BP
6212 Madinat Al Irfane - Rabat, Morocco

E-mail: amal.bouziane@gmail.com

Tel: +212-64582629

Fax: +212-37681163

This is an Open Access article distributed under the terms
of the Creative Commons Attribution Non-Commercial
License (<http://creativecommons.org/licenses/by-nc/3.0/>).

fects, the goals of periodontal therapy are to alter or eliminate the microbial aetiology and contributing risk factors for periodontitis.

The prevalence rates of aggressive periodontitis in epidemiologic studies are variable. Prevalence ranges from approximately 1% to a maximum of 15% [2]. In Morocco, this form of periodontal disease has a reported prevalence of 7.6% [3].

Aggressive periodontitis, especially in its severe form, was traditionally considered to have an unfavourable prognosis, which often led to radical treatments. Eickholz et al. [4] reported that initial diagnosis was identified as a statistically significant influence on tooth loss. For aggressive and generalized severe chronic periodontitis, the risk for tooth loss was doubled in comparison with moderate periodontitis.

On the other hand, many studies [5-10] have demonstrated a favourable healing potential in lesions associated with localized aggressive periodontitis, but not in the severe generalized form. Other studies [11,12] have reported a stabilization of aggressive periodontitis cases achieved with both nonsurgical and surgical debridement. As well, many other studies have addressed only short-term therapeutic outcomes [13-15].

The aim of our study was to show clinical and radiographic outcomes following nonsurgical periodontal treatment alone in severe generalized aggressive periodontitis (SGAP) in the short and the long term.

MATERIALS AND METHODS

The present article reports on a retrospective cohort study of cases of SGAP treated with nonsurgical periodontal therapy. The study protocol was approved by the local Ethics Committee for Biomedical Research of the Faculty of Medicine and Pharmacy, Mohammed V Souissi University.

Patient population

Patients with advanced generalized aggressive periodontitis were

included in this study. They were referred for treatment at the Department of Periodontology, Ibn Sina University Hospital-Rabat.

The initial diagnosis was based on severe periodontal tissue destruction and angular bone destruction in healthy subjects under 30 years of age (Figs. 1 and 2). There was generalized interproximal attachment loss affecting at least three permanent teeth other than the first molars and incisors.

The inclusion criteria were: (1) clinically healthy patients, except for the presence of periodontitis, with SGAP; (2) a periodontally compromised tooth with clinical attachment loss of ≥ 5 mm and/or probing depth (PD) of ≥ 7 mm; (3) presence of intrabony defect with at least 50% bone loss.

The noninclusion criteria were: (1) patients with systemic disease or drug therapy that could lead to impaired wound healing; (2) patients with periodontal risk factors (such as smoking and diabetes mellitus).

Patients were treated for deep periodontal pockets and angular bone loss. In order to demonstrate the favourable outcomes of nonsurgical periodontal therapy, sites with advanced lesions that were treated by periodontal surgery (open flap debridement) were not included in the study.

All the patients received nonsurgical periodontal treatment, including oral hygiene instructions and complete periodontal debridement followed by an antimicrobial therapy based on chlorhexidine (0.12%) mouth rinse and antibiotic combination of 1,000 mg of amoxicillin and 750 mg of metronidazole per day for seven days.

Patients were enrolled in a maintenance care program and were provided with supportive periodontal care every six months.

Clinical and radiographic assessment

Estimations of samples in the study sample were based on a subject-level analysis. The following parameters were measured at severely affected sites with a millimeter calibrated periodontal probe, by the same examiner (A.B.) not blinded to the procedure: PD of pockets ≥ 7 mm, and clinical attachment level (CAL) of sites



Figure 1. Initial state of one of the severe generalized aggressive periodontitis cases.




Figure 2. Severe angular alveolar bone loss and furcation defect degree II at 36. Probing depth: 3.3.13 (Buccal), 3.2.11 (Lingual).



Figure 3. Improved pocket depth at re-evaluation. Probing depth: 2.2.7 (Buccal), 2.2.6 (Lingual).



Figure 4. Stability of the lesion 5 years after the aetiological treatment. Probing depth: 3.2.2 (Buccal), lingual 3.3.3.

with attachment loss ≥ 5 mm. For each patient, the mean of multiple measurements was taken at baseline (prior to the treatment), at re-evaluation after the initial treatment (Fig. 3), and posttreatment (Fig. 4). Responding sites were defined as those showing at least 2-mm reduction in probing pocket depth or 1 mm of attachment gain after treatment. Resolution of inflammation and bone fill were recorded dichotomously as present or absent for each patient. The resolution of inflammation was determined by a decrease of gingival bleeding and the modification of gingival aspect at least to score one according to the Loe and Silness index. Pre- and postoperative radiographs were taken with the long cone parallel technique and bone fill was defined by bone densification and/or augmentation of bone level (Fig. 5).

Statistical analysis

The data were analyzed using the patient as a unit (subject based analysis). For each patient, mean values of PD and CAL were calculated before the analysis. The statistical analysis was performed with a statistical program (SPSS Inc., Chicago, IL, USA). The analysis compared baseline values with re-evaluation and post-




Figure 5. Radiographic improvement of the lesion.

treatment values.

The quantitative variables (PD, CAL) were expressed as medians and quartiles and were analysed between the three paired groups using the Friedman test followed by *post hoc* analysis. The Wilcoxon test with the Bonferroni correction was used for each paired group. A level of $P < 0.05$ was accepted for statistical significance.

RESULTS

Seven patients with 266 sites were examined (mean age, 26.14 ± 2.61 years; male/female sex ratio, 1/6). Two hundred and twenty-six of these sites had a favourable response to nonsurgical periodontal therapy. Clinical and radiographic data are shown in Table 1.

Re-evaluation after initial treatment was carried out at a mean of three months (3.71 ± 1.4), with a maximum of 6 months after initial treatment. The posttreatment recall was performed at a median of five years (5 [1.5–7]). The results of the treated teeth in the seven patients are summarized in (Table 2).

A significant difference was found between the values of PD at baseline (7.94 [7.33–8.19] mm) and both re-evaluation (4.33 [3.63–5.08] mm) and posttreatment (3.54 [3.33–4.11] mm) values ($P = 0.002$). A significant difference was also found between the values of CAL at baseline (9.02 [7.5–9.2] mm) and both re-evaluation (6.55 [6.30–6.87] mm) and posttreatment (6.45 [5.70–6.61] mm) values ($P = 0.002$) (Figs. 6 and 7).

The inflammation was resolved and angular bone defects were repaired in all cases. A radiographic analysis carried out on angular bone defects showed significant repair of prior angular bone defects.

DISCUSSION

Our study has demonstrated that SGAP can be successfully treated by nonsurgical therapy alone. This therapy resulted in the termination of disease progression, resolution of inflammation, de-

Table 1. Clinical and radiographic data.

Patient (n=7)	No. of examined sites	No. of responding sites ^{a)}	Initial		Re-evaluation		Posttreatment	
			PD (mm)	CAL (mm)	PD (mm)	CAL (mm)	PD (mm)	CAL (mm)
1	49	45	8.19	9.02	3.32	6.30	3.49	6.61
2	71	47	7.21	7.39	5.14	6.32	3.86	4.79
3	14	13	7.33	8.67	4.33	6.55	4.11	6.50
4	48	48	7.94	9.04	3.63	5.96	3.54	5.70
5	35	34	9.15	11.66	3.85	6.87	3.33	6.45
6	26	21	8.17	9.20	5.08	7.40	4.29	7.20
7	23	18	7.53	7.50	4.47	6.83	3.26	6.17

PD: probing depth, CAL: clinical attachment loss.
^{a)}Sites with reduction of PD or attachment gain.

Table 2. Changes in PD and CAL in mm during the observed stages (n = 7).

	Initial	At re-evaluation	Posttreatment	P-value
PD (mm)	7.94 (7.33–8.19)*	4.33 (3.63–5.08)	3.54 (3.33–4.11)	0.002
CAL (mm)	9.02 (7.50–9.20)*	6.55 (6.30–6.87)	6.45 (5.70–6.61)	0.002

Values are presented as median (interquartile range).
 PD: probing depth, CAL: clinical attachment loss.

*P<0.05: re-evaluation and posttreatment vs. initial (Wilcoxon with Bonferroni correction).




Figure 6. Diagram demonstrating changes in pocket depth in millimeters (median and quartiles) at baseline, re-evaluation, and posttreatment after using Friedman and Wilcoxon tests. *P=0.05, re-evaluation vs. initial. †P=0.05, posttreatment vs. initial. ‡P=0.12, re-evaluation vs. posttreatment. PD: probing depth.

creased pocket depths, gains in clinical attachment and significant repair of alveolar defects associated with increased height of the supporting alveolar bone of the affected teeth. Aggressive periodontitis, especially in its severe generalized form, was previously considered to have an unfavourable prognosis. In some studies [11,12] the stabilization of periodontal health in aggressive periodontitis was achieved in sites with advanced disease after surgical




Figure 7. Diagram demonstrating changes in clinical attachment loss (CAL) in millimeters at baseline, at re-evaluation, and posttreatment after using Friedman and Wilcoxon tests. *P=0.05, re-evaluation vs. initial. †P=0.05, posttreatment vs. initial. ‡P=0.27, re-evaluation vs. posttreatment.

therapy. Aimetti et al. [13], Aimetti et al. [14], Darby et al. [16], Hughes et al. [17], and Yek et al. [18], have showed that nonsurgical treatment can be a highly effective treatment for patients and sites affected by aggressive periodontitis. Their results dealt with short-term outcomes. Further studies [15,19–22] have demonstrated the short-term clinical outcomes of nonsurgical periodontal debridement in subjects with generalized aggressive periodontitis.

In the present study, the posttreatment exam demonstrated a statistically significant decrease in PDs of deep periodontal pockets and significant gains in CAL. Changes in PD and CAL were observed from baseline to posttreatment. There was no relapse of the disease at the posttreatment stage. This observation suggests that nonsurgical treatment may have a long-term positive effect on the periodontal status of patients with SGAP. This is in accordance with the results of a five year longitudinal study demonstrating favourable healing following surgical and nonsurgical treatment of juvenile periodontitis [23].

Our patients were very pleased to retain their own natural teeth despite the advanced, seemingly hopeless state of their teeth prior to treatment. Pocket depth and attachment loss continued to decrease in most of the patients after the completion of nonsurgical periodontal therapy while alveolar bone repair continued to increase. The postoperative healing was continuous in six of the seven patients and was not dependent on recall frequency. The post-treatment radiographs demonstrate a significant formation of the mineralized component of the supporting mineralized tissues. The reparative potential associated with this form of periodontitis can be explained by the good healing potential of the periodontium in SGAP and the effectiveness of antimicrobials in modifying the microbiological profile of this form of periodontitis. The architecture of periodontal lesions in the severe aggressive form of periodontitis can also explain the positive therapeutic outcomes. The rapid rate of disease progression often leads to angular lyses and deep periodontal pockets. Increases in pocket depth were associated with a significant reduction after periodontal therapy. Machtei et al. [24] demonstrated that the greater the initial pocket depth was, the greater the potential of regeneration. This could explain the striking improvement of severe periodontal lesions.

Our study has demonstrated favourable radiographic and clinical outcomes of nonsurgical periodontal debridement in subjects with SGAP. The results of this study suggest that nonsurgical periodontal therapy, based on oral instructions, periodontal debridement, and the use of antimicrobials, may be an effective procedure to improve the healing potential of severe aggressive periodontitis cases. The reparative potential associated with severe aggressive periodontitis may exceed that observed in moderate aggressive and chronic periodontitis and suggests a conservative therapeutic approach.

This study provides evidence of favourable outcomes following nonsurgical therapy alone of SGAP. However, the results of the study are limited by the small study population, the retrospective design of the study and the heterogeneity of the time points of posttreatment recalls.

Overall, although this study provided some indications of the beneficial effect of nonsurgical therapy on SGAP cases in the short and the long term, a broader comparative study is necessary to more fully assess the effects of nonsurgical periodontal therapy compared to surgical therapy.

The therapeutic result of SGAP cases suggests that this form of periodontitis can have positive outcomes. The reparative potential associated with SGAP should encourage clinicians to save seemingly hopeless teeth in this type of periodontitis cases.

CONFLICT OF INTEREST

No potential conflict of interest relevant to this article was reported.

ACKNOWLEDGEMENTS

The authors thank Mr Driss Tarrouzi and Mrs Hafida Sahli, English teachers, for their clarifications and valuable contribution.

ORCID

Amal Bouziane <http://orcid.org/0000-0002-6723-5467>
Latifa Benrachadi <http://orcid.org/0000-0002-8571-0444>
Redouane Abouqal <http://orcid.org/0000-0002-6117-4341>
Oumkeltoum Ennibi <http://orcid.org/0000-0002-1224-9124>

REFERENCES

1. Parameter on aggressive periodontitis. American Academy of Periodontology. *J Periodontol* 2000;71(5 Suppl):867-9.
2. Demmer RT, Papapanou PN. Epidemiologic patterns of chronic and aggressive periodontitis. *Periodontol* 2000 2010;53:28-44.
3. Haubek D, Ennibi OK, Poulsen K, Poulsen S, Benzarti N, Kilian M. Early-onset periodontitis in Morocco is associated with the highly leukotoxic clone of *Actinobacillus actinomycetemcomitans*. *J Dent Res* 2001;80:1580-3.
4. Eickholz P, Kaltschmitt J, Berbig J, Reitmeir P, Pretzl B. Tooth loss after active periodontal therapy. 1: patient-related factors for risk, prognosis, and quality of outcome. *J Clin Periodontol* 2008; 35:165-74.
5. Dubrez B, Baehni P, Cimasoni G. A case of localized juvenile periodontitis: treatment and 3 years follow-up with superimposable radiographs. *J Clin Periodontol* 1996;23:557-62.
6. Sewon LA. Rapid bony healing in localized juvenile periodontitis: a case report. *Scand J Dent Res* 1993;101:371-4.
7. Novak MJ, Stamatelakys C, Adair SM. Resolution of early lesions of juvenile periodontitis with tetracycline therapy alone: long-term observations of 4 cases. *J Periodontol* 1991;62:628-33.
8. Mattout P, Moskow BS, Fourel J. Repair potential in localized juvenile periodontitis: a case in point. *J Periodontol* 1990;61:653-60.
9. Moskow BS. Healing potential in periodontal lesions: localized juvenile periodontitis. *J N J Dent Assoc* 1986;57:45-51.
10. Lindhe J, Liljenberg B. Treatment of localized juvenile periodontitis: results after 5 years. *J Clin Periodontol* 1984;11:399-410.
11. Buchmann R, Nunn ME, Van Dyke TE, Lange DE. Aggressive periodontitis: 5-year follow-up of treatment. *J Periodontol* 2002;73: 675-83.
12. Kamma JJ, Baehni PC. Five-year maintenance follow-up of early-onset periodontitis patients. *J Clin Periodontol* 2003;30:562-72.
13. Aimetti M, Romano F, Guzzi N, Carnevale G. One-stage full-mouth disinfection as a therapeutic approach for generalized aggressive periodontitis. *J Periodontol* 2011;82:845-53.
14. Aimetti M, Romano F, Guzzi N, Carnevale G. Full-mouth disinfection and systemic antimicrobial therapy in generalized aggressive periodontitis: a randomized, placebo-controlled trial. *J Clin Peri-*

- odontol 2012;39:284-94.
15. Griffiths GS, Ayob R, Guerrero A, Nibali L, Suvan J, Moles DR, et al. Amoxicillin and metronidazole as an adjunctive treatment in generalized aggressive periodontitis at initial therapy or re-treatment: a randomized controlled clinical trial. *J Clin Periodontol* 2011;38:43-9.
 16. Darby IB, Hodge PJ, Riggio MP, Kinane DF. Clinical and microbiological effect of scaling and root planing in smoker and non-smoker chronic and aggressive periodontitis patients. *J Clin Periodontol* 2005;32:200-6.
 17. Hughes FJ, Syed M, Koshy B, Marinho V, Bostanci N, McKay IJ, et al. Prognostic factors in the treatment of generalized aggressive periodontitis: I. Clinical features and initial outcome. *J Clin Periodontol* 2006;33:663-70.
 18. Yek EC, Cintan S, Topcuoglu N, Kulekci G, Issever H, Kantarci A. Efficacy of amoxicillin and metronidazole combination for the management of generalized aggressive periodontitis. *J Periodontol* 2010;81:964-74.
 19. Guerrero A, Griffiths GS, Nibali L, Suvan J, Moles DR, Laurell L, et al. Adjunctive benefits of systemic amoxicillin and metronidazole in non-surgical treatment of generalized aggressive periodontitis: a randomized placebo-controlled clinical trial. *J Clin Periodontol* 2005;32:1096-107.
 20. Kaner D, Bernimoulin JP, Hopfenmuller W, Kleber BM, Friedmann A. Controlled-delivery chlorhexidine chip versus amoxicillin/metronidazole as adjunctive antimicrobial therapy for generalized aggressive periodontitis: a randomized controlled clinical trial. *J Clin Periodontol* 2007;34:880-91.
 21. Mestnik MJ, Feres M, Figueiredo LC, Duarte PM, Lira EA, Faveri M. Short-term benefits of the adjunctive use of metronidazole plus amoxicillin in the microbial profile and in the clinical parameters of subjects with generalized aggressive periodontitis. *J Clin Periodontol* 2010;37:353-65.
 22. Baltacioglu E, Aslan M, Sarac O, Saybak A, Yuva P. Analysis of clinical results of systemic antimicrobials combined with nonsurgical periodontal treatment for generalized aggressive periodontitis: a pilot study. *J Can Dent Assoc* 2011;77:b97.
 23. Wennstrom A, Wennstrom J, Lindhe J. Healing following surgical and non-surgical treatment of juvenile periodontitis: a 5-year longitudinal study. *J Clin Periodontol* 1986;13:869-82.
 24. Machtei EE, Cho MI, Dunford R, Norderyd J, Zambon JJ, Genco RJ. Clinical, microbiological, and histological factors which influence the success of regenerative periodontal therapy. *J Periodontol* 1994;65:154-61.