

First Evaluation of an Outbreak of Bovine Babesiosis and Anaplasmosis in Southern Brazil Using Multiplex PCR

Mariana Feltrin Canever¹, Luisa Lemos Vieira¹, Carolina Reck¹, Luisa Richter², Luiz Claudio Miletti^{1,*}

¹Laboratório de Bioquímica de Hemoparasitas e Vetores, Centro de Ciências Agroveterinárias, Universidade do Estado de Santa Catarina, Lages, Brazil; ²Companhia Integrada de Desenvolvimento Agrícola de Santa Catarina, Brazil

Abstract: Outbreaks of tick-borne disease cases in Santa Catarina, Brazil are known, but the presence of the pathogen DNA has never been determined. In this study, the first survey of *Anaplasma marginale*, *Babesia bigemina*, and *Babesia bovis* DNA on blood samples of 33 cattle from an outbreak in Ponte Alta Municipality, Santa Catarina, Brazil, has been carried out. A multiplex PCR detected 54.5% of animals were co-infected with 2 or 3 parasites, while 24.2% were infected with only 1 species. The most prevalent agent was *B. bigemina* (63.6%) followed by *A. marginale* (60.6%). This is the first report of tick-borne disease pathogens obtained by DNA analysis in Southern Brazil.

Key words: *Babesia bovis*, *Babesia bigemina*, *Anaplasma marginale*, multiplex-PCR, Brazil

INTRODUCTION

Babesia bovis and *Babesia bigemina* in conjunction with *Anaplasma marginale* are intraerythrocytic pathogens that are responsible for the most prevalent and costly tick-borne diseases (TBDs) of cattle worldwide [1]. The Brazilian cattle industry estimates that losses reach about 500 million US dollars annually, according to data from Ministry of Agriculture, with these agents [2].

B. bovis and *B. bigemina* belong to the phylum Apicomplexa, Order Piroplasmid, and are the causative agents of bovine babesiosis in Latin America, where the only known vector is the host tick *Rhipicephalus (B.) microplus* [3]. Cattle suffering from babesiosis show massive destruction of erythrocytes because of the multiplication of hemoparasites inside these cells [4] with high morbidity and mortality worldwide [5].

A. marginale, is an intraerythrocytic rickettsia from family Anaplasmataceae [6] responsible for important economic losses in cattle in tropical and subtropical areas throughout the world, including Latin America [7]. Anaplasmosis can also

be transmitted by bloodsucking flies, horseflies, and mosquitoes [8]. *A. marginale* infection in the clinical signs consist of hemolytic anemia, jaundice, dyspnea, tachycardia, fever, fatigue, lacrimation, salivation, diarrhea, frequent urination, anorexia, and abortion, which in some cases can lead the animal to death in less than 24 hr [9].

The state of Santa Catarina in Southern Brazil has presented several outbreaks of TBD but without an efficient differential diagnosis. This lack of information on the causative agent of TBD causes small farmers use incorrect drugs for treatment of the disease. The aim of this work is to assess what are the causative agents of TBD in 1 outbreak in Santa Catarina, using the multiplex PCR. We report here the first molecular detection of *A. marginale*, *B. bovis*, and *B. bigemina* from cattle in Southern Brazil.

MATERIALS AND METHODS

Description of an outbreak

From November 2010 to March 2011, a total of 33 bovine blood samples were collected from Ponte Alta Municipality, Santa Catarina, Brazil, where an outbreak of tick-borne disease occurred. Samples were collected at 3 different properties of beef cattle from Limousin and Charolais bred with ages between 8 months and 4 years. Three animals showed characteristic clinical signs of tick-borne diseases, and other samples were

•Received 24 April 2014, revised 5 August 2014, accepted 18 August 2014.

*Corresponding author (lcmiletti@yahoo.com.br)

© 2014, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

collected from animals of the aforementioned properties. The properties had the presence of *Rhipicephalus (B.) microplus* ticks.

Sample size determination

The determination of the sample size was based on statistical and epidemiological concepts, considering the population of cattle in Ponte Alta Municipality of 16,200 animals. The minimum number of samples (n=28) was calculated based on consideration of the disease prevalence of 10% TBD, using the proposal methodology by De Canno Roe [10], for an infinite population (n), with 95% reliability.

Extraction of DNA

DNA was extracted from whole blood using phenol-chloroform method. Briefly, 200 µl of whole blood was mixed with lysis buffer (10 mM Tris pH 7.4, 10 mM NaCl, 25 mM EDTA, 1% SDS) with 100 µg/ml of proteinase K (Sigma, St. Louis, Missouri, USA) and incubated at 42°C for 12 hr. Subsequently, 3 washes with phenol, phenol-chloroform (1:1), and chloroform, respectively, were performed, centrifuging to 14,000 g for 10 min and removing the supernatant. The DNA was precipitated with isopropanol and then washed with 70% ethanol. The tubes were placed in an oven at 37°C until complete evaporation of the alcohol. The resulting DNA was eluted in 50 µl of Milli-Q water-free DNase and stored at -20°C.

Multiplex PCR conditions

PCR amplification was carried out on genomic DNA extracted from cattle blood samples, and analyzed by multiplex PCR for the screening of *B. bovis*, *B. bigemina*, and *A. marginale*.

Briefly, PCR assay was performed in a total volume of 25 µl containing 80 ng of the genomic DNA and 8.5 pmoles of each primer pair (Table 1); 1U *Taq* DNA polymerase, GoTaq® Hot Start Polymerase (Promega; Madison, Wisconsin, USA), 0.2 mM dNTPs, 25 mM MgCl₂, 5 µl buffer (5X Green GoTaq® Flexi buffer (Promega) and ultra-pure water to make the final volume. Amplification involved a hot start of 5 min at 95°C,

followed by 35 cycles of 1 min at 95°C, 1 min at 58°C, 1 min at 72°C, and a final extension step of 10 min at 72°C.

The amplified PCR products were subjected to electrophoresis on 2% agarose gel stained with ethidium bromide, visualized under UV light, and photographed in trans-illuminator apparatus. The amplicons molecular sizes were estimated by including a 100 bp ladder standard (Ludwig Biotecnologia, Porto Alegre, Brazil). DNA positive controls of *A. marginale*, *B. bigemina*, and *B. bovis* were kindly provided by Dr. Luciana Brito Gatto, Embrapa Porto Velho, RO and Dr. Odilon Vidotto, UEL, PR. Negative control was distilled water.

Sequencing

Purified PCR products were then ligated into the p-GEM T easy® vector (Promega) and cloned by transformation into DH5 α *Escherichia coli* calcium competent cells. Single picked colonies were grown overnight, and plasmids were purified using the QIAGEN plasmid purification kit (QIAGEN, Hilden, Germany). To confirm plasmids contain the correct insert, between 1 and 2 µg DNA was purified from positive colonies and sequenced by a commercial sequencing service (ACTGene, Porto Alegre, Brazil). All high quality DNA sequences (Phred P20) were analyzed using the Phred/Phrap/Consed package [11]. The identity of the sequences were confirmed with the BLAST tool (www.ncbi.nlm.nih.gov/blast) [12].

RESULTS

Thirty-three blood samples from Ponte Alta Municipality, Santa Catarina, Brazil were examined for infection with *A. marginale*, *B. bigemina*, and *B. bovis* using multiplex PCR method. A summary of the results is shown in Table 2. When the samples were assayed using multiplex PCR, 3 and 5 animals were found to have single infections with *A. marginale* and *B. bigemina*, respectively. Further 1, 2, and 12 animals had double infections with *B. bigemina/B. bovis*, *A. marginale/B. bovis*, and *A. marginale/B. bigemina*, respectively. Three animals had

Table 1. Oligonucleotide primer sequences used in multiplex PCR

Pathogen	Primer	Sequence (5'-3')	Amplicon size (bp)	References
<i>Babesia bigemina</i>	BiIA	CAT CTA ATT TCT CTC CAT ACC CCT CC	278 pb	[30]
	BiIB	CCT CGG CTT CAA CTC TGA TGC CAA AG		
<i>Babesia bovis</i>	BoF BoR	CAC GAG GAA GGA ACT ACC GAT GTT GA	356 pb	[31]
		CCA AGG AGC TTC AAC GTA CGA GGT CA		
<i>Anaplasma marginale</i>	1773F	TGT GCT TAT GGC AGA CAT TTC C	1,000 pb	[32]
	2957R	AAA CCT TGT AGC CCC AAC TTA TCC		

Table 2. Multiplex PCR test on field bovine blood samples

	No. of PCR positive animals							No. of PCR negative animals	Total no. of animals
	Am	Bbi	Bbo	Am/Bbi	Am/Bbo	Bbi/Bbo	Am/Bbi/Bbo		
Multiplex PCR	3	5	0	12	2	1	3	7	33
Percentage (%)	24.2 (n=8)			54.6 (n=18)				21.2	100

Am, *Anaplasma marginale*; Bbi, *Babesia bigemina*; Bbo, *Babesia bovis*.

Table 3. Blood samples from 33 cattle using specific PCR amplification and number of samples showing simultaneous specific PCR amplification of 2 or 3 of the targeted micro-organisms

Detection of pathogen DNA			Co-detection of pathogen DNA			
Number of animals infected/ number of samples (%)			Number of animals infected/ number of samples (%)			
Am	Bbi	Bbo	Am/Bbi	Am/Bbo	Bbi/Bbo	Am/Bbi/Bbo
20/33 (60.6)	21/33 (63.6)	6/33 (18.2)	12/33 (36.4)	2/33 (6.1)	1/33 (3.0)	3/33 (9.1)

Am, *Anaplasma marginale*; Bbi, *Babesia bigemina*; Bbo, *Babesia bovis*.




Fig. 1. Analysis of mPCR products by agarose gel electrophoresis. M, 100 bp molecular size marker; lanes 1, 8, and 9, positive mPCR controls; lane 6, negative mPCR control (distilled water); lanes 2-3 (*B. bigemina*/*B. bovis*), 4 (*B. bigemina*/*A. marginale*), and 5 (*B. bigemina*/*A. marginale*), template DNA. Arrows indicate 278, 356, and 1,000 base pair amplicons, respectively, for *B. bigemina*, *B. bovis*, and *A. marginale* generated using primer sets described in materials and methods.

multiple infections with the 3 organisms, and 7 were negative. As shown in Table 3, *B. bigemina* is the most prevalent agent (63.6%) followed by *A. marginale* (60.6%). An example of the results obtained using multiplex PCR on field samples is illustrated in Fig. 1.

The identity of the amplicons was guaranteed by sequencing, demonstrating identity with sequences deposited in GenBank, respectively, with 93% (DQ501244.1) to *A. marginale*, 99% (AF027149) to *B. bovis*, and 83% of (S45366) for *B. bigemina*.

DISCUSSION

In this study, multiplex PCR was used for the first time to

detect TBD agents in cattle blood samples from an outbreak occurred in Santa Catarina, Brazil. The multiplex PCR showed high specificity for use in epidemiological studies as described [13-16].

The state of Santa Catarina is in enzootic stability for TBD [17], since the distribution of *A. marginale* and *Babesia* sp. were closely associated with the presence of the cattle tick *R. (B.) microplus* which is highly prevalent in Santa Catarina state [18] and by the high level of seroprevalence for *Babesia* sp. [19]. However, due to climatic factors, the study area can be considered in some periods of the year in enzootic instability [20].

The prevalence of TBD agents in Southern Brazil are mainly based on serological analysis [2,21,22]. In the state of Santa Catarina, only 2 papers were published, and they demonstrated high seroprevalence to *B. bigemina* and *B. bovis* [23,24]. In the same region, no studies were conducted on the prevalence of *A. marginale*. Despite knowledge of the circulation of TBD agents, to date, no work on molecular analysis was conducted in the state of Santa Catarina.

The results obtained using the field samples clearly indicate the prevalence of mixed infections in the field, so the multiplex PCR will facilitate the assessment of risk factors associated with TBD and this is important for improved design and implementation of cost-effective control strategies in endemic and non-endemic regions [25]. Multiplex PCR results obtained from field samples showed that 54.5% of animals were co-infected with 2 or 3 parasites, while 24.2% were infected with only 1 species. In Turkey [16], it was found that 24% of animals were co-infected with *A. marginale* and/or *B. bovis*, and a prevalence of 12% dual infection with *B. bovis* and *B. bigemi-*

na was observed in single PCR in Egypt [26]. Comparisons with other data are not possible due to the lack of researches using multiplex PCR for detection of these pathogens.

Of the 3 tick-borne diseases investigated in the present study, *B. bigemina* and *A. marginale* were found to be most prevalent with 63.6% and 60.6% prevalence rate, respectively. The low infection rate for *B. bovis* (18.2%) infections in cattle was related with low parasitemia during the acute phase of the disease, a feature of these agent [27]. Cattle that survive initial infection and also calves partially resistant to this disease remain persistently infected [28] with very low parasite levels that are sufficient for transmitting infection to ticks. Such cattle are potentially an important factor in the maintenance of *B. bovis* in endemic areas as well as a suspect source for introduction and spread of the disease to non-endemic areas where competent vectors are present [29]. These results highlight the situation in Santa Catarina where, although infection with *A. marginale* and *B. bigemina* appears to be the main risk, there is also a clear risk of co-infection between the 3 pathogens.

In conclusion, we could confirm that in Santa Catarina, like other states in Southern Brazil, the 3 TBD pathogens are circulating and knowledge of these data should be available to producers and government agencies.

ACKNOWLEDGMENTS

This work was supported by Universidade do Estado de Santa Catarina (UDESC), Brazil. Mariana Feltrin Canever was a fellowship of PIC program from UDESC, Brazil.

CONFLICT OF INTEREST

We have no conflict of interest related to this work.

REFERENCES

1. Suarez CE, Noh S. Emerging perspectives in the research of bovine babesiosis and anaplasmosis. *Vet Parasitol* 2011; 180: 109-125.
2. Almeida MB, Tortelli FP, Riet-Correa B, Ferreira JLM, Soares MP, Farias NAR, Riet-Correa F, Schild AL. Tristeza parasitária bovina na região sul do Rio Grande do Sul: estudo retrospectivo de 1978-2005. *Pesq Vet Bras* 2006; 26: 237-242.
3. Guglielmone AA. Epidemiology of babesiosis and anaplasmosis in South and Central America. *Vet Parasitol* 1995; 57: 109-119.
4. Bock R, Jackson L, de Vos A, Jorgensen W. Babesiosis of cattle. *Parasitology* 2004; 129: S247-S269.
5. McCosker PJ. The global importance of babesiosis. In: Ristic M, Krier JP, eds. *Babesiosis*. New York, USA. Academic Press. 1981, p 1-24.
6. Hornok S, Földvári G, Elek V, Naranjo V, Farkas R, de La Fuente J. Molecular identification of *Anaplasma marginale* and rickettsial endosymbionts in blood-sucking flies (Diptera: Tabanidae, Muscidae) and hard ticks (Acari: Ixodidae). *Vet Parasitol* 2008; 154: 354-359.
7. Harrus S, Baneth G. Drivers for the emergence and re-emergence of vector-borne protozoal and bacterial diseases. *Int J Parasitol* 2005; 35: 1309-1318.
8. Kocan KM, de La Fuente J, Blouin EF, Coetzee JF, Ewing SA. The natural history of *Anaplasma marginale*. *Vet Parasitol* 2010; 167: 95-107.
9. Vidotto O, Marana ERM. Diagnóstico em anaplasmose bovina. *Ciência Rural (Santa Maria)* 2001; 31: 361-368.
10. Thrusfield M. *Veterinary Epidemiology*. 3rd ed. Oxford, UK. Blackwell Publishing Co. 2003, p 46-69.
11. Ewing B, Green P. Base-calling of automated sequencer traces using phred. II. Error probabilities. *Gen Res* 1998; 8: 186-194.
12. Altschul SF, Madden TL, Schäffer AA, Zhang J, Zhang Z, Miller W, Lipman DJ. Gapped BLAST and PSI-BLAST: a new generation of protein database search programs. *Nucl Acids Res* 1997; 25: 3389-3402.
13. Figueroa JV, Chieves LP, Jonhson GS, Buening GM. Multiplex polymerase chain reaction based assay for the detection of *Babesia bigemina*, *Babesia bovis*, and *Anaplasma marginale* DNA in bovine blood. *Vet Parasitol* 1993; 50: 69-81.
14. Oliveira-Sequeira TCG, Oliveira, MCS, Araujo JP Jr, Amarante AFT. PCR-based detection of *Babesia bovis* and *Babesia bigemina* in their natural host *Boophilus microplus* and cattle. *Int J Parasitol* 2005; 35: 105-111.
15. Silva MG, Henriques G, Sánchez C, Marques, PX, Suarez CE, Oliva A. First survey for *Babesia bovis* and *Babesia bigemina* infection in cattle from Central and Southern regions of Portugal using serological and DNA detection methods. *Vet Parasitol* 2009; 166: 66-72.
16. Bilgic HB, Karagenç T, Simuunza M, Shiels B, Tait A, Eren H, Weir W. Development of a multiplex PCR assay for simultaneous detection of *Theileria annulata*, *Babesia bovis*, and *Anaplasma marginale* in cattle. *Exp Parasitol* 2013; 133: 222-229.
17. Nuerberg S, Serra-Freire NM. O carrapato e a tristeza parasitária bovina no estado de Santa Catarina. *Rev Bras Med Vet* 1995; 17: 257-259.
18. Souza AP, Gonzales JC, Ramos CI, Paloschi CG, Moraes AN. Variação sazonal de *Boophilus microplus* no Planalto Catarinense. *Pesq Agrop Bras* 1988; 23: 627-630.
19. Madruga CR, Aycardi E, Kessler RH, Schenk MAM, de Figueiredo GR, Curvo JBE. Níveis de anticorpos anti-*Babesia bigemina* e *Babesia bovis* em bezerros da raça Nelore, Ibagé e cruzamento de Nelore. *Pesq Agrop Bras* 1984; 19: 1163-1168.
20. Mahoney DF, Ross DR. Epizootiological factors in the control of

- bovine babesiosis. Aust Vet J 1972; 48: 292-298.
21. Schettlers TPM, Kleuskens J, Scholtes N, Gorenflot A. Parasite localization and dissemination in the *Babesia*-infected host. Ann Trop Med Parasitol 1998; 92: 513-519.
 22. Osaki SC, Vidotto O, Marana, ERM, Vidotto, M.C, Yoshihara E, Pacheco RC, Igarashi M, Minho AP. Occurrence of antibodies against *Babesia bovis* and studies on natural infection in Nelore cattle, in Umuarama municipality, Paraná State, Brazil. Rev Bras Med Vet 2002; 11: 77-83.
 23. Dalagnol AA, Martins E, Madruga CR, Gomes R, Schenk MAM, Kessler RH, Gratão G, Gales ME, Schenk JAP, Andreade M, Bianchini I, Míguita, M. Prevalência de agentes da tristeza parasitária bovina em bovinos de corte na região de clima cfb-SC. Agrop Cat 1999; 12: 46-47.
 24. Souza AP, Bellato V, Sartor AA, Farias LM, Prevalência de anticorpos anti-Babesia em bovinos no Planalto Norte de Santa Catarina. Rev Ciênc Agrovet 2002; 1: 21-23.
 25. Simuunza M, Weir W, Courcier E, Tait A, Shiels B. Epidemiological analysis of tick-borne diseases in Zambia. Vet Parasitol 2011; 175: 331-342.
 26. Adham FK, Abd-El-Samie EM, Gabre RM, Hussein HE. Detection of tick blood parasites in Egypt using PCR assay I—*Babesia bovis* and *Babesia bigemina*. Parasitol Res 2009; 105: 721-730.
 27. Kessler RH, Madruga CR, Schenk MAM, Ribeiro OC. Babesiose cerebral por *Babesia bovis* (Babés 1888 Starcovici 1893) em bezerras no Estado de Mato Grosso do Sul. Pesq Agrop Bras 1983; 18: 931-935.
 28. Brown WC, Norimine J, Knowles DP, Goff WL. Immune control of *Babesia bovis* infection. Vet Parasitol 2006; 138: 75-87.
 29. Howell JM, Ueti MW, Palmer GH, Scoles GA, Knowles DP. Persistently infected calves as reservoirs for acquisition and transovarial transmission of *Babesia bovis* by *Rhipicephalus (Boophilus) microplus*. J Clin Microbiol 2007; 45: 3155-3159.
 30. Figueroa JV, Chieves LP, Johnson GS, Buening GM. Detection of *Babesia bigemina*-infected carriers by polymerase chain reaction amplification. J Clin Microbiol 1992; 30: 2576-2582.
 31. Suarez CE, Palmer GH, Jasmer DP, Hines SA, Perryman LE, McElwain TF. Characterization of the gene encoding a 60-kilodalton *Babesia bovis* merozoite protein with conserved and surface exposed epitopes. Mol Biochem Parasitol 1991; 46: 45-52.
 32. Lew AE, Bock RE, Minchin CM, Masaka S. A *msp1α* polymerase chain reaction assay for specific detection and differentiation of *Anaplasma marginale* isolates. Vet Microbiol 2002; 86: 325-335.