

Three Echinostome Species from Wild Birds in the Republic of Korea

Seongjun Choe¹, Dongmin Lee¹, Hansol Park¹, Mihyeon Oh¹, Hyeong-Kyu Jeon¹, Youngsun Lee², Ki-Jeong Na^{2,3}, Youngjun Kim⁴, Hang Lee⁵, Keeseon S. Eom^{1,*}

¹Department of Parasitology, Medical Research Institute and Parasite Resource Bank, Chungbuk National University School of Medicine, Cheongju 361-763, Korea; ²The Wildlife Center of Chungbuk, Cheongju 361-763, Korea; ³College of Veterinary Medicine, Chungbuk National University, Cheongju 361-763, Korea; ⁴Chungnam Wild Animal Rescue Center, Kongju National University, Yesan 340-702, Korea; ⁵Conservation Genome Resource Bank for Korean Wildlife, Research Institute for Veterinary Science and College of Veterinary Medicine, Seoul National University, Seoul 151-742, Korea

Abstract: Three echinostome species, i.e., *Patagifer bilobus*, *Petasiger neocomense*, and *Saakotrema metatestis*, are newly recorded in the trematode fauna of the Republic of Korea. They were recovered from 3 species of migratory birds (*Platalea minor*, *Podiceps cristatus*, and *Egretta garzetta*), which were donated by the Wildlife Center of Chungbuk (WCC) and the Conservation Genome Resource Bank for Korean Wildlife (CGRB). Only 1 *P. bilobus* specimen was recovered from the intestine of a black-faced spoonbill (*P. minor*), and characterized by the bilobed head crown with a deep dorsal incision and 54 collar spines. Twenty *P. neocomense* were recovered from the intestine of a great crested grebe (*P. cristatus*), and they had a well-developed head crown with 19 spines and 2 testes obliquely located at the posterior middle of the body. Total 70 *S. metatestis* were collected from the bursa of Fabricius of 1 little egret (*E. garzetta*). It is characterized by stout tegumental spines covered in the entire leaf-shaped body, posterior extension of the uterus, presence of the uroproct and a well-developed head crown with 12 pairs of collar spines on each side. By the present study, these 3 echinostome species are newly added to the trematode fauna in Korea.

Key words: *Petasiger neocomense*, *Patagifer bilobus*, *Saakotrema metatestis*, echinostome, intestinal trematode, migratory bird

INTRODUCTION

Echinostomes (=Family Echinostomatidae) are flukes that can be easily distinguished from other flukes thanks to their typical head collars as well as the presence of collar spines around the anterior part [1]. This trematode group is composed of 355 species and 6 subspecies in 50 genera, and they are parasites of birds, mammals, and humans [2]. In Korea, total 14 echinostome species in 7 genera have been reported: *Acanthoparyphium marilae* Yamaguti, 1934; *Acanthoparyphium tyosenense* Yamaguti, 1939; *Echinochasmus japonicus* Tanabe, 1926; *Echinochasmus perfoliatus* (Ratz, 1908) Dietz, 1909; *Echinoparyphium recurvatum* (Von Linstow, 1873) Dietz, 1909; *Echinostoma cinetorchis* Ando and Ozaki, 1923; *Echinostoma gotoi*

Ando and Ozaki, 1923; *Echinostoma hortense* Asada, 1926; *Echinostoma miyagawai* Ishii, 1932; *Echinostoma revolutum* (Froelich, 1802) Looss, 1899; *Euparyphium murinum* Tubanguui, 1931; *Himasthla alincia* Dietz, 1909; *Himasthla kusiasigi* Yamaguti, 1939, and *Stephanoprora* sp. [3-13].

Echinostomes of birds have been investigated only in a few avian species in Korea. Total 8 species, i.e., *A. marilae*, *A. tyosenense*, *E. japonicus*, *E. recurvatum*, *E. gotoi*, *E. miyagawai*, *E. revolutum*, and *H. kusiasigi*, have been reported as the avian echinostomes. Their hosts were recorded as follows: *A. marilae* from the black scoter, *A. tyosenense* from the ruddy turnstone, great knot, black-tailed gull, white-winged scoter, and black scoter, *E. japonicus* from the domestic duck and great egret, *E. recurvatum* from the mallard, spot-billed duck, and great crested grebe, *E. gotoi* from the mallard and spot-billed duck, *E. miyagawai* from the mallard, domestic duck, and spot-billed duck, *E. revolutum* from the falcated teal, mallard, spot-billed duck, and great scaup and *H. kusiasigi* from the dulin [3,5,7,9,14-16].

The present report provides a detailed description of the morphological features and measurements of 3 so far unrecorded

•Received 6 December 2013, revised 24 June 2014, accepted 15 July 2014.

*Corresponding author (kseom@chungbuk.ac.kr)

© 2014, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

echinostome species in Korea: *Patagifer bilobus* (Rudolphi, 1819) Dietz, 1909; *Petasiger neocomense* Fuhrmann, 1927, and *Saakotrema metatestis* (Saakova, 1952) Skrjabin & Baschkirova, along with a review of those that were previously reported.

MATERIALS AND METHODS

Parasite specimens were recovered from the visceral organs of their hosts. Bird carcasses were donated by the Wildlife Center of Chungbuk (WCC) and the Conservation Genome Resource Bank for Korean Wildlife (CGRB). A black-faced spoonbill (*Platalea minor*), coded as CGRB12252, was collected from Yeocha-ri, Hwado-myeon, Ganghwa-gun, Incheon Metropolitan City, Korea in May 2010. A great crested grebe (*Podiceps cristatus*) was collected from a factory district in Cheongju-si, Chungcheongbuk-do, Korea in February 2011. The host animal had a fractured clavicle and hypothermia. Veterinarians of the WCC tried to save the bird but it eventually died. A little egret (*Egretta garzetta*, WCC no. 20120224) with severe fractures was found in Yeongdong-gun, Chungcheongbuk-do, Korea, in September 2013 and was necropsied immediately after death. Carcasses of 2 additional birds were donated to our laboratory for parasitological examinations and stored at -20°C.

Visceral organs of the hosts were dissected and washed with tap water. Parasitic worms were recovered by filtration through sieves (mesh size: 250 µm to 1 mm). Collected worms were flattened with a coverslip pressure and fixed with 10% neutral buffered formalin. The specimens were stained with Semichon's acetocarmine and were dehydrated through a graded alcohol series. After clearing with xylene, morphological characteristics of each specimen were observed with a light microscope with a reticle.

Examined specimens were deposited at the National Institute of Biological Resources (NIBR), Incheon Metropolitan City, Korea and the Parasite Resource Bank (PRB), Cheongju City, Chungcheongbuk-do, Korea.

RESULTS

Family Echinostomatidae Looss, 1899

Genus *Patagifer* Dietz, 1909

Patagifer bilobus (Rudolphi, 1819) Dietz, 1909

One specimen was recovered from the intestine of a black-faced spoonbill (*Platalea minor*, CGRB12252). NIBR specimen number is KOSPIV0000193866.

Description (n=1): A ribbon-shaped body with blunt end. The longitudinal muscle was well developed. Tegumental spines were absent. The body length was 12.3 mm and the width was 1.6 mm at the level of the uterus and 1.3 mm at the ventral sucker. The ratio between the forebody and body length was 12.2%. The head collar was large and well developed, bilobed, fan shaped, and with a deep incision. The collar was 937 µm in length and 1,415 µm in width. A total of 54 collar spines were present. Lateral spines were similar in size to the angle spines. The oral sucker was small, measured 283×370 µm. The pharynx was small, oval in shaped and located immediately behind the oral sucker, measured 253×224 µm. The esophagus was 341 µm, with an intestinal bifurcation anterior to the cirrus sac. The cecum ran through the lateral side of the body to the posterior extremity. The cirrus sac was elongate-oval in shape, measured 468×263 µm and was located anteriodorsally to the ventral sucker. The ventral sucker was large, well developed, and measured 1,190×995 µm. The uterus was filled with numerous eggs, loops intercaecal, and located between the ventral sucker and ovary. Eggs in the uterine loops (n=23) measured 95-105 (96±18)×52.5-57.5 (54±1) µm. The ovary was oval in shape, displaced by the uterine loops, and measured 175×335 µm. Mehlis' gland was oval in shape, located behind the ovary, and measured 458×292 µm. The vitelline follicles were located laterally on the body, between the ventral sucker and the posterior extremity and overlapped the cecum. The excretory vesicle and the pore were obscured. There were 2 testes, tandem: the anterior testis measured 1,035×332 µm and the posterior testis measured 1,308×332 µm (Fig. 1A-C).

Genus *Petasiger* Dietz, 1909

Petasiger neocomense Fuhrmann, 1927

Total 20 specimens were recovered from the intestine of a great crested grebe (*Podiceps cristatus*). NIBR specimen number is KOSPIV0000192593.

Description (n=12): The body was fusiform, 1,191-1,470 (1,348±91) µm in length and 347-425 (395±26) µm in width at the level of the ventral sucker. It was covered with minute tegumental spines. The head collar was well developed, being 132-177 (145±14) µm in length, 213-272 (247±17) µm in width, and armed with 19 spines; 8 angle spines, 6 lateral spines and 5 dorsal spines. The angle spine was larger than the other spines, measuring 81-107 (93±7, n=37) µm long and 14-18 (15±1, n=37) µm wide, whereas the lateral spines were 53-61 (56±2, n=9) µm long and 8-13 (9±1, n=




Fig. 1. Light microscopic view of *Patagifer bilobus* (A-C) and *Petasiger neocomense* (D-F). (A) Photo of *P. bilobus*. (B) Drawing of *P. bilobus*. (C) Head part of the worm. (D) Photo of *P. neocomense*. (E) Drawing of *P. neocomense*. (F) Collar spines of the worm.

9) μm wide and the dorsal spines were 58-77 (67 ± 5 , $n=16$) μm long and 9-15 (12 ± 1 , $n=16$) μm wide. The oral sucker was subterminal, circular to oval in shape, 74-113 (88 ± 12) μm long and 73-117 (88 ± 13) μm wide. The prepharynx was 23-99 (55 ± 20) μm long. The pharynx was oval, 65-79 (71 ± 4) μm long and 49-61 (53 ± 6) μm wide. The esophagus was 179-357 (262 ± 55) μm in length. The intestinal bifurcation was anterior to the cirrus sac, which was muscular, located in the anteriopdorsal region of the ventral sucker, and measured 120-181 (154 ± 23) μm in length and 74-100 (84 ± 8) μm in width. The cirrus, as observed in almost all specimens, was thin and long, measuring 17-30 (23 ± 4) μm in width. The ventral sucker was large, located in the center of the body, measured 241-294 (274 ± 18) μm long and 232-313 (281 ± 26) μm wide. 2

testes were present, elongate-oval in shape, and positioned oblique or symmetrical. The anterior testis measured 107-181 (145 ± 22) μm in length and 55-87 (71 ± 10) μm in width, and the posterior testis measured 107-181 (145 ± 22) μm in length and 49-87 (68 ± 10) μm in width. The ovary was round, dextral, close to ventral sucker and measured 38-74 (54 ± 10) μm in diameter. Mehlis' gland was elongate-oval, larger than ovary, 45-63 (57 ± 6) μm long and 104-179 (133 ± 22) μm wide. Vitelline follicles were only present in the hind body. The uterus was short and located behind the ventral sucker, and it contained only 1 or 2 eggs. The excretory vesicle was Y-shaped. Eggs were oval and measured 66-86 (73 ± 7) μm long and 38-50 (43 ± 4) μm wide (Fig. 1D-F).

Genus *Saakotrema* Skrjabin & Baschkirova, 1956
Saakotrema metatestis (Saakova, 1952) Skrjabin & Baschkirova, 1956

Total 70 specimens were collected from the bursa of Fabricius of 1 little egret (*Egretta garzetta*). Most of the worms recovered were damaged except for 3 intact specimens. NIBR specimen number is KOSP1V0000192596.

Description (n=3): A leaf-shaped body measuring 1.7-2.0 (1.86) mm long and 0.5-0.6 (0.55) mm wide, with the widest point at the level of the ventral sucker and narrowing at the posterior end. The entire body was covered with strong tegumental spines, 17 μ m in length. The collar was well developed and measured 88-133 (115) \times 185-223 (203.3) μ m. There were 24 collar spines in a single row. The oral sucker was well developed and measured 43-80 (66.7) \times 55-80 (68.3) μ m. The phar-

ynx was oval shaped, larger than oral sucker and measured 105-110 (108.3) \times 63-80 (70.0) μ m. The intestinal bifurcation was located between the pharynx and the ventral sucker. Ventral sucker was large, well developed and measuring 278-300 (285.0) \times 260-303 (280.8) μ m. The cirrus sac was obscure, oval shaped, and measured 135 \times 75 μ m (measurement taken from only 1 specimen). The ovary was oval in shape and measured 75-130 (108.3) \times 113-158 (132.5) μ m. The testes were tandem, round to oval in shape, and were located immediately behind the ovary. The anterior testis measured 244-254 (240.6) \times 283-342 (306.0) μ m and the posterior testis measured 322-332 (325.5) \times 234-322 (266.9) μ m. The uterine coil was long and extended posteriorly through the ventral side of the worm. Eggs were oval and measured 70-83 (74.2 \pm 4.1) \times 43-48 (45.1 \pm 2.9) μ m (Fig. 2A-D).




Fig. 2. Light microscopic view of *Saakotrema metatestis*. (A) Photo of an adult worm. (B) Description of adult worm. (C, D) Collar spines of *S. metatestis*.

DISCUSSION

The present study revealed 3 unrecorded echinostome species in Korea. Since the first report of *Echinostoma revolutum* by Issiki [3], more than 14 echinostome species have been identified in Korea. Park [4] recovered *Echinostoma hortense* from rats, while Yamaguti [5] identified *Acanthoparyphium marilae* in *Melanitta nigra*, as well as *A. tyosenense* in *M. nigra americana* and *M. fusca stejnegeri* as new species in Korea. *E. cinetorchis* and *Euparyphium murinum* were also recovered from wild rats by Seo et al. [6]. Chu et al. [7] reported the presence of *E. revolutum*, *E. gotoi*, *E. miyagawai*, and *Echinoparyphium koizumi* in wild birds. *Echinochasmus perfoliatus* was found from cats in Daegu-si [8]. Eom et al. [9] found *E. miyagawai* and *E. japonicus* in domestic ducks in Korea. However, *E. miyagawai* was considered as synonym of *E. revolutum* by Yamaguti [2,17] and Bykhovskaya-Pavlovskaya [18], but this species was re-validated upon closer examination of various characteristics, including the morphology and life cycle [19,20]. Lee et al. [11] found *E. revolutum* and *Echinoparyphium recurvatum* from house rats in Yangyang-gun, Gangwon-do and considered *E. koizumi* that reported by Chu et al. [7] as a synonym of *E. recurvatum*.

On the other hand, *Himasthla kususigi* metacercariae were detected in the marine clam, *Meretrix lusoria*, and subsequently adults were recovered in the black-tailed gull, *Larus crassirostris*, experimentally infected in Korea [10]. This fluke was also recovered from the dunlin, *Calidris alpina sakhalina*, naturally infected in Ganghwa-gun, Gyeonggi-do [16]. Sohn and Chai [13] found *E. revolutum*, *E. hortense*, *E. japonicus*, and *Stephanoprora* in feral cats purchased in a market in Busan Metropolitan City. Metacercariae of *Himasthla alincia* were collected from an intermediate host, *Mactra veneriformis*, and a study was performed by infecting chicks [12]. However, the natural definitive hosts of this fluke have not been reported in Korea. Therefore, total 14 echinostome species were described in the Republic of Korea before the present study. Including the 3 new faunas in this study, a total of 17 species of 10 genera, i.e., *Acanthoparyphium* (2 species), *Echinochasmus* (2), *Echinoparyphium* (1), *Echinostoma* (5), *Euparyphium* (1), *Himasthla* (2), *Petasiger* (1), *Patagifer* (1), *Saakotrema* (1), and *Stephanoprora* (1), have been recorded to be distributed in Korea until now [6-16,21-26] (Table 1).

The worms recovered from *Podiceps cristatus* had 19 collar spines, including obvious longer angle spines, oblique to symmetrical testes, an elongate-oval cirrus sac and a tubular cirrus. These morphological characters were in line with those of *P.*

neocomense, including size variation [27]. *P. neocomense* was first recovered from *P. cristatus* in Lake Neuchâtel by Fuhmann [28]. In Korea, the host of *P. neocomense* is also *P. cristatus*. *P. neocomense* resembles *P. nitidus* morphologically but can be easily distinguished by the shape of the cirrus and the cirrus sac, as well as by geography (Eurasia vs North America) [27].

P. bilobus is the type-species of the genus *Patagifer* and was recovered from the Threskiornithidae birds in Europe, Asia, Africa, Australia, and America [29]. *P. bilobus* was the only known species found in spoonbills (genus *Platalea*) [29]. In this study, we recovered only 1 damaged specimen, which had similar characteristics to *P. bilobus*; a ribbon-like body, non-confluent vitelline fields, 4 pairs of angle spines on the ventral lappet and a deep dorsal incision on the head collar [29]. Our specimen was missing some of the tegumental spines and a section of the collar spines; however, we could confirm traces of these spines and identified the specimen as *P. bilobus* based on the morphology and host ranges. *P. minor* was added to the list as a new host of *P. bilobus* and was the first reported case of an echinostome fluke from the Threskiornithidae birds in Korea.

S. metatestis was first described by Saakova in 1952 and named *Opisthotrema metatesti*. However, in 1956, Skrjabin and Baschkirova [30] identified it as an echinostome species based on its echinostomid characteristics. They created a new genus, *Saakotrema*, for the fluke of Saakova and assigned it to the genus *Saakotrema* in the family Echinostomatidae [30]. The specimens collected in this study were identical morphologically (leaf-shaped body, posterior extension of the uterus, presence of the uroproct) and in size to *S. metatestis* (Table 2). *S. metatestis* has been recovered from piscivorous birds in Europe and Asia [31], including China [32]. This is, so far, the first report of *S. metatestis* in Far-East Asia.

Among echinostomes reported in Korea, 7 species (*A. tyosenense*, *E. japonicus*, *E. perfoliatus*, *E. cinetorchis*, *E. hortense*, *E. revolutum*, and *E. recurvatum*) were known as zoonotic parasites infecting humans [33]. There was no evidence that the 3 present species could cause human infections. However, in the case of *A. tyosenense*, it was primarily considered only as a parasite of birds [5], but 60 years later it was revealed to be a zoonotic parasite [34]. Thus, further monitoring is required in aspects of public health.

The Korean peninsula bridges China and Japan, so we can assume that helminth fauna overlaps the 2 countries. However, the investigation of parasitic fauna in Korea is not thorough enough. In China [32] and Japan, some faunistic revisions are

Table 1. Definitive hosts of echinostomes other than humans in Korea

<i>Echinostome</i> spp.	Common name of hosts	Scientific name of hosts	References
<i>Acanthoparyphium marilae</i>	Black scoter	<i>Melanitta nigra americana</i>	[5]
<i>Acanthoparyphium tyosenense</i>	Ruddy Turnstone	<i>Arenaria interpres</i>	[14]
	Great Knot	<i>Calidris tenuirostris</i>	[14]
	Black-tailed gull	<i>Larus crassirostris</i> (exp.)	[23,25]
	White-winged scoter	<i>Melanitta fusca stejnegeri</i>	[5]
	Black scoter	<i>Melanitta nigra americana</i>	[5]
<i>Echinochasmus japonicus</i>	Domestic duck	<i>Anas platyrhynchos</i> var. <i>domestica</i>	[9]
	Great egret	<i>Ardea alba modesta</i>	[15]
	Black-striped field mouse	<i>Apodemus agrarius ningpoensis</i>	[24]
	White-toothed shrew	<i>Crocidura lasiura</i>	[26]
	Cat	<i>Felis catus</i>	[13]
<i>Echinochasmus perfoliatus</i>	Cat	<i>Felis catus</i>	[8]
<i>Echinoparyphium recurvatum</i>	Mallard	<i>Anas platyrhynchos platyrhynchos</i>	[7]
	Spot-billed duck	<i>Anas poecilorhyncha zonorhyncha</i>	[7]
	Great crested grebe	<i>Podiceps cristatus</i>	[7]
	House rat	<i>Rattus norvegicus</i>	[11]
<i>Echinostoma cinetorchis</i>	Black-striped field mouse	<i>Apodemus agrarius ningpoensis</i>	[26]
	Dog	<i>Canis lupus</i>	[22]
	House rat	<i>Rattus norvegicus</i>	[11]
<i>Echinostoma gotoi</i>	Mallard	<i>Anas platyrhynchos platyrhynchos</i>	[7]
	Spot-billed duck	<i>Anas poecilorhyncha zonorhyncha</i>	[7]
<i>Echinostoma hortense</i>	Black-striped field mouse	<i>Apodemus agrarius ningpoensis</i>	[6,13]
	Dog	<i>Canis lupus</i>	[22]
	Cat	<i>Felis catus</i>	[13]
	Manchurian reed vole	<i>Microtus fortis pellceus</i>	[6]
	Mouse	<i>Mus musculus</i>	[6]
	House rat	<i>Rattus norvegicus</i>	[4,6,11,21]
	Black rat	<i>Rattus rattus alexandrinus</i>	[6]
	Mallard	<i>Anas platyrhynchos platyrhynchos</i>	[7]
	Domestic duck	<i>Anas platyrhynchos</i> var. <i>domestica</i>	[9]
Spot-billed duck	<i>Anas poecilorhyncha zonorhyncha</i>	[7]	
<i>Echinostoma revolutum</i>	Falcated teal	<i>Anas falcata</i>	[7]
	Mallard	<i>Anas platyrhynchos platyrhynchos</i>	[3,7]
	Spot-billed duck	<i>Anas poecilorhyncha zonorhyncha</i>	[7]
	Greater Scaup	<i>Aythya marila</i>	[7]
	Cat	<i>Felis catus</i>	[13]
	House rat	<i>Rattus norvegicus</i>	[11]
<i>Euparyphium murinum</i>	Black-striped field mouse	<i>Apodemus agrarius ningpoensis</i>	[6,26]
<i>Himasthla alincia</i>	Chicken	<i>Gallus gallus domesticus</i> (exp.)	[12]
<i>Himasthla kusiasigi</i>	Dunlin	<i>Calidris alpina sakhalina</i>	[16]
	Black-tailed gull	<i>Larus crassirostris</i> (exp.)	[10]
<i>Petasiger neocomense</i>	Great crested grebe	<i>Podiceps cristatus</i>	Present study
<i>Patagifer bilobus</i>	Black-faced spoonbill	<i>Platalea minor</i>	Present study
<i>Saakotrema metatestis</i>	Little egret	<i>Egretta garzetta</i>	Present study
<i>Stephanoprora</i> sp.	Cat	<i>Felis catus</i>	[13]

available, but not in Korea. In this study, we found 3 unrecorded echinostome flukes from migratory birds in Korea and presented new area records for them. It is likely that many more unrecorded parasites are present in Korea, underlining the needs for additional studies.

ACKNOWLEDGMENTS

This work was supported by a grant from the National Institute of Biological Resources (NIBR), funded by the Ministry of Environment (MOE), the Republic of Korea (NIBR No. 2013-

Table 2. Comparison of *Saakotrema metatestis* morphometrics with those of previous studies

Host	<i>Egretta garzetta</i>	<i>Egretta garzetta</i>	<i>Egretta garzetta</i>
Reference	32	33	Present study
No.	6	4	3
Body (L)	1,500-2,470	1,600-2,300 (2,000)	1,720-2,000 (1,860)
Body (W)	0.45-0.58	500-700 (600)	500-600 (550)
Head collar (L)	-	108-150 (132)	87.5-132.5 (115)
Head collar (W)	-	198-270 (220)	185.0-222.5 (203.3)
Oral sucker (L)	40-55	79-120	42.5-80.0 (66.7)
Oral sucker (W)	54-61	66-90	55.0-80.0 (68.3)
Pharynx (L)	89-105	102-168	105.0-110.0 (108.3)
Pharynx (W)	68-88	60-72	62.5-80.0 (70.0)
Ventral sucker (L)	251-282	252-300	277.5-300.0 (285.0)
Ventral sucker (W)	244-282	276-300	260.0-302.5 (280.8)
Anterior testis (L)	251-322	240-330	244.1-253.9 (240.6)
Anterior testis (W)	253-310	180-330	283.2-341.8 (306.0)
Posterior testis (L)	311-345	330-432	322.3-332.0 (325.5)
Posterior testis (W)	270-321	180-330	234.4-322.3 (266.9)
Cirrus sac (L)	-	116-126	135
Cirrus sac (W)	-	96-116	75
Ovary (L)	117-150	90-180	75.0-130.0 (108.3)
Ovary (W)	108-120	108-150	112.5-157.5 (132.5)
Egg (L)	74.5-84.3	76-81	70.0-82.5 (74.2)
Egg (W)	41.8-47.6	45-54	42.5-47.5 (45.1)

L, length; W, width.

02-001). Parasite materials used in this study were provided by the Parasite Resource Bank of Korea, National Research Resource Center (2012-0000037), the Republic of Korea.

CONFLICT OF INTEREST

We have no conflict of interest related to this work.

REFERENCES

- Toledo R, Esteban JG, Fried B. Immunology and pathology of intestinal trematodes in their definitive hosts. *Adv Parasitol* 2006; 63: 285-365.
- Yamaguti S. *Systema Helminthum*. Vol. 1, Part 1. The Digenetic Trematodes of Vertebrates. New York, USA. Interscience Publishers. 1958.
- Issiki O. On the trematode, *Echinostoma revolutum* Fröhlich, 1802 from the Korean wild duck. *Central Vet Med J (Japan)* 1934; 47: 639-647 (in Japanese).
- Park JT. A rat trematode, *Echinostoma hortense* Asada, from Korea. *Keijo J Med* 1938; 9: 283-286.
- Yamaguti S. Studies on the helminth fauna of Japan. Part 25. Trematodes of birds, IV. *Jpn J Zool* 1939; 8: 131-210.
- Seo BS, Rim HJ, Lee CW. Studies on the parasitic helminths of Korea I. Trematodes of rodents. *Korean J Parasitol* 1964; 2: 20-26.
- Chu JK, Cho YJ, Chung SB, Won BO, Yoon MB. Study on the trematode parasites of the birds in Korea. *Korean J Parasitol* 1973; 22: 70-75.
- Lee HS. A survey on helminth parasites of cats in Gyeongbug Area. *Korean J Vet Res* 1979; 19: 57-61.
- Eom KS, Rim HJ, Jang DH. A study on the parasitic helminths of domestic duck (*Anas platyrhynchos* var. *domestica* Linnaeus) in Korea. *Korean J Parasitol* 1984; 22: 215-221.
- Kim YG, Chun SK. Studies on a trematode parasitic in bivalves IV. On the metacercaria of *Himasthla kusanagi* Yamaguti, 1939 (Trematoda) found in the clam *Meretrix lusoria* Röding. *Bull Korean Fish Soc* 1984; 17: 61-67.
- Lee SH, Sohn WM, Chai JY. *Echinostoma revolutum* and *Echinoparyphium recurvatum* recovered from house rats in Yangyang-gun, Kangwon-do. *Korean J Parasitol* 1990; 28: 235-240.
- Han ET, Han KY, Chai JY. Tegumental ultrastructure of the juvenile and adult *Himasthla alincia* (Digenea: Echinostomatidae). *Korean J Parasitol* 2003; 41: 17-25.
- Sohn WM, Chai JY. Infection status with helminthes in feral cats purchased from a market in Busan, Republic of Korea. *Korean J Parasitol* 2005; 43: 93-100.
- Chung OS, Joo KH, Lee WS. *Acanthoparyphium tyosenense* infection in great knots and turnstone on the western coast of Korea. *J Ecol Field Biol* 2007; 30: 97-99.
- Ryang YS, Ahn YK, Yoon MB. Trematode infections in the small intestine of *Egretta alba modesta* in Kangwon-do. *Korean J Parasitol* 1991; 29: 227-223.

16. Chu JP, Cho YJ, Yo JC, Park SG. *Himasthla kususigi* (Trematoda, Echinostomatidae) recovered from the intestine of the dulin, *Calidris alpina sakhalina*, in Korea. *Korean J Syst Zool* 2000; 16: 125-131.
17. Yamaguti S. Synopsis of Digenetic Trematodes of Vertebrates. Tokyo, Japan. Keigaku Publishing Co. 1971, p 1074.
18. Bykhovskaya-Pavlovskaya IE. Trematodes of Birds in the USSR. Moscow-Leningrad. Izdatel'stvo Akademii Nauk SSSR. 1962, p 407.
19. Kostadinova A, Gibson DI, Biserkov V, Chipev N. Re-validation of *Echinostoma miyagawai* Ishii, 1932 (Digenea: Echinostomatidae) on the basis of the experimental completion of its life-cycle. *Syst Parasitol* 2000; 45: 81-108.
20. Kostadinova A, Gibson DI, Biserkov V, Ivanova R. A quantitative approach to the evaluation of the morphological variability of two echinostomes, *Echinostoma miyagawai* Ishii, 1932 and *E. revolutum* (Frölich, 1802), from Europe. *Syst Parasitol* 2000; 45: 1-15.
21. Seo BS, Rim HJ, Lee CW, Yoon JS. Studies on the parasitic helminths of Korea II. Parasites of the rat, *Rattus norvegicus* Exl. in Seoul, with the description of *Capillaria hepaticai* (Bancroft, 1893) Travassos (1915). *Korean J Parasitol* 1964; 2: 55-62.
22. Cho SY, Kang SY, Ryang YS. Helminthes infections in the small intestine of stray dogs in Ejungbu City, Kyunggi do, Korea. *Korean J Parasitol* 1981; 19: 55-59.
23. Kim YG. Studies on a trematode parasitized on bivalves V. On metacercaria of Echinostomatidae detected from *Macra veneriformis*, *Cyclina sinensis*, and *Solen strictus*. *Bull Korean Soc Fish Pathol* 1988; 1: 31-37.
24. Yong TS, Chung KH, Ree HI. Infection status of intestinal parasites of field rodents in Korea. *Yonsei Reports Trop Med* 1991; 22: 55-59.
25. Kim YG, Yu JE, Chung EY, Chung PR. *Acanthoparyphium tyosenense* (Digenea: Echinostomatidae): experimental confirmation of the cercaria and its complete life history in Korea. *J Parasitol* 2004; 90: 97-102.
26. Chai JY, Park JH, Jung BK, Guk SM, Kim JL, Shin EH, Klein TA, Kim HC, Chong ST, Baek LJ, Song JW. Echinostome infections in the striped-field mouse *Apodemus agarius*, and the Ussuri white-toothed shrew, *Crocidura lasiura*, caught near the demilitarized zone, Gyeonggi-do (Province), Republic of Korea. *Korean J Parasitol* 2009; 47: 311-314.
27. Faltýnková A, Gibson DI, Kostadinova A. A revision of *Petasiger* Dietz, 1909 (Digenea: Echinostomatidae) and a key to its species. *Syst Parasitol* 2008; 71: 1-40.
28. Fuhrmann O. *Petasiger neocomense* nov. sp. Une nouvelle espe'ce d'Echinostomides. *Bulletin de la Socie'te' Neucha'teloise des Sciences Naturelles (Nouvelle se'rie)*, 1927; 1: 3-6.
29. Faltýnková A, Gibson DI, Kostadinova A. A revision of *Petasiger* Dietz, 1909 (Digenea: Echinostomatidae) and a key to its species. *Syst Parasitol* 2008; 70: 159-183.
30. Skrjabin KI, Baschkirova EY. Family Echinostomatidae Dietz, 1909. In Skrjabin KI, eds, Trematodes of Animals and Man, Principles of Trematology (Vol. 12). Moscow, USSR. Izdatelstro Akademi Nauk S.S.S.R. 1956, p 501-546.
31. Sitko J. Trematodes of herons (Aves: Ciconiiformes) in the Czech Republic. *Helminthologia* 2012; 49: 33-42.
32. Chen HT. Fauna Sinica. Platyhelminthes, Trematoda, Digenea (I). Beijing, China. Science Press. 1985, p 333-497.
33. Chai JY. Echinostomes in humans (Chapter 7). In Fried B, Toledo R, eds, The Biology of Echinostomes. New York, USA. Springer. 2009, p 147-183.
34. Chai JY, Han ET, Park YK, Guk SM, Lee SH. *Acanthoparyphium tyosenense*: the discovery of human infections and identification of its source. *J Parasitol* 2001; 87: 794-800.