

Case Report

Contralateral Juxtafacet Cyst Development after the Spontaneous Resolution of a Previous Facet Cyst

Hyeun Sung Kim, M.D., Ph.D.,^{1*} Chang Il Ju, M.D., Ph.D.,² Seok Won Kim, M.D., Ph.D.,² Sung Hoon Kim, M.D., Ph.D.³

Department of Neurosurgery,¹ Heori Sarang Hospital, Daejeon, Korea

Department of Neurosurgery,² College of Medicine, Chosun University, Gwangju, Korea

Department of Rehabilitation Medicine,³ Wonju Christian Hospital, Yonsei University Wonju College of Medicine, Wonju, Korea

Juxtafacet cysts are implicated in neural compression. Thus far, it is known that surgical removal is the definitive treatment for symptomatic juxtafacet cyst because spontaneous regression is rare, and the failure rate of conservative treatment is high. We have reported a rare case of right-sided juxtafacet cyst development after the spontaneous resolution of contralateral left-sided facet cyst. The left-sided facet cyst resolved spontaneously without surgical treatment, but a juxtafacet cyst developed on the contralateral facet on the right side, as illustrated on 4-year follow-up magnetic resonance images. To the best of our knowledge, this is the first report of newly developed contralateral juxtafacet cyst after spontaneous regression. Herein, we have discussed the natural history and the management of this rare case.

Key Words : Lumbar · Facet · Cyst.

INTRODUCTION

Advances in imaging have facilitated the diagnosis of intraspinal cystic lesions in addition to lesions in the facet joint of the lumbar spine. The symptoms of lumbar juxtafacet cyst include unilateral or bilateral radiculopathy, as reported in 55–97% of the cases^{5,6,8,11,16}. The pathogenesis of facet cyst is closely related to the degenerative instability of the lumbar spine and the degenerative changes of the facet joints. The cysts are formed because of arthrotic disruption of the facet joint and are commonly associated with degeneration of joints, typically observed in mobile segments³. Good results have been reported after surgical excision of the symptomatic lesions. There are limited data, however, on nonsurgical management of these cysts, and only a few studies have reported spontaneous regression of these lesions⁷. We report a rare case of a patient who experienced spontaneous regression of symptomatic lumbar facet cyst and in whom a new contralateral juxtafacet cyst developed after 4 years.

CASE REPORT

A 58-year-old male patient was admitted to our institute for intermittent low back pain and radiating pain in the right leg. He had been treated with conservative treatment for left facet cyst at the L3–4 level 4 years ago. A plain radiographs showed mild instability on the L3–4 level (Fig. 1A, B). Magnetic resonance (MR) imaging obtained 4 years ago revealed an extradural mass consistent with a facet cyst, arising from the medial left L3–4 facet joint, that compressed the posterolateral aspect of the dural sac and showed combined stenosis (Fig. 1C, D). He refused to undergo the recommended surgical treatment. In compliance with his wishes, he was administered conservative treatment. His symptoms resolved gradually. Approximately 4 weeks after presentation, he experienced near-total resolution of his symptoms. After 4 years, he revisited the hospital without evident acute or chronic back injury. He complained of enduring low back pain and of radiating pain in the right leg. MR image of the lumbar spine showed an intraspinal extradural mass, near at the medial right L3–4 facet joint, compressing the posterolateral aspect of the dural sac, and combined stenosis, but the previous left facet cyst had regressed completely (Fig. 2). He

• Received : May 23, 2014 • Revised : July 23, 2014 • Accepted : August 13, 2014

* Move to : Department of Neurosurgery, Nanoori Hospital, Suwon, Korea

• Address for reprints : Seok Won Kim, M.D., Ph.D.

Department of Neurosurgery, College of Medicine, Chosun University, 365 Pilmun-daero, Dong-gu, Gwangju 61453, Korea

Tel : +82-62-220-3126, Fax : +82-62-227-4575, E-mail : chosunns@chosun.ac.kr

• This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.




Fig. 1. A and B : Simple dyanmic flexion and extension views show mild instability at the L3–4 level. C and D : T2-weighted magnetic resonance images showing cyst on the left L3–4 facet joint and compression of the L5 root.




Fig. 2. T2-weighted magnetic resonance images taken after 4 years show the presence of a cyst on the right L3–4 facet joint. The previous left facet cyst is absent.

had received conservative treatment, which included anti-inflammatory medication, muscle relaxants, and physical therapy. This time, however, the pain was progressive and was unresponsive to conservative treatment. We surgically excised the juxtafacet cyst and performed interbody fusion of L3–4 and percutaneous screw fixation for combined spinal stenosis (Fig. 3). The patient’s symptoms resolved immediately after surgery, and the postoperative course was uneventful. At 12 months after surgery, he complained of mild back discomfort but remained free of leg pain.

DISCUSSION

Juxtafacet cysts, also known as synovial or ganglion cysts, are rare intraspinal lesions that may elicit symptoms of disc herniation or stenosis. The term “juxtafacet cyst” was initially used by Kao et al.⁷⁾ to describe both synovial and ganglion cysts that are adjacent to facet joint or arise from or grow into the ligamentum flavum. The exact etiology of intraspinal juxtafacet cyst remains unknown¹⁴⁾. These lesions have been arised most frequently in patients aged above 50 years and have severely degenerated facet joints and instability. In the lumbar spine, jux-



Fig. 3. Simple radiographs showing interbody fusion of L3–4 with percutaneous screw fixation after surgery.

tafacet cysts appear in association with facet joint degeneration and juxtafacet cysts are thought to be caused by repetitive trauma and microinstability, thereby leading to focal weakness in the facet capsule^{3,9,15)}. The natural history of juxtafacet cysts is unknown, and there is scant literature on their conservative treatment. Surgical removal has been considered the definitive treatment for symptomatic intraspinal cysts because conservative treatment is not widely documented, and the reported success rates for conservative treatment are substantially lower than those for surgical treatment^{4,10,12,13)}. However, extensive conservative treatment should always be considered as the first therapeutic option, in case of no severe neurological deficits. Some cases of facet cysts spontaneously regress with corresponding changes in clinical symptoms. One plausible explanation for this observation is cyst rupture, which occurs in other regions of the body in conjunction with radiographically documented resolution of lesions. Another possible explanation for cyst regression is the alteration of the local forces driving the cyst formation⁴⁾. In our case, at first admission, the patient had spinal

stenosis with mild instability and left facet cyst at the same level. According to the patient's wishes, conservative treatment was administered that resolved his symptoms. After 4 years, he visited our institute again for radiating pain in the right leg. MR images showed that the left facet cyst had regressed and that a contralateral right facet cyst had developed. As conservative treatment had failed to improve his symptoms, surgical treatment was performed for removal of the facet cyst and coexistent spinal stenosis. It is believed that juxtafacet cysts grow owing to facet joint instability, which increases intra-articular pressure and induces herniation of the synovium through a focal area of weakened joint capsule^{1,3}. Symptomatic juxtafacet cysts can be treated effectively with a minimally invasive microsurgical approach. When the cysts are associated with spondylolisthesis or stenosis, they may need to be stabilized in order to minimize the risk of instability. In one study, patients with persistent low back pain and radiating pain did not experience relief despite aggressive conservative treatment, and fusion surgery was found to be necessary for alleviating their symptoms².

CONCLUSION

We report a rare case of a patient who experienced spontaneous regression of facet cyst and in whom a contralateral juxtafacet cyst developed. Accordingly, careful observation should be conducted in case of symptomatic facet cyst.

References

- Alicioglu B, Sut N : Synovial cysts of the lumbar facet joints : a retrospective magnetic resonance imaging study investigating their relation with degenerative spondylolisthesis. *Prague Med Rep* 110 : 301-309, 2009
- Deinsberger R, Kinn E, Ungersböck K : Microsurgical treatment of juxta facet cysts of the lumbar spine. *J Spinal Disord Tech* 19 : 155-160, 2006
- Doyle AJ, Merrilees M : Synovial cysts of the lumbar facet joints in a symptomatic population : prevalence on magnetic resonance imaging. *Spine (Phila Pa 1976)* 29 : 874-878, 2004
- Houten JK, Sanderson SP, Cooper PR : Spontaneous regression of symptomatic lumbar synovial cysts. Report of three cases. *J Neurosurg* 99 (2 Suppl) : 235-238, 2003
- Howington JU, Connolly ES, Voorhies RM : Intraspinal synovial cysts : 10-year experience at the Ochsner Clinic. *J Neurosurg* 91 (2 Suppl) : 193-199, 1999
- Hsu KY, Zucherman JF, Shea WJ, Jeffrey RA : Lumbar intraspinal synovial and ganglion cysts (facet cysts). Ten-year experience in evaluation and treatment. *Spine (Phila Pa 1976)* 20 : 80-89, 1995
- Kao CC, Uihlein A, Bickel WH, Soule EH : Lumbar intraspinal extradural ganglion cyst. *J Neurosurg* 29 : 168-172, 1968
- Kim SW, Lee SM : Cauda equina syndrome caused by bilateral facet cyst accompanying spinal stenosis. *J Korean Neurosurg Soc* 37 : 76-78, 2005
- Kouten JK, Sanderson SP, Cooper PR : Spontaneous regression of symptomatic lumbar synovial cysts : report of three cases. *J Neurosurg* 99 (2 Suppl) : 235-238, 2003
- Lee JC, Kang SH, Park SW : A case of spontaneous regression of lumbar synovial cyst and literature review. *Korean J Spine* 7 : 212-214, 2010
- Lyons MK, Atkinson JL, Wharen RE, Deen HG, Zimmerman RS, Lemmens SM : Surgical evaluation and management of lumbar synovial cysts : the Mayo Clinic experience. *J Neurosurg* 93 (1 Suppl) : 53-57, 2000
- Ngo T, Decina P, Hsu W : Spontaneous resolution of symptoms associated with a facet synovial cyst in an adult female - a case report. *J Can Chiropr Assoc* 57 : 87-92, 2013
- Swartz PG, Murtagh FR : Spontaneous resolution of an intraspinal synovial cyst. *AJNR Am J Neuroradiol* 24 : 1261-1263, 2003
- Silbergleit R, Gebarski SS, Brunberg JA, McGillicuddy J, Blaiivas M : Lumbar synovial cysts : correlation of myelographic, CT, MR, and pathologic findings. *AJNR Am J Neuroradiol* 11 : 777-779, 1990
- Violaris K, Karakyriou M : Presentation of a rare case of bilateral lumbar synovial cysts. *Open J Mod Neurosurg* 2 : 25-27, 2012
- Yarde WL, Arnold PM, Kepes JJ, O'Boynick PL, Wilkinson SB, Batnitzky S : Synovial cysts of the lumbar spine : diagnosis, surgical management, and pathogenesis. Report of eight cases. *Surg Neurol* 43 : 459-464; discussion 465, 1995