



# Ethics in the Intensive Care Unit

**Jae Young Moon, M.D., Ph.D. and Ju-Ock Kim, M.D., Ph.D.**

Department of Internal Medicine, Chungnam National University Hospital, Chungnam National University School of Medicine, Daejeon, Korea

The intensive care unit (ICU) is the most common place to die. Also, ethical conflicts among stakeholders occur frequently in the ICU. Thus, ICU clinicians should be competent in all aspects for ethical decision-making. Major sources of conflicts are behavioral issues, such as verbal abuse or poor communication between physicians and nurses, and end-of-life care issues including a lack of respect for the patient's autonomy. The ethical conflicts are significantly associated with the job strain and burn-out syndrome of healthcare workers, and consequently, may threaten the quality of care. To improve the quality of care, handling ethical conflicts properly is emerging as a vital and more comprehensive area. The ICU physicians themselves need to be more sensitive to behavioral conflicts and enable shared decision making in end-of-life care. At the same time, the institutions and administrators should develop their processes to find and resolve common ethical problems in their ICUs.

**Keywords:** Ethics; Behavior; Terminal Care; Intensive Care Units

## Introduction

Physicians often experience ethical dilemmas in the clinical field. The intensive care unit (ICU) is the most common place of ethical conflicts, and many respiratory physicians face with these conflicts frequently. Recently, the intensive care dedicated system was implemented, and the critical care environment is rapidly changing in Korea. In this paper, we will describe the impact of ethical dilemmas on the quality of care and the role of the ICU physicians with respect to such dilemmas.

**Address for correspondence: Ju-Ock Kim, M.D., Ph.D.**

Department of Internal Medicine, Chungnam National University School of Medicine, 266 Munhwa-ro, Jung-gu, Daejeon 301-747, Korea

**Phone:** 82-42-280-7158, **Fax:** 82-42-257-5753

**E-mail:** jokim@cnu.ac.kr

**Received:** Feb. 2, 2015

**Revised:** Feb. 17, 2015

**Accepted:** Feb. 23, 2015

© It is identical to the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>).

Copyright © 2015

The Korean Academy of Tuberculosis and Respiratory Diseases.

All rights reserved.

## Impact of Ethical Conflicts in the ICU

Today's health care environment is increasingly complex. The complexity of technology-driven modern health care, value heterogeneity, individual rights, and the number of choices according to individual values are less conducive to good decision making<sup>1</sup>. Despite considerable technologic breakthroughs in the provision of intensive care medicine, mortality in the ICU remains high<sup>2</sup> and ethical conflicts among stakeholders occur frequently in contemporary health care settings. These may have negative impacts on healthcare workers, patients, and their families, and lower the quality of intensive care.

In a previous study, over 70% of ICU workers reported perceived conflicts, which were often considered severe and were significantly associated with job strain<sup>3</sup>. Nurses perceived distressing situations more frequently than physicians did; additionally, 45% of the registered nurses surveyed reported having left or having considered leaving a position because of moral distress<sup>4</sup>. The ICU is one of the places in the hospital where family members suffer. More than two-thirds of family members visiting ICU patients have symptoms of anxiety or depression. In a French study, it was found that 75.5% of family members and 82.7% of spouses had symptoms of anxiety or depression in a multicenter study including 78 ICUs and 544 family members<sup>5</sup>.

## Major Sources of Ethical Conflicts in the ICU

The incidence of ethical conflicts is higher in the ICU compared with the general wards. In a 3-year period during which a large urban teaching hospital had a total of 181,558 admissions, the hospital ethics committee staffed 0.16% of consultations. The services with the highest incidence of ethics committee consultations were the trauma ICU (0.88%) and medical ICU (0.56%)<sup>6</sup>. In another 1-year study, 4,968 patients were admitted to the adult ICUs at Columbia University Medical Center and 168 ethical consultations (3.3%) were requested<sup>7</sup>. Although there are few reports on domestic data, according to a single center study by Park et al.<sup>8</sup>, the incidences of perceived conflicts that registered nurses reported during two different study periods were 2.1% and 0.5%, respectively.

Swetz et al.'s study<sup>9</sup> showed the common diagnoses of cases referred to consultation at the Mayo Clinic in Rochester, Minnesota, during a 10-year period, including malignancy (18%) and neurologic disease (18%), followed by cardiovascular disease (17%), multi-organ failure (11%), pulmonary disease (9%), and others. According to Romano et al.'s study<sup>7</sup>, the most common diagnosis was cardiovascular disease, followed by neurologic disease, pulmonary disease, renal disease, gastrointestinal disease, malignancy, multi-organ system failure, psychiatric disease, developmental or genetic disease, and miscellaneous.

The 1-day cross-sectional questionnaire survey from 397 ICUs in 29 countries showed that conflicts were perceived by 5,268 respondents (71.6%) and that nurse-physician conflicts were the most common (32.6%)<sup>3</sup>. The major sources of conflicts

in ICUs were behavior-related conflicts and conflicts associated with end-of-life (EOL) care. The most common conflict-causing behaviors were personal animosity, mistrust, and communication gaps. During EOL care, the main sources of perceived conflict were a lack of psychological support, absence of staff meetings, and problems with the decision-making process.

In particular, verbal abuse was one of the conflict-causing behaviors, and nurse-physician disruptive behavior was found to have a negative impact on patient safety<sup>10</sup>. As Sofield and Salmond's study reported<sup>11</sup>, 51% of respondents reported patient errors occurred as a result of such behavior. The most common forms of verbal abuse and/or disruptive behavior was condescending language. Other common types of disruptive behavior included disrespect and a failure to return phone calls<sup>12-15</sup>. In many studies, between 17% and 41.9% of health-care providers reported a specific adverse event that occurred as a result of disruptive behavior<sup>16,17</sup>. As a result, eleven recommendations have been made by The Joint Commission to help organizations implement the group's standards (Table 1)<sup>18</sup>.

## Burnout Syndrome and Moral Distress of Health Care Professionals

ICU caregivers work in a stressful environment, which can lead to burnout syndrome<sup>19</sup>. In addition, the ethical environment of their unit and ethical conflicts cause moral distress, which is believed to be associated with burnout.

It is well known that burnout can affect up to 45% of ICU nurses and physicians<sup>20,21</sup>. Burnout is a psychological syn-

**Table 1. The Joint Commission Recommendations for managing disruptive behavior**

No.	Joint Commission Recommendations
1	Educate all team members on appropriate behavior defined by the organizations code of conduct
2	Enforce the code of conduct consistently and equitably among all staff
3	Develop and implement policies that address <ul style="list-style-type: none"> <li>- "Zero tolerance" for disruptive behavior</li> <li>- Complementary policies for physicians and non-physicians</li> <li>- Non-retaliation clauses</li> <li>- Response to patients and/or families who experience or witness disruptive behavior</li> <li>- Disciplinary actions</li> </ul>
4	Develop a process for addressing disruptive behavior with input from medicine, nursing, administration, and other employees
5	Provide training in conflict resolution
6	Assess staff perceptions of disruptive behavior and threat to patient safety
7	Develop a reporting/surveillance system for identifying disruptive behavior
8	Support surveillance with tiered strategies - move toward discipline if pattern persists
9	Conduct interventions with a commitment to well-being of all staff
10	Encourage interprofessional dialogues
11	Document all attempts to address behavior

Source: The Joint Commission<sup>18</sup>. [http://www.jointcommission.org/SentinelEvents/SentinelEventAlert/sea\\_40.htm](http://www.jointcommission.org/SentinelEvents/SentinelEventAlert/sea_40.htm).

drome that occurs in response to chronic emotional and interpersonal stressors at work. It can lead to emotional instability, commitment difficulties, a feeling of failure, and an urge to leave one's job. Consequences to caregivers' health include insomnia, irritability, and depressive symptoms, which in turn can impact the quality of the care they provide<sup>20,22</sup>. Burnout in physicians tends to manifest as the provider caring less for his or her patients, and the number of medical errors increases<sup>22,23</sup>.

The Maslach Burnout Inventory, which measured the burnout of the nurses who work in Korean university hospitals, demonstrated significant emotional exhaustion and high-burnout conditions in that professional environment and profession<sup>24,25</sup>.

In Canada, it was found that internal medicine residents value the opportunity to learn from dying patients, but often lack supervision and then experience emotional distress<sup>26</sup>. The majority of respondents agreed the learning experience gained from providing EOL care was meaningful, although 48.1% felt guilty and 40.6% felt failure at least some point after a patient's death. A domestic single center study reported that 69.1% of residents encountered serious ethical dilemmas at least once a year, but the majority of these residents claimed to have resolved those dilemmas either alone or through discussions with colleagues without an attending physician's advice<sup>27</sup>.

The high rate of physician suicide is also well known; the suicide rate among male doctors is 40% higher than among men in general, whereas the rate among female doctors is 130% higher than among women in general<sup>28</sup>. Health care professionals are frequently exposed to emotional stress and exhaustion.

## Type of Ethical Problems and Education in Ethics

When we examine the behavior of the health professionals involved, it can be seen that different types of moral behavior can give rise to these problem. Types of ethical problems are presented in Table 2<sup>29</sup>.

A lack of knowledge or education in health ethics is a common cause of ethical dilemmas occurring in ICU. This manifests when the health professional does not see a situation

as being an ethical or moral problem. An example is do not resuscitate (DNR) orders on chronically ill patients. The problems arising from knowledge deficits could decrease considerably if professionals receive training or education in recognizing the associated moral problems and how to be competent in moral standards of care<sup>29</sup>. This is the reason why education in ethics is important.

Doctors' and patients' values sometimes differ, and even though medical decision making is complex, it should be approached using the principles of bioethics. Such ethical decision making training in the clinic would make patients' best interests and health professionals' values consistent with the fundamental values of our society.

## Ethical Issues of EOL Care in the ICU

During EOL care, the main cause of ethical conflict was found to be the inappropriateness of care, including a lack of psychological support and problems with the decision-making process. In Korea, most terminal cancer patients die in medical institutions. However, the quality of EOL care in Korean ICUs does not ensure the terminally ill patient's autonomy and dignity at the end of the patient's life.

One of the most important EOL issues is how health care professionals communicate the patients' prognoses to them<sup>30,31</sup>. Accurate prognostic information helps patients and physicians make sound medical decisions and care choices<sup>31,32</sup>. Yun et al.'s study<sup>31</sup> found that patients who learned their prognoses from their physicians had better outcomes, such as lower rates of emotional distress and a higher health-related quality of life (QOL), but a substantial proportion of patients did not receive such information from their physicians. Despite the fact that most patients are resistant to chemotherapy regardless of cancer type near the EOL, chemotherapy was still frequently administered in Korea<sup>33</sup>. However, a multicenter prospective cohort study of patients with cancer entering the terminal stage of their disease shows that the patients' awareness of terminal illness, use of palliative care, and admission to an ICU do not influence the survival of terminally ill patients<sup>34</sup>.

Surrogate decision-making was frequently observed among Korean cancer patients, especially when the patient's death was imminent, and for decisions related to EOL care. The proportion of patient self-determination for DNR directives was extremely rare near the EOL period, far from an ethical ideal<sup>35</sup>. Annually, thousands of terminal cancer patients still die under cardio-pulmonary resuscitation in Korean ICUs. Among nurses and physicians who work in adults ICUs in Europe, perceptions of inappropriate care have been frequently reported. The main reported reason for perceived inappropriateness of care was a mismatch between the level of care and the expected patient outcome, usually in the direction of perceived excess intensity of care<sup>36</sup>.

**Table 2. Various types of moral behavior of health professionals**

No.	Type
1	Problems caused by knowledge deficits
2	Problems caused by authority
3	Problems arising from 'groupie' moral standards
4	Problems caused by moral insensitivity
5	Problems caused by amoral behavior
6	Problems caused by differences of opinion
7	Problems caused by oppositional views

Despite all the literature, there is little discussion about the QOL of terminal cancer patients in their end-stages of life or the optimal triage of admission to an ICU. According to US Medicare database, from 1993 to 2002, an increasing proportion of patients who had been diagnosed with advanced lung cancer received ICU care near the end of their lives. Two-thirds of patients died either during or within 1 month of hospitalization<sup>37</sup>. Hwang et al.'s study<sup>38</sup> showed that ICU mortality and in-hospital mortality rates of patients with lung cancer admitted to the medical ICU were 53.6 and 61.8%, respectively; very poor values.

## Shared Decision Making

The ICU is not the adequate place for the care of terminally ill patients because it is not designed for EOL care. If the disease state of the patient is not curable, how can we determine the best treatment goals for the patient? The physician used to be the best situated to make decisions reflecting the patient's best interests. However, it is no longer appropriate to assume a concordance between physician and patient values. Under the shared decision-making model, physicians work together with patients and their families to define the patient's health-care values, beliefs, and treatment preferences. The physician not only provides information about treatments but also recommends the treatments he or she believes to be most concordant with the patient's values and goals<sup>39</sup>.

## The Role of ICU Professionals

There are many barriers to overcome ethical dilemmas in different cultures. ICU leaders or dedicated professionals have the responsibility to find and resolve the 'hidden' barriers which cause conflicts and lead to a decline in the quality of care in their respective units.

In order to do that, it is important to make a constant effort to communicate with colleagues, as well as with patients and their families. At the same time, ethics training programs for residents, critical care nurses, and fellows are necessary and should be administered in their hospital and ICU trainings.

Multiple recent studies have documented that clinical practices such as educational interventions, family conferences, and the introduction of protocols might be necessary to resolve conflicts and to improve the quality of care<sup>40</sup>.

## Conclusion

Physicians frequently confront ethical issues in clinical practice, especially in the ICU. However experience, common sense, and simply being a good person, do not guarantee that

physicians can identify or resolve ethical dilemmas<sup>41</sup>.

Good clinical decision-making is only possible when the medical decision-making and ethical decision-making are correctly balanced. Knowledge about common ethical dilemmas and training in ethical decision-making is essential. At the same time, the ICU leaders and hospital directors should prepare protocols to find and resolve the common ethical problems in their hospital.

## Conflicts of Interest

No potential conflict of interest relevant to this article was reported.

## References

1. Aulisio MP, Arnold RM, Youngner SJ. Ethics consultation from theory to practice. Baltimore: The Johns Hopkins University Press; 2003.
2. Angus DC, Barnato AE, Linde-Zwirble WT, Weissfeld LA, Watson RS, Rickert T, et al. Use of intensive care at the end of life in the United States: an epidemiologic study. *Crit Care Med* 2004;32:638-43.
3. Azoulay E, Timsit JF, Sprung CL, Soares M, Rusinova K, Lafabrie A, et al. Prevalence and factors of intensive care unit conflicts: the conflictus study. *Am J Respir Crit Care Med* 2009; 180:853-60.
4. Hamric AB, Blackhall LJ. Nurse-physician perspectives on the care of dying patients in intensive care units: collaboration, moral distress, and ethical climate. *Crit Care Med* 2007;35: 422-9.
5. Pochard F, Darmon M, Fassier T, Bollaert PE, Cheval C, Colloigner M, et al. Symptoms of anxiety and depression in family members of intensive care unit patients before discharge or death. A prospective multicenter study. *J Crit Care* 2005;20: 90-6.
6. Tapper EB, Vercler CJ, Cruze D, Sexson W. Ethics consultation at a large urban public teaching hospital. *Mayo Clin Proc* 2010;85:433-8.
7. Romano ME, Wahlander SB, Lang BH, Li G, Prager KM. Mandatory ethics consultation policy. *Mayo Clin Proc* 2009;84: 581-5.
8. Park DW, Moon JY, Ku EY, Kim SJ, Koo YM, Kim OJ, et al. Ethical issues recognized by critical care nurses in the intensive care units of a tertiary hospital during two separate periods. *J Korean Med Sci* 2015;30:495-501.
9. Swetz KM, Crowley ME, Hook C, Mueller PS. Report of 255 clinical ethics consultations and review of the literature. *Mayo Clin Proc* 2007;82:686-91.
10. Saxton R, Hines T, Enriquez M. The negative impact of nurse-physician disruptive behavior on patient safety: a review of

- the literature. *J Patient Saf* 2009;5:180-3.
11. Sofield L, Salmond SW. Workplace violence: a focus on verbal abuse and intent to leave the organization. *Orthop Nurs* 2003; 22:274-83.
  12. Rosenstein AH. Original research: nurse-physician relationships: impact on nurse satisfaction and retention. *Am J Nurs* 2002;102:26-34.
  13. Smetzer JL, Cohen MR. Intimidation: practitioners speak up about this unresolved problem. *Jt Comm J Qual Patient Saf* 2005;31:594-9.
  14. Rosenstein AH, O'Daniel M. Impact and implications of disruptive behavior in the perioperative arena. *J Am Coll Surg* 2006;203:96-105.
  15. Weber DO. Poll results: doctors' disruptive behavior disturbs physician leaders. *Physician Exec* 2004;30:6-14.
  16. Rosenstein AH, O'Daniel M. Disruptive behavior and clinical outcomes: perceptions of nurses and physicians. *Am J Nurs* 2005;105:54-64.
  17. Veltman LL. Disruptive behavior in obstetrics: a hidden threat to patient safety. *Am J Obstet Gynecol* 2007;196:587.e1-4.
  18. The Joint Commission. Behaviors that undermine a culture of safety [Internet]. The Joint Commission; [cited 2008 Nov 26]. Available from: [http://www.jointcommission.org/SentinelEvents/SentinelEventAlert/sea\\_40.htm](http://www.jointcommission.org/SentinelEvents/SentinelEventAlert/sea_40.htm).
  19. Merlani P, Verdon M, Businger A, Domenighetti G, Pargger H, Ricou B, et al. Burnout in ICU caregivers: a multicenter study of factors associated to centers. *Am J Respir Crit Care Med* 2011; 184:1140-6.
  20. Poncet MC, Toullic P, Papazian L, Kentish-Barnes N, Timsit JF, Pochard F, et al. Burnout syndrome in critical care nursing staff. *Am J Respir Crit Care Med* 2007;175:698-704.
  21. Embriaco N, Azoulay E, Barrau K, Kentish N, Pochard F, Loundou A, et al. High level of burnout in intensivists: prevalence and associated factors. *Am J Respir Crit Care Med* 2007;175: 686-92.
  22. Shanafelt TD, Bradley KA, Wipf JE, Back AL. Burnout and self-reported patient care in an internal medicine residency program. *Ann Intern Med* 2002;136:358-67.
  23. Fahrenkopf AM, Sectish TC, Barger LK, Sharek PJ, Lewin D, Chiang VW, et al. Rates of medication errors among depressed and burnt out residents: prospective cohort study. *BMJ* 2008;336:488-91.
  24. Kang JH, Kim CW. Evaluating applicability of Maslach burnout Inventory among university hospitals nurses. *Korean J Adult Nurs* 2012;24:31-7.
  25. Poghosyan L, Aiken LH, Sloane DM. Factor structure of the Maslach burnout inventory: an analysis of data from large scale cross-sectional surveys of nurses from eight countries. *Int J Nurs Stud* 2009;46:894-902.
  26. Schroder C, Heyland D, Jiang X, Rocker G, Dodek P; Canadian Researchers at the End of Life Network. Educating medical residents in end-of-life care: insights from a multicenter survey. *J Palliat Med* 2009;12:459-70.
  27. Park SH, Koh YS. Attitudes toward medical ethics among resident physicians in one Korean University Hospital. *Korean J Med Ethics Educ* 2007;10:109-16.
  28. Schernhammer ES, Colditz GA. Suicide rates among physicians: a quantitative and gender assessment (meta-analysis). *Am J Psychiatry* 2004;161:2295-302.
  29. Hawley G. Ethics in clinical practice: an interprofessional approach. Edinburgh: Pearson Education; 2007. p. 45-9.
  30. Steinhäuser KE, Christakis NA, Clipp EC, McNeilly M, McIntyre L, Tulsky JA. Factors considered important at the end of life by patients, family, physicians, and other care providers. *JAMA* 2000;284:2476-82.
  31. Yun YH, Kwon YC, Lee MK, Lee WJ, Jung KH, Do YR, et al. Experiences and attitudes of patients with terminal cancer and their family caregivers toward the disclosure of terminal illness. *J Clin Oncol* 2010;28:1950-7.
  32. Back AL, Arnold RM. Discussing prognosis: "how much do you want to know?" talking to patients who are prepared for explicit information. *J Clin Oncol* 2006;24:4209-13.
  33. Yun YH, Kwak M, Park SM, Kim S, Choi JS, Lim HY, et al. Chemotherapy use and associated factors among cancer patients near the end of life. *Oncology* 2007;72:164-71.
  34. Yun YH, Lee MK, Kim SY, Lee WJ, Jung KH, Do YR, et al. Impact of awareness of terminal illness and use of palliative care or intensive care unit on the survival of terminally ill patients with cancer: prospective cohort study. *J Clin Oncol* 2011;29: 2474-80.
  35. Lee JK, Keam B, An AR, Kim TM, Lee SH, Kim DW, et al. Surrogate decision-making in Korean patients with advanced cancer: a longitudinal study. *Support Care Cancer* 2013;21: 183-90.
  36. Piers RD, Azoulay E, Ricou B, Dekeyser Ganz F, Decruyenaere J, Max A, et al. Perceptions of appropriateness of care among European and Israeli intensive care unit nurses and physicians. *JAMA* 2011;306:2694-703.
  37. Sharma G, Freeman J, Zhang D, Goodwin JS. Trends in end-of-life ICU use among older adults with advanced lung cancer. *Chest* 2008;133:72-8.
  38. Hwang KE, Seol CH, Hwang YR, Jo HG, Park SH, Yoon KH, et al. The prognosis of patients with lung cancer admitted to the medical intensive care unit. *Asia Pac J Clin Oncol* 2013 Dec 2 [Epub]. <http://dx.doi.org/10.1111/ajco.12157>.
  39. Thompson DR, Kaufman D. Critical care ethics: a practice guide. 3rd ed. Mount Prospect: Society of Critical Care Medicine; 2014.
  40. Truog RD, Campbell ML, Curtis JR, Haas CE, Luce JM, Rubenfeld GD, et al. Recommendations for end-of-life care in the intensive care unit: a consensus statement by the American College [corrected] of Critical Care Medicine. *Crit Care Med* 2008;36:953-63.
  41. Longo DL, Fauci A, Kasper D, Hauser S, Jameson J, Loscalzo J. Harrison's principles of internal medicine. 18th ed. New York: McGrawHill Medical; 2012.