

Article

Isolation of secondary metabolites from an Arctic bacterium, *Pseudomonas aeruginosa* and their antimicrobial activities

Ui Joung Youn*, Min Ju Kim, Se Jong Han, and Jung Han Yim

Division of Life Sciences, Korea Polar Research Institute, KIOST, Incheon 21990, Republic of Korea

북극유래 박테리아, *Pseudomonas aeruginosa*로 부터 대사산물들의 분리 및 항진균 활성

윤익중* · 김민주 · 한세종 · 임정한

극지연구소 생명과학 연구부

(Received September 27, 2016; Revised November 1, 2016; Accepted November 3, 2016)

ABSTRACT: Chemical study of an Arctic bacterium, *Pseudomonas aeruginosa* (Pseudomonadaceae) led to the isolation of two diketopiperazines 1 and 2, two phenazine alkaloids 3 and 4, and an indole carbaldehyde 5, along with a benzoic acid derivative 6. The structures of the compounds were confirmed by 1D and 2D NMR, and MS experiments, as well as by comparison of their data with published values. Among the isolates, compounds 5 and 6 were isolated for the first time from *P. aeruginosa* of the seawater of Arctic Chuckchi Sea. Antimicrobial activities of compounds 1–6 against a *Staphylococcus aureus* and *Candida albicans* were evaluated.

Key words: *Pseudomonas aeruginosa*, antimicrobial activity, diketopiperazine, phenazine, Pseudomonadaceae

Pseudomonas is a diverse genus of Gammaproteobacteria with more than 60 species exhibiting varied life styles in a wide range of environments, including soil, water, plant surfaces, and animals (Moore *et al.*, 2006). The genus *Pseudomonas* has had an important role in the biotechnology and pharmaceutical industries due to its metabolic versatility (Gross and Loper, 2009). The vast metabolic diversity of *Pseudomonas* spp. is reflected on the production of a wide variety of bioactive molecules such as mupirocin, fosfadecin, fosfocytosin, and xantholysins (Fuller *et al.*, 1971; Gross and Loper, 2009; Buckingham, 2011; Li *et al.*, 2013). In particular, *P. aeruginosa* produces a number of toxic substances that have been implicated towards its pathogenicity (Olgerts and Arligues, 1974; David *et al.*, 1986; Robert *et al.*, 1987). It shows antibacterial, antiarchaeal, antialgal and antifungal activity due to production of various compounds (Hassan and Irwin, 1980;

Baron and Rowe, 1981; Dakhama *et al.*, 1993; Hernandez *et al.*, 2004; Rane *et al.*, 2007). The compounds like phenazine serve as electron shuttles in reduction and solubilization of Fe(III) (Hernandez *et al.*, 2004), and function as antibiotics (Baron and Rowe, 1981). A previous study reported secondary metabolites, such as diketopiperazines and phenazine derivatives from the Antarctic strain, *P. aeruginosa*, along with their antibacterial activity (Jayatilake *et al.*, 1996).

In the course of an ongoing chemical and biochemical studies on the Arctic microorganism, a bacterial strain, *P. aeruginosa* B15H6 displayed potent antimicrobial activity. Although previous research mentioned about metabolites from an Antarctic sponge-associated bacterium, *P. aeruginosa*, this study aims to find other active compounds from an Arctic bacterium, *P. aeruginosa* B15H6. Here, we report the chemical structure elucidation and bioactivity of compounds 1–6 isolated from the Arctic bacterium, *P. aeruginosa* B15H6.

*For correspondence. E-mail: ujyoun@kopri.re.kr;
Tel.: +82-32-760-5562; Fax: +82-32-760-5509

Materials and Methods

Isolation and identification of the bacterium

The bacterial strain (B15H6) was isolated from the seawater of Chukchi Sea in 2012. Seawater (50 μ l) was swabbed in freshly prepared SZB medium (glucose 20 g/L, peptone 5 g/L, yeast extract 1 g/L, FePO₄ 10 mg/L, and sea water 750 ml/L) and incubated at 25°C for 4 days. Among several selected colonies, a unique bacterial colony with a faint creamy color, and a round thin surface was isolated. The bacterium was identified as *P. aeruginosa* B15H6 by phylogenetic analysis of the 16S rDNA sequence (Baek *et al.*, 2015). Stock cultures of the *P. aeruginosa* B15H6 used in this study are preserved at -70°C in KOPRI institute.

Extraction and isolation of the metabolites

The bacterial strain B15H6 was cultivated in 20 L Zobell medium enriched with 18 g/L peptone for 4 days using a 30-L fermenter (Biotron Co.). Cells were removed from the cultures by centrifugation and filtration through a 0.45- μ m filter, and the filtrate (20 L) was passed through an Amberlite XAD-20 resin (40 \times 8 cm), using a gradient solvent system of H₂O-MeOH (100:0 to 0:100) to give crude extracts (4.5 g). The extracts were diluted with H₂O (1 L) and partitioned using EtOAc (0.5 L \times 3 times) to yield EtOAc extracts (1.5 g). A Thin-layer chromatography (TLC) experiment was performed using a solvent mixture, CHCl₃-MeOH (9:1), to establish isolation method by silica gel column chromatography. The EtOAc extracts (1.4 g) were separated by column chromatography over silica gel (Column Chromatography, CC; ϕ 10 \times 90 cm; 230-400 mesh, 2 kg) using a gradient solvent system of CHCl₃-MeOH (99:1 to 50:50), to afford 20 fractions (E1-E20). A fraction combined from E4 to E8 (300 mg), showing similar TLC patterns was subjected to silica gel separation (CC; ϕ 3 cm; 230-400 mesh, 500 g), with CHCl₃-MeOH (100:0 to 70:30) as the solvent system, yielding ten subfractions (E4S1 to E4S10). Subfraction E4S5 (70 mg) was chromatographed over a Sephadex LH-20 column and eluted with H₂O-MeOH solvents (90:10 to 100% MeOH), to afford five subfractions (E5S5L1 to E5S5L5). Compounds **1** (10.0 mg) and **2** (8.0 mg) were recrystallized using a pure chloroform (CHCl₃), from subfractions

E5S5L2 and E5S5L4, respectively. Fraction E10 (100 mg) was chromatographed on a Sephadex LH-20 gel (300 g) column, and eluted with an H₂O-MeOH (90:10 to 0:100) solvent system, to afford five subfractions (E10L1 to E10L5). Subfraction E10L2 was purified by semi-preparative HPLC on a RP-18 column, using MeOH-H₂O mixtures (60:40 to 0:100), to yield **3** (6.0 mg, *t_R* 85 min) and **4** (5.5 mg, *t_R* 87 min). Fraction E12 was subjected to semi-preparative HPLC [MeOH/H₂O (50:50 to 0:100)] to yield **6** (2.5 mg, *t_R* 75 min) and **5** (1.5 mg, *t_R* 77 min).

Phenazine-1-carboxylic Acid (1) – Yellow needle; UV (MeOH) λ_{\max} (log *e*) 365 (3.7) nm; IR ν_{\max} (KBr) 3410, 1625 cm⁻¹; LCMS *m/z* 225 [M + H]⁺; ¹H-NMR (600 MHz, CD₃OD) δ H 8.98 (1H, dd, *J* = 7.2, 1.6 Hz, H-2), 8.53 (1H, dd, *J* = 8.8, 1.6 Hz, H-4), 8.33 (1H, dd, *J* = 8.8, 1.6 Hz, H-9), 8.28 (1H, dd, *J* = 8.0, 2.0 Hz, H-6), 8.04 (1H, overlapping signal, H-3), 8.02 (1H, m, H-7), 8.00 (1H, m, H-8); ¹³C-NMR (150 MHz, CD₃OD) δ C 166.0 (C=O), 125.0 (C-1), 137.5 (C-2), 130.3 (C-3), 135.2 (C-4), 143.5 (C-4a), 139.9 (C-5a), 128.0 (C-6), 133.3 (C-7), 131.8 (C-8), 130.1 (C-9), 144.2 (C-9a), 140.1 (C-10a).

Phenazine-1-carboxamide (2) – Pale yellow needle; UV (MeOH) λ_{\max} (log *e*) 365 (3.9) nm; LCMS *m/z* 224 [M + H]⁺; ¹H-NMR (600 MHz, CD₃OD) δ H 9.01 (1H, d, *J* = 7.2 Hz, H-2), 8.42 (1H, d, *J* = 8.4 Hz, H-4), 8.28 (1H, dd, *J* = 7.2, 2.4 Hz, H-9), 8.22 (1H, dd, *J* = 7.2, 2.4 Hz, H-6), 7.97 (1H, t, *J* = 7.2 Hz, H-3), 7.92 (1H, m, H-7), 7.92 (1H, m, H-8); ¹³C-NMR (150 MHz, CD₃OD) δ C 166.8 (C=O), 129.0 (C-1), 136.1 (C-2), 130.1 (C-3), 134.5 (C-4), 143.6 (C-4a), 141.6 (C-5a), 129.3 (C-6), 131.9 (C-7), 131.3 (C-8), 129.9 (C-9), 143.3 (C-9a), 140.9 (C-10a).

cyclo-(L-Prolyl-L-Phenylalanyl) (3) – Colorless amorphous powder; $[\alpha]_D^{20}$ -110.5 (*c* 0.2, MeOH); UV (MeOH) λ_{\max} (log *e*) 254 (3.5) nm; IR ν_{\max} (KBr) 3315, 1675, 1420 cm⁻¹; LCMS *m/z* 245 [M + H]⁺; ¹H-NMR (600 MHz, CD₃OD) δ H 7.21-7.29 (5H, m, H-1' to H-6'), 4.44 (1H, br t, *J* = 5.4 Hz, H-9), 4.06 (1H, dd, *J* = 10.8, 6.6 Hz, H-6), 3.53 (1H, ddd, *J* = 20.4, 12.0, 8.4 Hz, H-3b), 3.36 (1H, dddd, *J* = 18.6, 12.0, 7.2, 2.0 Hz, H-3a), 3.16 (2H, m, H-10), 2.08 (1H, m, H-5a), 1.80 (2H, m, H-4), 1.20 (1H, m, H-5b); ¹³C-NMR (150 MHz, CD₃OD) δ C 167.0 (C-1), 46.0 (C-3), 22.9 (C-4), 29.5 (C-5), 60.2 (C-6), 171.0 (C-7), 57.8 (C-9), 38.3 (C-10), 137.4 (C-1'), 129.5 (C-2' & C-6'), 131.2 (C-3' & C-5'), 128.2 (C-4').

cyclo-(L-Prolyl-L-Leucine) (4) – Colorless amorphous powder; $[\alpha]_D^{20}$ –120.3 (*c* 0.2, MeOH); UV (MeOH) λ_{\max} (log *e*) 225 (3.8) nm; LCMS *m/z* 211 [M + H]⁺; ¹H-NMR (600 MHz, CD₃OD) δ H 4.26 (1H, t, *J* = 7.2 Hz, H-6), 4.13 (1H, m, H-9), 3.50 (1H, dd, *J* = 13.2, 12.0 Hz, H-3), 2.29 (1H, m, H-5a), 2.04 (1H, m, H-5b), 2.01 (1H, m, H-4a), 1.94 (1H, m, H-4b), 1.93 (1H, m, H-11), 1.89 (1H, m, H-10a), 1.50 (1H, m, H-10b), 0.96 (1H, d, *J* = 6.6 Hz, H-12), 0.95 (1H, d, *J* = 6.0 Hz, H-13); ¹³C-NMR (150 MHz, CD₃OD) δ C 169.0 (C-1), 46.5 (C-3), 23.8 (C-4), 29.2 (C-5), 60.4 (C-6), 172.9 (C-7), 54.7 (C-9), 39.5 (C-10), 25.9 (C-11), 23.4 (C-12), 22.3 (C-13).

1H-Indole-3-carbaldehyde (5) – White amorphous powder, LCMS *m/z* 146 [M + H]⁺; ¹H-NMR (600 MHz, CD₃OD) δ H 10.07 (1H, s, CHO), 8.32 (1H, br d, *J* = 6.0 Hz, H-4), 7.86 (1H, d, *J* = 3.0 Hz, H-2), 7.45 (1H, br d, *J* = 6.6 Hz, H-7), 7.34 (2H, m, H-5/H6); ¹³C-NMR (150 MHz, CD₃OD) δ C 185.1 (CHO), 135.1 (C-2), 137.0 (C-1a), 124.4 (C-6), 124.0 (C-3a), 123.0 (C-5), 122.0 (C-4), 117.0 (C-3), 111.4 (C-7).

p-Hydroxybenzaldehyde (6) – White amorphous powder, LCMS *m/z* 123 [M + H]⁺; ¹H-NMR (600 MHz, CD₃OD) δ H 7.81 (2H, d, *J* = 8.4 Hz, H-2 & H-6), 6.95 (2H, d, *J* = 8.4 Hz, H-3 & H-5); ¹³C-NMR (150 MHz, CD₃OD) δ C 190.7 (CHO), 130.1 (C-1), 132.3 (C-2 & C-6), 115.8 (C-3 & C-5), 161.0 (C-4).

General experimental procedures

Optical rotations were measured on a Rudolph Research Autopol IV multi wavelength polarimeter. UV spectra were recorded on a Shimadzu PharmaSpec-1700 UV-visible spectrophotometer. IR spectra were measured on a Bruker Tensor-27 spectrophotometer. 1D and 2D NMR spectra were recorded on a Bruker AVANCE (600 MHz) spectrometer. Mass spectra and high-resolution MS spectra were performed with a BioTOF II ESI mass spectrometer. Thin-layer chromatography (TLC) was performed on silica gel 60 F₂₅₄ (0.25 mm, Merck). Silica gel (230–400 mesh, Merck) and RP-18 (YMC·GEL ODS-A, 12 nm, S-150 μ m) were used for column chromatography. Semi-preparative HPLC was conducted on YL9100 HPLC system (Young Lin) equipped with a UV/Vis detector using an Alltech reversed-phase YMC-Pak C-18 column (10 μ m, 20 \times 250 mm) with a flow rate of 2 ml/min.

Antimicrobial assays

The antimicrobial activities were tested with *Staphylococcus aureus* KCTC 3881 and *Candida albicans* KCTC 27242 (Korean Collection for Type Cultures) in 96-well-plate. Cell culture (95 μ l) was distributed in each well, and compounds dissolved in DMSO were added up to final concentrations (0.5, 1, 2, 5, 10, 20, and 50 μ M). Total culture (0.1 ml) was incubated at 25°C for 16 h. Cell inhibition was measured at 600 nm using Multiskan™ GO Microplate Spectrophotometer (Thermo Scientific). The IC₅₀ value was calculated using exponential trend line in Excel (Microsoft). Kanamycin and nystatin were used as positive controls against the bacterium and yeast, respectively.

Results and Discussion

Repeated chromatography of the culture broth, *P. aeruginosa* on silica gel and YMC-pack C₁₈ columns led to the isolation of six compounds (1–6) (Fig. 1).

Compound 1 was obtained as a yellow needle and its molecular weight was evaluated as *m/z* 225 [M + H]⁺, by the positive LCMS spectrometry. The UV spectrum showed absorption maxima at 365 nm. The IR spectrum displayed characteristic absorption bands at 3410 and 1625 cm⁻¹, corresponding to the hydroxy group(s) and conjugated C=O group(s), respectively. The ¹³C NMR spectrum revealed 13 carbon signals, including an ester carbonyl carbon at δ C 166.0 (COOH), four N-atom substituted aromatic quaternary carbons at δ C 144.2 (C-9a)/143.5 (C-4a)/140.1 (C-10a)/139.9 (C-5a), and an aromatic quaternary

Table 1. Antimicrobial activities of compounds 1–6

Compounds	<i>S. aureus</i>	<i>C. albicans</i>
	IC ₅₀ (μ g/ml)	IC ₅₀ (μ g/ml)
1	7.17	20.03
2	>50 ^a	44.85
3	>50	>50
4	>50	>50
5	13.01	>50
6	>50	>50
Kanamycin ^b	2.84	
Nystatin ^c		1.33

^a >50, considered to be inactive

^{b,c} positive controls for antibacterial and antifungal assays, respectively.

at δ_C 125.0 (C-1), indicating a characteristic diketopiperazine molecule (Jayatilake *et al.*, 1996). The 1H and 1H - 1H COSY NMR spectra showed three mutually coupled aromatic proton signals at δ_H 8.98 (1H, dd, $J = 7.2, 1.6$ Hz, H-2)/8.53 (1H, dd, $J = 8.8, 1.6$ Hz, H-4)/8.04 (1H, overlapped, H-3), indicating an 1,2,3-trisubstituted benzene moiety (Fig. 2), and additionally displayed four mutually coupled aromatic proton signals at δ_H 8.28 (1H, dd, $J = 8.0, 0.8$ Hz, H-6)/ δ_H 8.02 (1H, overlapped, H-7)/ δ_H 8.00 (1H, overlapped, H-8)/ δ_H 8.33 (1H, dd, $J = 8.8, 1.6$ Hz, H-9), corresponding to an 1,2-disubstituted benzene moiety. The HMBC spectrum confirmed the position of the carboxylic acid group at C-1 position of 1,2,3-trisubstituted benzene moiety, based on three-bond correlation of H-2 with C-10a and the carbonyl carbon (COOH). The other supporting HMBC correlations were displayed in Fig. 2. Accordingly, compound **1** was assigned as phenazine-1-carboxylic acid, by comparison of their physicochemical data with literature values (Jayatilake *et al.*, 1996).

Compound **2** was obtained as a pale yellow needle. The




Fig. 1. Structures of compounds **1-6** isolated from *P. aeruginosa*.




Fig. 2. Important HMBC and COSY correlations of **1** and **3**.

LC-MS spectrum of **2** gave a molecular ion $[M + H]^+$ at m/z 224. The NMR spectra of **2** were quite similar to those of **1**, except for a downfield shifted aromatic quaternary carbon at δ_C 129.0 (C-1). In addition, the even-numbered mass of **2** indicated that a NH_2 group was attached at carboxylic carbon instead of hydroxy group (-OH) in the carboxylic acid moiety of **1**. Thus, compound **2** was identified as phenazine-1-carboxamide (Jayatilake *et al.*, 1996).

Compound **3** was obtained as a colorless amorphous powder. The LCMS spectrum showed a molecular ion peak at m/z 245 $[M + H]^+$. The 1H NMR spectrum of **3** displayed five mutually coupled aromatic proton signals from δ_H 7.21 to δ_H 7.29, indicating 1-substituted benzene moiety. The diketopiperazine skeleton was apparent from the characteristic chemical shifts in 1H - and ^{13}C -NMR spectra; two amides at δ_C 171.0 (C-1) and 167.0 (C-7) and two N-C residues at [δ_H 4.06 (1H, dd, $J = 10.8, 6.6$ Hz)/ δ_C 60.2 (C-6) δ_H 4.44 (1H, br t, $J = 5.4$ Hz)/ δ_C 57.8 (C-9)]. Furthermore, the NMR and HSQC spectra of **3** revealed three methylene groups at [δ_H 3.53 (1H, ddd, $J = 20.4, 12.0, 8.4$ Hz), 3.36 (1H, dddd, $J = 18.6, 12.0, 7.2, 2.0$ Hz)/ δ_C 46.0 (C-3), δ_H 1.80 (2H, m)/ δ_C 22.9 (C-4), and δ_H 2.08, 1.20 (2H, m)/ δ_C 29.5 (C-5)], all of which confirmed to be closely connected to each other through the HMBC and COSY analyses (Fig. 2). In addition, NMR spectra showed remaining signals for a methylene group at δ_H 3.16 (2H, m)/ δ_C 38.3 (C-10), displayed two- and three-bond HMBC correlations with C-1/C-2' (Fig. 2). The relative stereo chemistry of asymmetric carbons was determined as 6*R* and 9*R*, by comparison of the optical rotation, $[\alpha]_D^{20} - 110.5$ (c 0.2, MeOH) of **3** with that of the literature value (Adamczeski *et al.*, 1995). Accordingly, the structure of **3** was identified as *cyclo*-(L-Prolyl-L-Phenylalanyl) (Adamczeski *et al.*, 1995).

Compound **4** was obtained as a colorless amorphous powder. The 1H and ^{13}C NMR spectra of **4** showed similar patterns compare to those of **3**. However, **4** revealed additional signals for a methine group at δ_H 1.93 (1H, m)/ δ_C 25.9 (C-11) and two symmetrical methyl groups at [δ_H 0.96 (1H, d, $J = 6.6$ Hz)/ δ_C 23.4 (C-12), 0.95 (1H, d, $J = 6.0$ Hz)/ δ_C 22.3 (C-13)] in NMR spectra, indicating an isopropyl group, confirmed the attachment to C-10 position, instead of the benzyl group in **3**. The relative stereo chemistry of **4** was confirmed to be the same with that of **3**. On the basis of the above evidence, **4** was identified as *cyclo*-

(L-Prolyl-L-Leucine), by comparison of their physicochemical data with published values (Adamczeski *et al.*, 1995).

Compound **5** was obtained as a white amorphous powder. The LCMS spectrum showed a molecular ion peak at m/z 146 $[M + H]^+$. The NMR and HSQC spectra of **5** displayed signals for four mutually coupled aromatic signals at [δ H 8.32 (1H, br d, $J=6.0$ Hz)/ δ C 122.0 (C-4), 7.86 (1H, d, $J=3.0$ Hz)/ δ C 135.1 (C-2), 7.45 (1H, br d, $J=6.6$ Hz)/ δ C 111.4 (C-7), δ H 7.34 (1H, m)/ δ C 123.0 (C-5), and δ H 7.34 (1H, m)/ δ C 124.4 (C-6)], a tri-substituted olefinic group at [δ H 7.86 (1H, d, $J=3.0$ Hz)/ δ C (138.0, C-2) and δ C (117.0, C-3)], an aromatic quaternary carbon at δ C 124.0 (C-3a), and a downfield shifted aromatic quaternary carbon at δ C 137.0 (C-1a) due to a nitrogen atom attachment, implying the presence of an indole skeleton. Furthermore, the 1 H and 13 C NMR spectra revealed downfield shifted signals at δ H 10.07 (1H, s)/ δ C 185.1 (CHO), indicating an aldehyde group. The HMBC spectrum displayed two- to three-bonds correlations from H-2 to the aldehyde carbon (CHO)/C-3a and from H-4 to C-3/C-1a/C-3a (Fig. 2), confirmed the identification of compound **5** as 1*H*-indole-3-carbaldehyde (Wang *et al.*, 2013).

The 1 H and 13 C NMR spectra of **6** indicated a benzoic acid derivative, which were in good agreement with those of *p*-hydroxybenzaldehyde (**6**) (Kim *et al.*, 2003). To the best of our knowledge, compounds **5** and **6** were isolated for the first time from *P. aeruginosa* of the seawater of Arctic Chuckchi Sea.

The EtOAc-soluble extract from the culture broth of *P. aeruginosa* B15H6 exhibited potent antimicrobial activity against *Candida albicans* and *Staphylococcus aureus*, respectively (data not shown).

The secondary metabolites, **1-6** isolated from the extract were also evaluated for their antimicrobial activity. Among the isolates, compound **1** displayed potent antibacterial activity with an IC_{50} value of 7.17 μ M against the Gram-positive bacteria, *S. aureus*, and also showed moderate antiyeast activity with an IC_{50} value of 20.03 μ M against *C. albicans*; the indol alkaloid **5** exhibited moderate antibacterial activity (IC_{50} value, 13.01 μ M) against *S. aureus*. Meanwhile the phenazine amide **2** showed only weak inhibitory activity (IC_{50} value, 44.85 μ M) against *C. albicans*. Other remaining compounds were inactive. In study of antibacterial activities for phenazine alkaloids, all compounds of the phenazine class have been reported to have antibacterial

properties (Dahiya *et al.*, 1988; Dakhama *et al.*, 1993; Jayatilake *et al.*, 1996). These results suggest the phenazine carboxylic acid **1** could possess antimicrobial activity.

적 요

북극 유래 박테리아인 *Pseudomonas aeruginosa* 균주의 대사산물에 대한 화학적 연구는 벤조산 유도체 **6**번을 포함하여, 두 개의 diketopiperazine **1**과 **2**, 두 개의 phenazine alkaloid **3**과 **4**, indole carbaldehyde **5**번을 분리하였다. 화합물들의 구조는 1D 과 2D NMR, 및 MS 기법들과 기존 보고된 문헌 값과의 비교에 의하여 동정되었다. 분리된 화합물들 중 **5**번과 **6**번은 북극 척지해 해수의 *P. aeruginosa*로부터 처음으로 보고되었다. 대사산물들 **1-6**의 항균 활성은 *Staphylococcus aureus*과 *Candida albicans*에 대하여 측정하였다.

Acknowledgements

This work was supported by a grant to the Korea Polar Research Institute (KOPRI), under a project PE16350. We thank H. S. Shin, National center for inter University Research Facilities, Seoul National University, for the provision of the Mass Spectrometry Facility used in this study.

References

- Adamczeski, M., Reed, A.R., and Crews, P. 1995. New and known diketopiperazines from the caribbean sponge, *Calyx cf. podatypa*. *J. Nat. Prod.* **58**, 201-208.
- Back, K., Lee, Y.M., Hwang, C.Y., Park, H., Jung, Y.J., Kim, M.K., Hong, S.G., Kim, J.H., and Lee, H.K. 2015. *Psychroserpens jangbogonensis* sp. nov., a psychrophilic bacterium isolated from Antarctic marine sediment. *Int. J. Syst. Evol. Microbiol.* **65**, 183-188.
- Baron, S.S. and Rowe, J.J. 1981. Antibiotic action of pyocyanin. *Antimicrob. Agents Chemother.* **20**, 814-820.
- Buckingham, J. 2011. Dictionary of Natural Products on DVD, CRC Press, BocaRaton, FL, USA.
- Dahiya, J.S., Woods, D.L., and Tewari, J.P. 1988. Control of *Rhizoctonia solani*, causal agent of brown girdling root rot of rapeseed, by *Pseudomonas fluorescens*. *Bot. Bull. Acad. Sin.* **29**, 135-142.
- Dakhama, A., de la Noue, J., and Lavoie, M.C. 1993. Isolation and

- identification of anti-fungal substances produced by *Pseudomonas aeruginosa*. *J. Appl. Phycol.* **5**, 297–306.
- David, W., John, M., Robert, W., Peter, C., and Graham, W.T.** 1986. Purification and structural analysis of pyocyanin and 1-hydroxyphenazine. *Eur. J. Biochem.* **159**, 309–313.
- Fuller, A.T., Mellows, G., Woolford, M., Banks, G.T., Barrow, K.D., and Chain, E.D.** 1971. Pseudomonic acid: an antibiotic produced by *Pseudomonas fluorescens*. *Nature* **234**, 416–417.
- Gross, H. and Loper, J.E.** 2009. Genomics of secondary metabolite production by *Pseudomonas* spp. *Nat. Prod. Rep.* **26**, 1408–1446.
- Hassan, H.M. and Irwin, F.** 1980. Mechanism of the antibiotic action of pyocyanine. *J. Bacteriol.* **141**, 156–163.
- Hernandez, M.E., Kappler, A., and Newman, D.K.** 2004. Phenazines and other redox-active antibiotics promote microbial mineral reduction. *Appl. Environ. Microbiol.* **70**, 921–928.
- Jayatilake, G.S., Thornton, M.P., Leonard, A.C., Grimwade, J.E., and Baker, B.J.** 1996. Metabolites from an Antarctic sponge-associated bacterium, *Pseudomonas aeruginosa*. *J. Nat. Prod.* **59**, 293–296.
- Kim, H., Ralph, J., Lu, F., Ralph, S.A., Boudet, A.M., MacKay, J.J., Sederoff, R.R., Ito, T., Kawai, S., Ohashi, H., et al.** 2003. NMR analysis of lignins in CAD-deficient plants. Part 1. Incorporation of hydroxycinnamaldehydes and hydroxybenzaldehydes into lignins. *Org. Biomol. Chem.* **1**, 268–281.
- Li, W., Rokni-Zadeh, H., De Vleeschouwer, M., Ghequire, M.G., Sinnaeve, D., Xie, G.L., Rozenski, J., Madder, A., Martins, J.C., and De Mot, R.** 2013. The antimicrobial compound xantholysin defines a new group of *Pseudomonas* cyclic lipopeptides. *PLoS One* **8**, e62946.
- Moore, E.R., Tindall, B.J., Dos Santos, V.A.M., Pieper, D.H., Ramos, J.L., and Palleroni, N.J.** 2006. *The Prokaryotes*, pp. 646–703. Springer, New York, USA.
- Olgerts, R.P. and Artigues, H.S.** 1974. *Pseudomonas aeruginosa* exotoxin in mice: localization and effects on protein synthesis. *Infect. Immun.* **9**, 540–546.
- Rane, M.R., Sarode, P.D., Chaudhari, B.L., and Chincholkar, S.B.** 2007. Detection, isolation identification of phenazine-1-carboxylic acid produced by biocontrol strains of *Pseudomonas aeruginosa*. *J. Sci. Ind. Res.* **66**, 627–631.
- Robert, W., Tyrone, P., Graham, T., David, W., John, M., David, S., David, R., and Peter, C.** 1987. Pyocyanin and 1-hydroxyphenazine produced by *Pseudomonas aeruginosa* inhibit the beating of human respiratory cilia in vitro. *J. Clin. Invest.* **79**, 221–229.
- Wang, C., Yang, Y., Mei, Z., and Yang, X.** 2013. Cytotoxic compounds from *Laminaria japonica*. *Chem. Nat. Compd.* **49**, 699–701.