

STRONG CONVERGENCE OF AN ITERATIVE ALGORITHM FOR A CLASS OF NONLINEAR SET-VALUED VARIATIONAL INCLUSIONS

XIE PING DING AND SALAHUDDIN

ABSTRACT. In this communication, we introduce an Ishikawa type iterative algorithm for finding the approximate solutions of a class of nonlinear set valued variational inclusion problems. We also establish a characterization of strong convergence of this iterative techniques.

1. Introduction

The variational inequalities were initially studied by Kinderlehrer and Stampacchia [1]. Since then, they have been widely investigated. They cover partial differential equations, optimal controls, optimizations, mathematical programming, mechanics, economics, transportation and finances, see [2-9]. In 1994, Hassouni and Moudafi [10] introduced a class of variational inequalities which includes various classes of variational inequalities as special cases. Since then, there are a great number of numerical methods for solving various variational inequalities and variational inclusions. It is well known that monotonicity, accretivity of the underlying operators and their generalizations plays indispensable roles for solving the generalized variational inequalities and

Received December 16, 2016. Revised December 30, 2016. Accepted February 07, 2017.

2010 Mathematics Subject Classification: 40H90.

Key words and phrases: Nonlinear set valued variational inclusions; Iterative algorithm; m -accretive mappings; ϕ -strongly accretive mappings; \mathcal{H} -generalized Lipschitz continuous mappings; \mathcal{H} -mixed Lipschitz continuous mapping.

© The Kangwon-Kyungki Mathematical Society, 2017.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

generalized variational inclusions. For example, see [11-41].

Let \mathcal{X} be a real Banach space with norm $\|\cdot\|$, \mathcal{X}^* be the topological dual space of \mathcal{X} and $\langle \cdot, \cdot \rangle$ be the generalized duality pairing between \mathcal{X} and \mathcal{X}^* . Let J denote the normalized duality mapping from \mathcal{X} to $2^{\mathcal{X}^*}$ defined by

$$Jx = \{f \in \mathcal{X}^* : \langle x, f \rangle = \|x\| \cdot \|f\|, \|f\| = \|x\|\}, x \in \mathcal{X}.$$

In this work, we shall denote the single valued normalized duality mapping by j . It is well known that if \mathcal{X} is a smooth then j is a single valued. Also we denote by $\mathcal{H}(\cdot, \cdot)$, the Hausdorff metric on $CB(\mathcal{X})$ defined by

$$\mathcal{H}(A, B) = \max\{\sup_{x \in B} \inf_{y \in A} d(x, y), \sup_{x \in A} \inf_{y \in B} d(x, y)\}, A, B \in CB(\mathcal{X})$$

where $CB(\mathcal{X})$ denote the family of all nonempty closed and bounded subsets of \mathcal{X} .

DEFINITION 1.1 A set valued mapping $A : D(A) \subset \mathcal{X} \rightarrow 2^{\mathcal{X}^*}$ is said to be

- (i) accretive, if for any $x, y \in D(A)$, there exists $j(x - y) \in J(x - y)$ such that for all $u \in Ax, v \in Ay$

$$\langle u - v, j(x - y) \rangle \geq 0;$$

- (ii) strictly accretive if A is accretive and

$$\langle u - v, j(x - y) \rangle = 0 \text{ if and only if } x = y;$$

- (iii) m -accretive if A is accretive and $(I + \rho A)(D(A)) = \mathcal{X}$ for every $\rho > 0$ where I is an identity mapping;

- (iv) ϕ -strongly accretive if for any $x, y \in D(A)$ there exists $j(x - y) \in J(x - y)$ such that for all $u \in Ax$ and $v \in Ay$

$$\langle u - v, j(x - y) \rangle \geq \phi(\|x - y\|)\|x - y\|,$$

where $\phi : [0, \infty) \rightarrow [0, \infty)$ is a strictly increasing continuous function with $\phi(0) = 0$;

- (v) ϕ -expansive, if any $x, y \in D(A), u \in Ax$ and $v \in Ay$

$$\|u - v\| \geq \phi(\|x - y\|).$$

REMARK 1.1

- (i) If A is ϕ -strongly accretive then A is ϕ -expansive.

- (ii) If $\mathcal{X} = H$ is a Hilbert space, then A is m -accretive if and only if A is maximal monotone. A is said to be maximal monotone if its graph is not properly contained in the graph of any other monotone mapping. In addition, the concepts of accretiveness and strong accretiveness reduce to the one of monotonicity and strong monotonicity, respectively.

DEFINITION 1.2 A mapping $T : \mathcal{X} \rightarrow CB(\mathcal{X})$ is said to be

- (i) \mathcal{H} -continuous if for any given $\epsilon > 0$ and each $\hat{x} \in \mathcal{X}$ there exists a $\delta = \delta(\epsilon, \hat{x}) > 0$ such that whenever $\|\hat{x} - y\| < \delta$ then

$$\mathcal{H}(T\hat{x}, Ty) < \epsilon,$$

- (ii) \mathcal{H} -uniformly continuous if for any given $\epsilon > 0$ there exists a $\delta = \delta(\epsilon) > 0$ such that whenever $\|x - y\| < \delta$ then

$$\mathcal{H}(Tx, Ty) < \epsilon.$$

DEFINITION 1.3 [34] A mapping $T : \mathcal{X} \rightarrow 2^{\mathcal{X}}$ is called

- (i) lower semi continuous if $T^{-1} = \{x \in E : Tx \cap V \neq \emptyset\}$ is open in \mathcal{X} whenever $V \subset \mathcal{X}$ is open;
- (ii) γ - \mathcal{H} -generalized Lipschitz continuous if there exists a constant $\gamma > 0$ such that for all $x, y \in \mathcal{X}$

$$\mathcal{H}(Tx, Ty) \leq \gamma(1 + \|x - y\|).$$

DEFINITION 1.4 A mapping $N : \mathcal{X} \times \mathcal{X} \rightarrow \mathcal{X}$ is said to be ϕ -strongly accretive with respect to $T : \mathcal{X} \rightarrow CB(\mathcal{X})$ in the first argument if for any $x, y \in D(T)$ there exists $j(x - y) \in J(x - y)$ such that for all $z \in \mathcal{X}, u \in Tx$ and $v \in Ty$

$$\langle N(u, z) - N(v, z), j(x - y) \rangle \geq \phi(\|x - y\|)\|x - y\|,$$

where $\phi : [0, \infty) \rightarrow [0, \infty)$ is a strictly increasing continuous function with $\phi(0) = 0$.

DEFINITION 1.5 Let \mathcal{X} be a uniformly smooth Banach space and let $N : \underbrace{\mathcal{X} \times \mathcal{X} \cdots \mathcal{X}}_l \rightarrow \mathcal{X}$ be a single valued mapping. N is said to be $(\xi_1, \xi_2, \dots, \xi_l)$ -mixed Lipschitz continuous if there exist constants $\xi_i >$

0, $i = 1, 2, \dots, l$ such that

$$\begin{aligned} & \|N(x_1, x_2, \dots, x_l) - N(y_1, y_2, \dots, y_l)\| \\ & \leq \sum_{i=1}^l \xi_i \|x_i - y_i\|, \quad \forall x_i, y_i \in \mathcal{X}, (i = 1, 2, \dots, l). \end{aligned}$$

LEMMA 1.1 [33] *Let \mathcal{X} be a real smooth Banach space, then for any $x, y \in \mathcal{X}$*

$$\|x + y\|^2 \leq \|x\|^2 + 2\langle y, j(x + y) \rangle,$$

where $j : \mathcal{X} \rightarrow \mathcal{X}^*$ is a normalized duality mapping on \mathcal{X} .

LEMMA 1.2 [30] *Let \mathcal{X} be a real smooth Banach space. Let $T, F : \mathcal{X} \rightarrow 2^{\mathcal{X}}$ be the two set valued mappings and $N(\cdot, \cdot) : \mathcal{X} \times \mathcal{X} \rightarrow \mathcal{X}$ be a nonlinear mapping satisfying the following conditions:*

- (i) *the mapping $x \rightarrow N(x, y)$ is ϕ -strongly accretive with respect to the mapping T ,*
- (ii) *the mapping $y \rightarrow N(x, y)$ is accretive with respect to the mapping F .*

Then the mapping $S : \mathcal{X} \rightarrow 2^{\mathcal{X}}$ defined by $Sx = N(Tx, Fx)$ is ϕ -strongly accretive mapping.

LEMMA 1.3 [26] *Let \mathcal{X} be a real Banach space and $S : \mathcal{X} \rightarrow 2^{\mathcal{X}} \setminus \{\phi\}$ be a lower semi continuous and ϕ -strongly accretive mapping. Then for any $x \in \mathcal{X}$, Sx is a one point set, i.e., S is a single valued mapping.*

LEMMA 1.4 [41] *Let \mathcal{X} be a metric space, $T : \mathcal{X} \rightarrow CB(\mathcal{X})$ be a set valued mapping. Therefore for any given $\epsilon > 0$ and for any given $x, y \in \mathcal{X}$, $u \in Tx$, there exists $v \in Ty$ such that,*

$$d(u, v) \leq (1 + \epsilon)\mathcal{H}(Tx, Ty)$$

where $\mathcal{H}(\cdot, \cdot)$ is a Hausdorff metric on $CB(\mathcal{X})$.

Let $T_1, T_2, \dots, T_l : \mathcal{X} \rightarrow CB(\mathcal{X})$ be the set valued mappings and $N : \underbrace{\mathcal{X} \times \mathcal{X} \cdots \mathcal{X}}_l \rightarrow \mathcal{X}$ be the nonlinear mapping. Let $F : \mathcal{X} \rightarrow 2^{\mathcal{X}}$ be

a set valued mapping and let $A : D(A) \subset \mathcal{X} \rightarrow 2^{\mathcal{X}}$ be a m -accretive mapping and let $g : \mathcal{X} \rightarrow D(A)$ and $h : \mathcal{X} \rightarrow \mathcal{X}$ be the single valued mappings. In this paper, we will consider the following nonlinear set

valued variational inclusion problems: for any given $f \in \mathcal{X}, \lambda > 0$, finding $x \in D(A)$ such that (x, w_1, \dots, w_l, v) is a solution set of

$$(1.1) \quad f \in h(v) + N(w_1, w_2, \dots, w_l) + \lambda(A(g(x)))$$

for any $w_i \in T_i(x), i = 1, 2, \dots, l$ and $v \in F(x)$.

2. Main Results

In this section, we define an Ishikawa type iterative algorithm for solving the nonlinear set valued variational inclusion problems (1.1). Moreover, we give a characterization of strong convergence of this algorithm to the solutions set of the nonlinear set valued variational inclusion problems (1.1)

ALGORITHM 2.1 Let $\epsilon > 0$ be a some given number. Let \mathcal{X} be a real smooth Banach space. Let $A : D(A) \subset \mathcal{X} \rightarrow 2^{\mathcal{X}}$ be a m -accretive mapping, $g : \mathcal{X} \rightarrow D(A)$ and $h : \mathcal{X} \rightarrow \mathcal{X}$ be the single valued mappings. Let $F : \mathcal{X} \rightarrow CB(\mathcal{X})$ and $T_1, T_2, \dots, T_l : \mathcal{X} \rightarrow CB(\mathcal{X})$ be the set valued mappings and $N : \underbrace{\mathcal{X} \times \mathcal{X} \cdots \mathcal{X}}_l \rightarrow \mathcal{X}$ be the nonlinear mapping. Let

$\{\alpha_n\}_{n=0}^{\infty}$ and $\{\beta_n\}_{n=0}^{\infty}$ be the two sequences in $[0, 1]$. For any given initial $x_0 \in D(A), \bar{w}_{0,i} \in T_i(x_0), i = 1, 2, \dots, l, \bar{v}_0 \in F(x_0)$ and $\bar{u}_0 \in A(g(x_0))$ define by

$$y_0 = (1 - \beta_0)x_0 + \beta_0(f + x_0 - h(\bar{v}_0) - N(\bar{w}_{0,1}, \bar{w}_{0,2}, \dots, \bar{w}_{0,l}) - \lambda\bar{u}_0).$$

Since $\bar{u}_0 \in A(g(x_0))$, by Lemma 1.4, there exists $u_0 \in A(g(y_0))$ such that

$$\|\bar{u}_0 - u_0\| \leq (1 + \epsilon) \mathcal{H}(A(g(x_0)), A(g(y_0))),$$

$$\bar{w}_{0,i} \in T_i(x_0), w_{0,i} \in T_i(y_0)$$

$$\|\bar{w}_{0,i} - w_{0,i}\| \leq (1 + \epsilon) \mathcal{H}(T_i(x_0), T_i(y_0)), \text{ for } i = 1, 2, \dots, l,$$

$$\text{and } \bar{v}_0 \in F(x_0), v_0 \in F(y_0)$$

$$\|\bar{v}_0 - v_0\| \leq (1 + \epsilon) \mathcal{H}(F(x_0), F(y_0)).$$

For any given $w_{0,i} \in T_i(y_0), i = 1, 2, \dots, l$ and $u_0 \in A(g(y_0)), v_0 \in F(y_0)$, define

$$x_1 = (1 - \alpha_0)x_0 + \alpha_0(f + y_0 - h(v_0) - N(w_{0,1}, \dots, w_{0,l}) - \lambda u_0).$$

Since $\bar{u}_0 \in A(g(x_0))$, by Lemma 1.4, there exists $\bar{u}_1 \in A(g(x_1))$ such that

$$\|\bar{u}_0 - \bar{u}_1\| \leq (1 + \epsilon) \mathcal{H}(A(g(x_0)), A(g(x_1))),$$

$$\bar{w}_{0,i} \in T_i(x_0), \bar{w}_{1,i} \in T_i(x_1)$$

$$\|\bar{w}_{0,i} - \bar{w}_{1,i}\| \leq (1 + \epsilon) \mathcal{H}(T_i(x_0), T_i(x_1)), \text{ for } i = 1, 2, \dots, l,$$

$$\text{and } \bar{v}_0 \in F(x_0), \bar{v}_1 \in F(x_1)$$

$$\|\bar{v}_0 - \bar{v}_1\| \leq (1 + \epsilon) \mathcal{H}(F(x_0), F(x_1)).$$

For any given $\bar{w}_{1,i} \in T_i(x_1)$, define

$$y_1 = (1 - \beta_1)x_1 + \beta_1(f + x_1 - h(\bar{v}_1) - N(\bar{w}_{1,1}, \bar{w}_{1,2}, \dots, \bar{w}_{1,l}) - \lambda\bar{u}_1).$$

Since $\bar{u}_1 \in A(g(x_1))$, by Lemma 1.4, there exists $u_1 \in A(g(y_1))$ such that

$$\|\bar{u}_1 - u_1\| \leq (1 + \epsilon) \mathcal{H}(A(g(x_1)), A(g(y_1))),$$

$$\bar{w}_{1,i} \in T_i(x_1), w_{1,i} \in T_i(y_1)$$

$$\|\bar{w}_{1,i} - w_{1,i}\| \leq (1 + \epsilon) \mathcal{H}(T_i(x_1), T_i(y_1)), \text{ for } i = 1, 2, \dots, l,$$

$$\text{and } \bar{v}_1 \in F(x_1), v_1 \in F(y_1)$$

$$\|\bar{v}_1 - v_1\| \leq (1 + \epsilon) \mathcal{H}(F(x_1), F(y_1)).$$

For any given $w_{1,i} \in T_i(y_1)$, $i = 1, 2, \dots, l$ and $u_1 \in A(g(y_1))$, $v_1 \in F(y_1)$, define

$$x_2 = (1 - \alpha_1)x_1 + \alpha_1(f + y_1 - h(v_1) - N(w_{1,1}, \dots, w_{1,l}) - \lambda u_1).$$

Continuing in this way, we can compute the sequences $\{x_n\}_{n=0}^\infty$, $\{v_n\}_{n=0}^\infty$ and $\{w_{n,i}\}_{n=0}^\infty$, $i = 1, 2, \dots, l$ by the iterative schemes such that for $n = 0, 1, 2, 3, \dots$

(i)

$$x_{n+1} = (1 - \alpha_n)x_n + \alpha_n(f + y_n - h(v_n) - N(w_{n,1}, \dots, w_{n,l}) - \lambda u_n),$$

$$\text{for } w_{n,i} \in T_i(y_n), \quad i = 1, 2, \dots, l, v_n \in F(y_n), u_n \in A(g(y_n));$$

(ii)

$$y_n = (1 - \beta_n)x_n + \beta_n(f + x_n - h(\bar{v}_n) - N(\bar{w}_{n,1}, \bar{w}_{n,2}, \dots, \bar{w}_{n,l}) - \lambda\bar{u}_n),$$

$$\text{for any } \bar{w}_{n,i} \in T_i(x_n), \quad i = 1, 2, \dots, l, \bar{v}_n \in F(x_n) \text{ and } \bar{u}_n \in A(g(x_n));$$

- (iii) $\bar{u}_n \in A(g(x_n)), u_n \in A(g(y_n))$
 $\|\bar{u}_n - u_n\| \leq (1 + \epsilon) \mathcal{H}(A(g(x_n)), A(g(y_n)));$
 $\bar{w}_{n,i} \in T_i(x_n), w_{n,i} \in T_i(y_n)$
 $\|\bar{w}_{n,i} - w_{n,i}\| \leq (1 + \epsilon) \mathcal{H}(T_i(x_n), T_i(y_n)), \text{ for } i = 1, 2, \dots, l,$
and $\bar{v}_n \in F(x_n), v_n \in F(y_n)$
 $\|\bar{v}_n - v_n\| \leq (1 + \epsilon) \mathcal{H}(F(x_n), F(y_n));$
- (iv) $\bar{u}_n \in A(g(x_n)), \bar{u}_{n+1} \in A(g(x_{n+1}))$
 $\|\bar{u}_n - \bar{u}_{n+1}\| \leq (1 + \epsilon) \mathcal{H}(A(g(x_n)), A(g(x_{n+1}))),$
 $\bar{w}_{n,i} \in T_i(x_n), \bar{w}_{n+1,i} \in T_i(x_{n+1});$
 $\|\bar{w}_{n,i} - \bar{w}_{n+1,i}\| \leq (1 + \epsilon) \mathcal{H}(T_i(x_n), T_i(x_{n+1})), \text{ for } i = 1, 2, \dots, l,$
and $\bar{v}_n \in F(x_n), \bar{v}_{n+1} \in F(x_{n+1})$
 $\|\bar{v}_n - \bar{v}_{n+1}\| \leq (1 + \epsilon) \mathcal{H}(F(x_n), F(x_{n+1})).$

The sequences $\{x_n\}_{n=0}^\infty$, $\{v_n\}_{n=0}^\infty$ and $\{w_{n,i}\}_{n=0}^\infty, i = 1, 2, \dots, l$ defined by (i)-(iv) is called the Ishikawa type iterative sequences.

THEOREM 2.1 *Let \mathcal{X} be a real smooth Banach space. Let $T_1, T_2, \dots, T_l : \mathcal{X} \rightarrow CB(\mathcal{X})$ and $A : D(A) \subset \mathcal{X} \rightarrow 2^{\mathcal{X}}$ be the set valued mappings. Let $g : \mathcal{X} \rightarrow D(A)$ and $h : \mathcal{X} \rightarrow \mathcal{X}$ be the single valued mappings. Let $F : \mathcal{X} \rightarrow CB(\mathcal{X})$ be the set valued mapping and let $N : \underbrace{\mathcal{X} \times \mathcal{X} \cdots \mathcal{X}}_l \rightarrow \mathcal{X}$ be a single valued continuous mapping satisfying the following conditions:*

- (i) $A \circ g : \mathcal{X} \rightarrow 2^{\mathcal{X}}$ is m -accretive and \mathcal{H} -uniformly continuous mapping;
- (ii) $T_1, T_2, \dots, T_l : \mathcal{X} \rightarrow CB(\mathcal{X})$ is lower semi continuous and $\gamma_i - \mathcal{H}$ -Lipschitz continuous mapping for $i = 1, 2, \dots, l$
- (iii) the mapping $x_1 \rightarrow N(x_1, x_2, \dots, x_l)$ is ϕ -strongly accretive and ξ_i -mixed Lipschitz continuous mapping with constants $\xi_i > 0, i = 1, 2, \dots, l$ with respect to T_i where $\phi : [0, \infty) \rightarrow [0, \infty)$ is a strictly increasing function with $\phi(0) = 0$;
- (iv) the mapping $x \rightarrow N(T_1x, \dots, T_lx)$ is uniformly continuous;
- (v) h is δ -Lipschitz continuous mapping;
- (vi) $F : \mathcal{X} \rightarrow CB(\mathcal{X})$ be the lower semi continuous and $\eta - \mathcal{H}$ -Lipschitz continuous mapping;

Suppose $\{\alpha_n\}_{n=0}^\infty$ and $\{\beta_n\}_{n=0}^\infty$ are the sequences in $[0, 1]$ such that $\sum_{n=0}^\infty \alpha_n = \infty$, $\lim_{n \rightarrow \infty} \alpha_n = 0$ and $\lim_{n \rightarrow \infty} \beta_n = 0$. Then the sequences $\{x_n\}_{n=0}^\infty$, $\{w_{n,i}\}_{n=0}^\infty$, $i = 1, 2, \dots, l$ and $\{v_n\}_{n=0}^\infty$ generated by Algorithm 2.1 converges strongly to $x^* \in X$, if and only if the sequences $\{x_n\}_{n=0}^\infty$, $\{w_{n,i}\}_{n=0}^\infty$, $\{v_n\}_{n=0}^\infty$, $\{\mathcal{H}(A(g(x_n)), A(g(x^*)))\}_{n=0}^\infty$, $\{\mathcal{H}(T_i(x_n), T_i(x^*))\}_{n=0}^\infty$, $\{\mathcal{H}(F(x_n), F(x^*))\}_{n=0}^\infty$, $i = 1, 2, \dots, l$ are bounded where for any $u^* \in A(g(x^*))$, $v^* \in F(x^*) \in \mathcal{X}$, $w_i^* \in T_i(x^*) \in \mathcal{X}$, (x^*, w_i^*, v^*) , $i = 1, 2, \dots, l$ is a solution set of nonlinear set valued variational inclusion problems (1.1).

Proof. Let $x^* \in \mathcal{X}$ such that $f \in h(v) + N(w_1, \dots, w_l) + \lambda Aog(x^*)$ for any $w_i \in T_i(x^*)$, $v \in F(x^*)$, $i = 1, 2, \dots, l$, whose existence is guaranted by [35, Theorem 3.1] without loss of generality, we may assume that $f \equiv 0$ and $\lambda = 1$. Take $w_i^* \in T_i(x^*)$, $i = 1, 2, \dots, l$, and $u^* \in A(g(x^*))$, $v^* \in F(x^*)$ such that

$$0 = h(v^*) + N(w_1^*, \dots, w_l^*) + u^*.$$

NECESSITY.

Suppose that

$$\|x_n - x^*\| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Therefore, from the \mathcal{H} -uniform continuity of Aog ,

$$\mathcal{H}(A(g(x_n)), A(g(x^*))) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Again from

$$\|w_{n,i} - w_i^*\| \rightarrow 0 \text{ as } n \rightarrow \infty \text{ for } i = 1, 2, \dots, l,$$

$$\|v_n - v^*\| \rightarrow 0 \text{ as } n \rightarrow \infty,$$

it follows from the \mathcal{H} -uniform continuity of T_i and F that

$$\mathcal{H}(T_i(x_n), T_i(x^*)) \rightarrow 0 \text{ as } n \rightarrow \infty, \text{ for } i = 1, 2, \dots, l,$$

$$\mathcal{H}(F(x_n), F(x^*)) \rightarrow 0 \text{ as } n \rightarrow \infty.$$

Thus the sequences $\{x_n\}_{n=0}^\infty$, $\{v_n\}_{n=0}^\infty$, $\{w_{n,i}\}_{n=0}^\infty$, $\{\mathcal{H}(A(g(x_n)), A(g(x^*)))\}_{n=0}^\infty$, $\{\mathcal{H}(T_i(x_n), T_i(x^*))\}_{n=0}^\infty$ for $i = 1, 2, \dots, l$ and $\{\mathcal{H}(F(x_n), F(x^*))\}_{n=0}^\infty$ are all bounded.

SUFFICIENCY.

Suppose that the sequences $\{x_n\}_{n=0}^\infty$, $\{w_{n,i}\}_{n=0}^\infty$, $\{\mathcal{H}(A(g(x_n)), A(g(x^*)))\}_{n=0}^\infty$, $\{\mathcal{H}(T_i(x_n), T_i(x^*))\}_{n=0}^\infty$, $\{\mathcal{H}(F(x_n), F(x^*))\}_{n=0}^\infty$ for $i = 1, 2, \dots, l$, are all bounded. Then we divide the proof of the sufficiency into five steps:

STEP 1. We claim that the sequences $\{\bar{w}_{n,i}\}_{n=0}^\infty, \{\bar{u}_n\}_{n=0}^\infty, \{\bar{v}_n\}_{n=0}^\infty, i = 1, 2, \dots, l$ are bounded. Indeed by Nadler's Theorem for each $n \geq 0$ there exist $\bar{w}_i \in T_i(x^*), i = 1, 2, \dots, l, \bar{v} \in F(x^*)$ and $\bar{u} \in A(g(x^*))$ such that

$$\begin{aligned} \|\bar{w}_{n,i} - \bar{w}_i\| &\leq (1 + \epsilon_i) \mathcal{H}(T_i(x_n), T_i(x^*)), \quad i = 1, 2, \dots, l; \\ \|\bar{u}_n - \bar{u}\| &\leq (1 + \epsilon) \mathcal{H}(A(g(x_n)), A(g(x^*))); \\ \|\bar{v}_n - \bar{v}\| &\leq (1 + \epsilon) \mathcal{H}(F(x_n), F(x^*)). \end{aligned}$$

Since $T_i : \mathcal{X} \rightarrow \mathcal{CB}(\mathcal{X})$ is $\gamma_i - \mathcal{H}$ -generalized Lipschitz continuous and $F : \mathcal{X} \rightarrow \mathcal{CB}(\mathcal{X})$ is $\eta - \mathcal{H}$ -generalized Lipschitz continuous mapping, we have

$$\begin{aligned} \|\bar{w}_{n,i} - w_i^*\| &\leq \|\bar{w}_{n,i} - \bar{w}_i\| + \|w_i^* - \bar{w}_i\| \\ &\leq (1 + \epsilon_i) \mathcal{H}(T_i(x_n), T_i(x^*)) + \text{diam}(T_i(x^*)), \quad i = 1, 2, \dots, l \\ (2.2) \quad &\leq (1 + \epsilon_i)\gamma_i(1 + \|x_n - x^*\|) + \text{diam}(T_i(x^*)), \quad i = 1, 2, \dots, l; \end{aligned}$$

$$\begin{aligned} \|\bar{v}_n - v^*\| &\leq \|\bar{v}_n - \bar{v}\| + \|v^* - \bar{v}\| \\ &\leq (1 + \epsilon) \mathcal{H}(F(x_n), F(x^*)) + \text{diam}(F(x^*)) \\ (2.3) \quad &\leq (1 + \epsilon)\eta(1 + \|x_n - x^*\|) + \text{diam}(F(x^*)). \end{aligned}$$

Thus the sequences $\{\bar{w}_{n,i}\}_{n=0}^\infty, \{\bar{v}_n\}_{n=0}^\infty$ are bounded. Also, we note that

$$\begin{aligned} \|\bar{u}_n - u^*\| &\leq \|\bar{u}_n - \bar{u}\| + \|u^* - \bar{u}\| \\ &\leq (1 + \epsilon) \mathcal{H}(A(g(x_n)), A(g(x^*))) + \text{diam}(A(g(x^*))) \\ (2.4) \quad &\leq (1 + \epsilon)(1 + \|A(g(x_n)) - A(g(x^*))\|) + \text{diam}(A(g(x^*))). \end{aligned}$$

Hence the sequence $\{u_n\}_{n=0}^\infty$ is a bounded.

STEP 2. We claim that $\|y_n - x_n\| \rightarrow 0$ as $n \rightarrow \infty$. Indeed, since the mapping $x \rightarrow N(w_1, w_2 \dots w_l)$ is $\xi_i - \mathcal{H}$ -mixed Lipschitz continuous mapping with respect to $T_i, i = 1, 2, \dots, l$ and from Algorithm 2.1, we have

$$\begin{aligned} \|y_n - x_n\| &= \|-\beta_n(h(\bar{v}_n) + N(\bar{w}_{n,1}, \dots, \bar{w}_{n,l}) + \bar{u}_n)\| \\ &\leq \beta_n \|h(\bar{v}_n) - h(v^*)\| + \beta_n \|N(w_1^*, \dots, w_l^*) - N(\bar{w}_{n,1}, \dots, \bar{w}_{n,l})\| + \beta_n \|\bar{u}_n - u^*\| \\ &\leq \beta_n \delta \|\bar{v}_n - v^*\| + \beta_n (\|N(\bar{w}_{n,1}, \dots, \bar{w}_{n,l}) - N(w_1^*, \bar{w}_{n,2} \dots \bar{w}_{n,l})\| \\ &\quad + \dots + \|N(w_1^*, \dots, \bar{w}_{n,l}) - N(w_1^*, \dots, w_l^*)\|) + \beta_n \|\bar{u}_n - u^*\| \\ &\leq \beta_n \delta \|\bar{v}_n - v^*\| + \beta_n (\xi_1 \|\bar{w}_{n,1} - w_1^*\| + \dots + \xi_l \|\bar{w}_{n,l} - w_l^*\|) + \beta_n \|\bar{u}_n - u^*\|, \end{aligned}$$

which implies that $\|y_n - x_n\| \rightarrow 0$ as $n \rightarrow \infty$.

STEP 3. We claim that the sequences $\{v_n\}_{n=0}^\infty, \{w_{n,1}\}_{n=0}^\infty, \dots, \{w_{n,l}\}_{n=0}^\infty, \{u_n\}_{n=0}^\infty$ are bounded. Indeed by Algorithm 2.1(iii), we have

$$(2.5) \quad \begin{aligned} \|u_n\| &\leq \|\bar{u}_n - u_n\| + \|\bar{u}_n\| \\ &\leq (1 + \epsilon) \mathcal{H}(A(g(x_n)), A(g(y_n))) + \|\bar{u}_n\|. \end{aligned}$$

$$(2.6) \quad \begin{aligned} \|w_{n,i}\| &\leq \|\bar{w}_{n,i} - w_{n,i}\| + \|\bar{w}_{n,i}\| \\ &\leq (1 + \epsilon_i) \mathcal{H}(T_i(x_n), T_i(y_n)) + \|\bar{w}_{n,i}\|, \text{ for } i = 1, 2, \dots, l. \end{aligned}$$

$$(2.7) \quad \begin{aligned} \|v_n\| &\leq \|\bar{v}_n - v_n\| + \|\bar{v}_n\| \\ &\leq (1 + \epsilon) \mathcal{H}(F(x_n), F(y_n)) + \|\bar{v}_n\|. \end{aligned}$$

From step 2 and \mathcal{H} -uniform continuity of $A \circ g, T_i,$ and $F,$ we have

$$\begin{aligned} \mathcal{H}(A(g(x_n)), A(g(y_n))) &\rightarrow 0 \text{ as } n \rightarrow \infty; \\ \mathcal{H}(T_i(x_n), T_i(y_n)) &\rightarrow 0 \text{ as } n \rightarrow \infty, \text{ for } i = 1, 2, \dots, l; \\ \mathcal{H}(F(x_n), F(y_n)) &\rightarrow 0 \text{ as } n \rightarrow \infty. \end{aligned}$$

Hence the sequences $\{\mathcal{H}(A(g(x_n)), A(g(y_n)))\}_{n=0}^\infty, \{\mathcal{H}(T_i(x_n), T_i(y_n))\}_{n=0}^\infty, \{\mathcal{H}(F(x_n), F(y_n))\}_{n=0}^\infty$ for $i = 1, 2, \dots, l$ are bounded. Thus it follows from the boundedness of $\{\bar{u}_n\}_{n=0}^\infty, \{\bar{w}_{n,i}\}_{n=0}^\infty, \{\bar{v}_n\}_{n=0}^\infty$ and (2.1) that $\{u_n\}_{n=0}^\infty, \{w_{n,i}\}_{n=0}^\infty, \{v_n\}_{n=0}^\infty$ are bounded for $i = 1, 2, \dots, l.$

On the other hand, by Nadler's Theorem for each $n \geq 0$ there exists $\hat{w}_i \in T_i(x^*), i = 1, 2, \dots, l, \hat{v} \in F(x^*)$ such that

$$\begin{aligned} \|v_n - \hat{v}\| &\leq (1 + \epsilon) \mathcal{H}(F(y_n), F(x^*)); \\ \|w_{n,i} - \hat{w}_i\| &\leq (1 + \epsilon_i) \mathcal{H}(T_i(y_n), T_i(x^*)), \quad i = 1, 2, \dots, l. \end{aligned}$$

Observe that

$$\begin{aligned} \|w_{n,i} - w_i^*\| &\leq \|w_{n,i} - \hat{w}_i\| + \|\hat{w}_i - w_i^*\| \\ &\leq (1 + \epsilon_i) \mathcal{H}(T_i(y_n), T_i(x^*)) + \text{diam}(T_i(x^*)) \\ &\leq (1 + \epsilon_i)\gamma_i(1 + \|y_n - x^*\|) + \text{diam}(T_i(x^*)), \text{ for } i = 1, 2, \dots, l. \\ \|v_n - v^*\| &\leq \|v_n - \hat{v}\| + \|\hat{v} - v^*\| \\ &\leq (1 + \epsilon) \mathcal{H}(F(y_n), F(x^*)) + \text{diam}(F(x^*)) \\ &\leq (1 + \epsilon)\eta(1 + \|y_n - x^*\|) + \text{diam}(F(x^*)). \end{aligned}$$

Since the sequence $\{x_n\}_{n=0}^\infty$ is bounded and $\|y_n - x_n\| \rightarrow 0$ as $n \rightarrow \infty,$ so $\{y_n\}_{n=0}^\infty$ is bounded. Hence $\{w_{n,i}\}_{n=0}^\infty, i = 1, 2, \dots, l$ and $\{v_n\}_{n=0}^\infty$ are bounded.

STEP 4. We claim that

$$\|h(v_n) + N(w_{n,1}, \dots, w_{n,l}) + u_n - h(\bar{v}_{n+1}) - N(\bar{w}_{n+1,1}, \dots, \bar{w}_{n+1,l}) - \bar{u}_{n+1}\| \rightarrow 0$$

as $n \rightarrow \infty$. Since

$$\|h(v_n) - h(v^*)\| \leq \delta \|v_n - v^*\|$$

and

$$\|N(w_{n,1}, \dots, w_{n,l}) - N(w_1^*, \dots, w_l^*)\| \leq \xi_1 \|w_{n,1} - w_1^*\| + \dots + \xi_l \|w_{n,l} - w_l^*\|.$$

It follows from the boundedness of $\{v_n\}_{n=0}^\infty$, $\{w_{n,i}\}_{n=0}^\infty$, $i = 1, 2, \dots, l$, that $\{h(v_n)\}_{n=0}^\infty$ and $\{N(w_{n,1}, \dots, w_{n,l})\}_{n=0}^\infty$ are bounded. Furthermore, from Algorithm 2.1 (i) we deduce that as $n \rightarrow \infty$,

$$\begin{aligned} \|x_{n+1} - x_n\| &= \alpha_n \|y_n - x_n - h(v_n) - N(w_{n,1}, \dots, w_{n,l}) - u_n\| \\ &\leq \alpha_n \|y_n - x_n\| + \alpha_n \|h(v_n) + N(w_{n,1}, \dots, w_{n,l}) + u_n\| \rightarrow 0 \end{aligned}$$

and hence

$$\|y_n - x_{n+1}\| \leq \|y_n - x_n\| + \|x_n - x_{n+1}\| \rightarrow 0.$$

From [36], the mapping $x \rightarrow h(F(x))$ and $x \rightarrow N(T_1(x), \dots, T_l(x))$ are single valued. Thus from condition (vii) and $\|y_n - x_{n+1}\| \rightarrow 0$ as ($n \rightarrow \infty$), we conclude that as $n \rightarrow \infty$

$$(2.8) \quad \|N(w_{n,1}, \dots, w_{n,l}) - N(\bar{w}_{n+1,1}, \dots, \bar{w}_{n+1,l})\| \rightarrow 0.$$

On the other hand, Algorithm 2.1 and the \mathcal{H} -uniform continuity of Aog , we conclude that as $n \rightarrow \infty$,

$$(2.9) \quad \begin{aligned} &\|u_n - \bar{u}_{n+1}\| \\ &\leq \|u_n - \bar{u}_n\| + \|\bar{u}_n - \bar{u}_{n+1}\| \\ &\leq (1 + \epsilon) \mathcal{H}(A(g(y_n)), A(g(x_n))) + (1 + \epsilon) \mathcal{H}(A(g(x_n)), A(g(x_{n+1}))) \rightarrow 0. \end{aligned}$$

Therefore from (2.7) and (2.8), the assertion is valid.

STEP 5. We claim that the sequence $\{x_n\}_{n=0}^\infty$ converges strongly to x^* . Indeed from [34], the set valued mapping

$$x \rightarrow h(F(x)) + N(T_1(x), \dots, T_l(x)) + A(g(x))$$

is ϕ -strongly accretive mapping. Hence, we have

$$\begin{aligned}
& \langle x_{n+1} - h(\bar{v}_{n+1}) - N(\bar{w}_{n+1,1}, \dots, \bar{w}_{n+1,l}) - \bar{u}_{n+1} - (x^* - h(v^*) \\
& \quad - N(w_1^*, \dots, w_l^*) - u^*), j(x_{n+1} - x^*) \rangle \\
(2.10) \quad & \leq \|x_{n+1} - x^*\|^2 - \langle h(\bar{v}_{n+1}) + N(\bar{w}_{n+1,1}, \dots, \bar{w}_{n+1,l}) + \bar{u}_{n+1} \\
& \quad - (h(v^*) + N(w_1^*, \dots, w_l^*) + u^*), j(x_{n+1} - x^*) \rangle \\
& \leq \|x_{n+1} - x^*\|^2 - \phi(\|x_{n+1} - x^*\|)\|x_{n+1} - x^*\|.
\end{aligned}$$

We observe that from Algorithm 2.1 (i), Lemma 1.6 and (2.9)

$$\begin{aligned}
(2.11) \quad & \|x_{n+1} - x^*\|^2 \\
& = \|(1 - \alpha_n)(x_n - x^*) + \alpha_n(y_n - x^* - h(v_n) - N(w_{n,1}, \dots, w_{n,l}) - u_n)\|^2 \\
& \leq (1 - \alpha_n)^2 \|x_n - x^*\|^2 + 2\alpha_n \langle y_n - x^* - h(v_n) - N(w_{n,1}, \dots, w_{n,l}) \\
& \quad - u_n, j(x_{n+1} - x^*) \rangle \\
& \leq (1 - \alpha_n)^2 \|x_n - x^*\|^2 + 2\alpha_n \langle y_n - h(v_n) - N(w_{n,1}, \dots, w_{n,l}) \\
& \quad - u_n - (x_{n+1} - h(\bar{v}_{n+1}) - N(\bar{w}_{n+1,1}, \dots, \bar{w}_{n+1,l}) - \bar{u}_{n+1}), j(x_{n+1} - x^*) \rangle \\
& \quad + 2\alpha_n \langle x_{n+1} - h(\bar{v}_{n+1}) - N(\bar{w}_{n+1,1}, \dots, \bar{w}_{n+1,l}) - \bar{u}_{n+1} - (x^* - h(v^*) \\
& \quad - N(w_1^*, \dots, w_l^*) - u^*), j(x_{n+1} - x^*) \rangle \\
& \leq (1 - \alpha_n)^2 \|x_n - x^*\|^2 + 2\alpha_n \|y_n - h(v_n) - N(w_{n,1}, \dots, w_{n,l}) - u_n - (x_{n+1} \\
& \quad - h(\bar{v}_{n+1}) - N(\bar{w}_{n+1,1}, \dots, \bar{w}_{n+1,l}) - \bar{u}_{n+1})\| \|x_{n+1} - x^*\| + 2\alpha_n \|x_{n+1} - x^*\|^2 \\
& \quad - 2\alpha_n \phi(\|x_{n+1} - x^*\|)\|x_{n+1} - x^*\|.
\end{aligned}$$

Since $\{x_n\}_{n=0}^\infty$ is bounded, we have

$$M = \sup\{\|x_n - x^*\| : n \geq 0\} < \infty.$$

Note that $\alpha_n \rightarrow 0$ as $n \rightarrow \infty$. Thus without loss of generality, we may assume that $1 - 2\alpha_n > 0$ for all $n \geq 0$. Hence it follows from (2.10) that

$$\begin{aligned}
(2.12) \quad & \|x_{n+1} - x^*\|^2 \\
& \leq \frac{(1 - \alpha_n)^2}{1 - 2\alpha_n} \|x_n - x^*\|^2 + \frac{2\alpha_n}{1 - 2\alpha_n} \|y_n - h(v_n) - N(w_{n,1}, \dots, w_{n,l}) \\
& \quad - u_n - (x_{n+1} - h(\bar{v}_{n+1}) - N(\bar{w}_{n+1,1}, \dots, \bar{w}_{n+1,l}) - \bar{u}_{n+1})\| \|x_{n+1} - x^*\| \\
& \quad - \frac{2\alpha_n}{1 - 2\alpha_n} \phi(\|x_{n+1} - x^*\|) \|x_{n+1} - x^*\| \\
& \leq \|x_n - x^*\|^2 + \frac{2\alpha_n}{1 - 2\alpha_n} \left\{ \frac{M^2\alpha_n}{2} + M \|y_n - h(v_n) - N(w_{n,1}, \dots, w_{n,l}) \right. \\
& \quad \left. - u_n - (x_{n+1} - h(\bar{v}_{n+1}) - N(\bar{w}_{n+1,1}, \dots, \bar{w}_{n+1,l}) - \bar{u}_{n+1})\| \right\} \\
& \quad - \frac{2\alpha_n}{1 - 2\alpha_n} \phi(\|x_{n+1} - x^*\|) \|x_{n+1} - x^*\|.
\end{aligned}$$

Next, we discuss two possible cases:

CASE(1).

$$\inf\{\|x_{n+1} - x^*\| : n \geq 0\} = \sigma > 0.$$

From step 4, we have

$$\begin{aligned}
M^2\alpha_n + 2M \|y_n - h(v_n) - N(w_{n,1}, \dots, w_{n,l}) - u_n - (x_{n+1} - h(\bar{v}_{n+1}) \\
- N(\bar{w}_{n+1,1}, \dots, \bar{w}_{n+1,l}) - \bar{u}_{n+1})\| \rightarrow 0 \text{ as } n \rightarrow \infty.
\end{aligned}$$

Hence there exists a possible integer N_0 such that for all $n \geq N_0$,

$$\begin{aligned}
(2.13) \quad & M^2\alpha_n + 2M \|y_n - h(v_n) - N(w_{n,1}, \dots, w_{n,l}) - u_n - (x_{n+1} - h(\bar{v}_{n+1}) \\
& \quad - N(\bar{w}_{n+1,1}, \dots, \bar{w}_{n+1,l}) - \bar{u}_{n+1})\| \\
& < \phi(\sigma)\sigma.
\end{aligned}$$

Thus, it follows from (2.11) and (2.12) that

$$\begin{aligned}
\|x_{n+1} - x^*\|^2 & \leq \|x_n - x^*\|^2 + \frac{\alpha_n}{1 - 2\alpha_n} \phi(\sigma)\sigma - \frac{2\alpha_n}{1 - 2\alpha_n} \phi(\sigma)\sigma \\
& \leq \|x_n - x^*\|^2 - \frac{\alpha_n}{1 - 2\alpha_n} \phi(\sigma)\sigma,
\end{aligned}$$

which implies that

$$\phi(\sigma)\sigma \sum_{n=N_0}^{\infty} \alpha_n \|x_{N_0} - x^*\|^2 < \infty.$$

This contradicts $\sum_{n=0}^{\infty} \alpha_n = \infty$. Therefore case (1) is false.

CASE(2).

$$\inf\{\|x_{n+1} - x^*\| : n \geq 0\} = 0.$$

In this case, there exists a subsequence $\{x_{n_{j+1}}\} \subset \{x_n\}$ such that

$$\|x_{n_{j+1}} - x^*\| \rightarrow 0 \text{ as } j \rightarrow \infty.$$

Hence, for any given $\epsilon > 0$ there exists a positive integer n_j such that for all $n \geq n_j$

$$(2.14) \quad \begin{aligned} & \|x_{n_{j+1}} - x^*\| \leq \epsilon \\ & M^2 \alpha_n + 2M \|y_n - h(v_n) - N(w_{n,1}, \dots, w_{n,l}) - u_n - (x_{n+1} - h(\bar{v}_{n+1}) \\ & \quad - N(\bar{w}_{n+1,1}, \dots, \bar{w}_{n+1,l}) - \bar{u}_{n+1})\| < \phi(\epsilon)\epsilon. \end{aligned}$$

We claim that

$$\|x_{n_{j+m}} - x^*\| \leq \epsilon, \quad \forall m \geq 1.$$

Indeed, we prove that

$$\|x_{n_{j+2}} - x^*\| \leq \epsilon.$$

If this false then

$$\|x_{n_{j+2}} - x^*\| \geq \epsilon.$$

Hence, we have

$$\phi(\|x_{n_{j+2}} - x^*\|) \geq \phi(\epsilon).$$

Thus, it follow from (2.11) that

$$\begin{aligned} \epsilon^2 < \|x_{n_{j+2}} - x^*\|^2 &\leq \|x_{n_{j+1}} - x^*\|^2 + \frac{\alpha_{n_{j+1}}}{1 - 2\alpha_{n_{j+1}}} \phi(\epsilon)\epsilon - \frac{2\alpha_{n_{j+1}}}{1 - 2\alpha_{n_{j+1}}} \phi(\epsilon)\epsilon \\ &= \|x_{n_{j+1}} - x^*\|^2 - \frac{\alpha_{n_{j+1}}}{1 - 2\alpha_{n_{j+1}}} \phi(\epsilon)\epsilon \\ &\leq \|x_{n_{j+1}} - x^*\|^2 \leq \epsilon^2. \end{aligned}$$

This contradict to show that

$$\|x_{n_{j+2}} - x^*\| \leq \epsilon.$$

By induction we can show that

$$\|x_{n_{j+m}} - x^*\| \leq \epsilon, \quad \forall m \geq 1.$$

Hence

$$\|x_n - x^*\| \rightarrow 0 \text{ as } n \rightarrow \infty.$$

This completes the proof. \square

References

- [1] D. Kinderlehrer and G. Stampacchia, *An introduction to variational inequalities and their applications*, Academic Press, 1980.
- [2] J. P. Aubin, *Mathematical methods of game theory and economics*, North Holland, Amsterdam, The Netherlands, 1982.
- [3] H. Brezis, *Operateurs maximaux monotone er semi groupes de contractions dans les espaces de Hilbert*, North-Holland Mathematics Studies 5 Notes de Matematica (50) North-Holland, Amsterdam, 1973.
- [4] R. E. Bruck and S. Reich, *Accretive operators, Banach limits and dual ergodic theorems*, Bull. Acad. Polon Sci. **12** (1981), 585–589
- [5] R. Glowinski, J. L. Lions and R. Tremolieres, *Numerical analysis of variational inequalities*, North-Holland, Amsterdam, 1981.
- [6] F. Giannessi and A. Mugerì, *Variational inequalities and network equilibrium problems*, Plenum Press, New York NY USA, 1995.
- [7] L. Gioranescu, *Geometry of Banach spaces, duality mapping and nonlinear problems*, kluwer Acad. Press, Amsterdam, 1990.
- [8] N. Kikuchi and J. T. Oden, *Contact problems in elasticity*, SIAM, Philadelphia, 1988.
- [9] P. D. Panagiotoupoulos and G. E. Stavroulakis, *New types of variational principles based on the notion of quasi differentiability*, Acta Mech. **94** (1992), 171–194.
- [10] A. Hassouni and A. Moudafi, *A perturbed algorithms for variational inequalities*, J. Math. Anal. Appl. **185** (1994), 706–712.
- [11] X. P. Ding, *Perturbed proximal point algorithms for generalized quasi variational inclusions*, J. Math. Appl. Appl. **201** (1997), 88–101.
- [12] X. P. Ding, *Proximal point algorithm with errors for generalized strongly nonlinear quasi-variational inclusions*, Appl. Math. Mech. **19** (7) (1998), 637–643.
- [13] X. P. Ding, *On a class of generalized nonlinear implicit quasivariational inclusions*, Appl. Math. Mech. **20** (10) (1999), 1087–1098.
- [14] X. P. Ding, *Generalized implicit quasivariational inclusions with fuzzy set-valued mappings*, Comput. Math. Appl. **38** (1) (1999), 71–79.
- [15] X. P. Ding, *Perturbed proximal point algorithms for general quasi-variational-like inclusions*, J. Computat. Appl. Math. **113** (2000), 153–165.
- [16] X. P. Ding, *Generalized quasi-variational-like inclusions with nonconvex functionals*, Appl. Math. Comput. **122** (2001), 267–282.
- [17] X. P. Ding, *Perturbed Ishikawa Type Iterative Algorithm for Generalized Quasi-variational Inclusions*, Appl. Math. Comput. **14** (2003), 359–373.
- [18] X. P. Ding, *Algorithms of solutions for completely generalized mixed implicit quasivariational inclusions*, Appl. Math. Comput. **148** (1) (2004), 47–66.
- [19] X. P. Ding, *Predictor-Corrector iterative algorithms for solving generalized mixed quasi-variational-like inclusion*, J. Comput. Appl. Math. **182** (1) (2005), 1–12.

- [20] X. P. Ding and Salahuddin, *On a system of general nonlinear variational inclusions in Banach spaces*, Appl. Math. Mech. **36** (12) (2015), 1663–1672, DOI:10.1007/s10483-015-1972-6.
- [21] X. P. Ding and H. R. Feng, *The p -step iterative algorithm for a system of generalized mixed quasi-variational inclusions with (A, η) -accretive operators in q -uniformly smooth Banach spaces*, J. Comput. Appl. Math. **220** (1-2) (2008), 163–174.
- [22] X. P. Ding and H. R. Feng, *Algorithm for solving a new class of generalized nonlinear implicit quasi-variational inclusions in Banach spaces*, Appl. Math. Comput. **208** (2009), 547–555.
- [23] X. P. Ding and Z. B. Wang, *Sensitivity analysis for a system of parametric generalized mixed quasi-variational inclusions involving (K, η) -monotone mappings*, Appl. Math. Comput. **214** (2009) 318–327.
- [24] X. P. Ding, Z. B. Wang, *Auxiliary principle and algorithm for a system of generalized set-valued mixed variational-like inequality problems in Banach spaces*, J. Comput. Appl. Math. **223** (2010), 2876–2883.
- [25] Y. P. Fang and N. J. Huang, *H -accretive operators and resolvent operators technique for solving variational inclusions in Banach spaces*, Appl. Math. Lett. **17** (6) (2004), 647–653.
- [26] X. He, *On ϕ -strongly accretive mapping and some set valued variational problems*, J. Math. Anal. Appl. **277** (2) (2003), 504–511.
- [27] N. J. Huang, *On the generalized implicit quasi variational inequalities*, J. Math. Anal. Appl. **216** (1997), 197–210.
- [28] J. S. Jung and C. H. Morales, *The Mann process for perturbed m -accretive operators in Banach spaces*, Nonlinear Anal. **46** (20) (2001), 231–243.
- [29] S. S. Chang, *Set valued variational inclusions in Banach Spaces*, J. Math. Anal. Appl. **248** (2000), 438–454.
- [30] S. S. Chang, J. K. Kim and H. K. Kim, *On the existence and iterative approximation problems of solutions for set valued variational inclusions in Banach spaces*, J. Math. Anal. Appl. **268** (2002), 89–108.
- [31] S. S. Chang, Y. J. Cho, B. S. Lee and I. J. Jung, *Generalized set valued variational inclusions in Banach spaces*, J. Math. Anal. Appl. **246** (2000), 409–422.
- [32] S. S. Chang, Salahuddin and Y. K. Tang, *A system of nonlinear set valued variational inclusions*, SpringerPlus 2014, 3:318, Doi:10.1186/2193-180-3-318.
- [33] S. S. Chang, Y. J. Cho, B. S. Lee, I. J. Jung and S. M. Kang, *Iterative approximations of fixed points and solutions for strongly accretive and strongly pseudo contractive mappings in Banach Spaces*, J. Math. Anal. Appl. **224** (1998), 149–165.
- [34] L. C. Ceng, S. S. Schaible and J. C. Yao, *On the characterization of strong convergence of an iterative algorithm for a class of multivalued variational inclusions*, Math. Math. Oper. Res. **70** (2009), 1–12.
- [35] Y. J. Cho, H. Y. Zhou, S. M. Kang, S. S. Kim, *Approximations for fixed points of ϕ -hemicontractive mappings by the Ishikawa iterative process with mixed errors*, Math. Comput. Model. **34** (2001), 9–18.

- [36] C. E. Chidume, H. Zegeye and K.R. Kazmi, *Existence and convergence theorem for a class of multivalued variational inclusions in Banach space*, *Nonlinear Anal.* **59** (2004), 649–656.
- [37] M. F. Khan and Salahuddin, *Generalized multivalued nonlinear co-variational inequalities in Banach spaces*, *Funct. Diff. Equations* **14** (2-4) (2007), 299–313.
- [38] M. K. Ahmad and Salahuddin, *Stable perturbed algorithms for a new class of generalized nonlinear implicit quasi variational inclusions in Banach spaces*, *Advances in Pure Math.* **2** (2) (2012), 139–148.
- [39] M. F. Khan and Salahuddin, *Generalized co-complementarity problems in p -uniformly smooth Banach spaces*, *JIPAM, J. Inequal. Pure Appl. Math.* **7** (2), Article 66, (2006), 11 pages.
- [40] R. U. Verma and Salahuddin, *Extended systems of nonlinear vector quasi variational inclusions and extended systems of nonlinear vector quasi optimization problems in locally FC-spaces*, *Commun. Appl. Nonlinear Anal.* **23** (1) (2016), 71–88.
- [41] Jr. S. B. Nadler, *Multivalued contraction mappings*, *Pacific J. Math.* **30** (1969), 475–487.

Xie Ping Ding

College of Mathematics and Software Science
Sichuan Normal University
Chengdu, Sichuan 610066, P.R. China
E-mail: dingxip@sicnu.edu.cn

Salahuddin

Department of Mathematics
Jazan University
Jazan, Kingdom of Saudi Arabia
E-mail: salahuddin12@mailcity.com