

# System-Wide Expression and Function of Olfactory Receptors in Mammals

S. June Oh\*

Department of Pharmacology, Inje University College of Medicine, Busan 47392, Korea

Olfactory receptors (ORs) in mammals are generally considered to function as chemosensors in the olfactory organs of animals. They are membrane proteins that traverse the cytoplasmic membrane seven times and work generally by coupling to heterotrimeric G protein. The OR is a G protein-coupled receptor that binds the guanine nucleotide-binding  $G_{\alpha_{olf}}$  subunit and the  $G\beta\gamma$  dimer to recognize a wide spectrum of organic compounds in accordance with its cognate ligand. Mammalian ORs were originally identified from the olfactory epithelium of rat. However, it has been recently reported that the expression of ORs is not limited to the olfactory organ. In recent decades, they have been found to be expressed in diverse organs or tissues and even tumors in mammals. In this review, the expression and expected function of olfactory receptors that exist throughout an organism's system are discussed.

**Keywords:** chemosensory receptor, ectopic expression, G protein-coupled receptor, olfactory receptor, system-wide expression

## Introduction

Animal olfactory receptors (ORs) make up a G protein-coupled chemosensory receptor (GPCR) family with 7 transmembrane alpha-helices located in the cytoplasmic membrane of cells. They are known to work by binding heterotrimeric guanine nucleotide-binding proteins (G proteins), composed of the olfactory  $G\alpha$  ( $G\alpha_{olf}$ ) subunit and the  $G\beta\gamma$  dimer. GPCRs are classified by their sequence homology [1] or based on their phylogenetic origin [2], and there are more than 16 types of G protein  $\alpha$  subunits [3, 4]. ORs are also known to recognize a wide spectrum of organic compounds in accordance with its cognate ligand. When it was originally identified from rat olfactory epithelium [5], the expression of the protein was thought to be confined to that tissue. After  $G\alpha_{olf}$  [6] and rat ORs [5], OR expression in mammalian germ cells [7, 8] and heart [9] was reported in the early 1990s. In the late 20th century, OR expression in an insulin-secreting cell line and the spleen of rats [10] and human erythroid cells [11] was reported. ORs are largely distributed in olfactory sensory neurons of the nasal epithelium but are also expressed in other non-olfactory tissues [12].

We discuss the physiological functions of ORs and suggest future perspectives for their research.

## Olfactory Receptors in the Brain and Heart-Related Systems

Since the expression of ORs in mammalian germ cells was reported [7], OR expression in various organs and tissues in animals has been observed. The brain and the heart are the fundamental organs that form the basis of animal life. For example, controlling the fluctuation of blood pressure and pulse of an animal is accomplished by endocrine action through autonomic reflex and peptide hormone secretion. Brain and heart are closely connected to each other by various mechanisms, including the nervous system.

For the first time, prostate-specific G protein-coupled receptors (PSGRs), also known as RA1c, were reported to be expressed in the brain of rat [13] and mouse [14] and the heart of rat and mouse [15]. It was not expected to be expressed in human tissues other than prostate.

### Brain

After it was demonstrated that RA1c homologs—Olf59 in rat [13] and Olf78 in mouse [14]—were expressed in areas

Received March 6, 2018; Revised March 13, 2018; Accepted March 13, 2018

\*Corresponding author: Tel: +82-51-893-3569, Fax: +82-51-797-8569, E-mail: o@biophilos.org

Copyright © 2018 by the Korea Genome Organization

© It is identical to the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>).

of the brain and olfactory epithelium, much effort has been made to confirm the expression of ORs in the brain. MOL2.3, called Olf78, was reported to be expressed in the ganglia of the autonomic nervous system [16]. Some mouse ORs—M71 (Olf151), C6 (Olf49), and OR3—are detected in the cerebral cortex and might play a role in developmental processes, such as axon guidance and target recognition during the postnatal period [17]. OR expression (OR1E1, OR2J3, OR2L13, OR11H1, and OR52L1) in the frontal cortex is downregulated in Parkinson disease (PD) patients [18].

The expression of a mouse OR, Olf110, was detected in the cerebral cortex of wild-type mice for 1 postnatal year, and its mRNA levels did not vary during that period [19]. It was observed that 8 ORs and 6 taste receptors (TASRs) are constitutively expressed in the frontal cortex, entorhinal cortex, and cerebellum in human control brains. These data imply that variable dysregulation of certain ORs and TASRs is common in several neurodegenerative diseases, including Alzheimer disease, Progressive Supranuclear Palsy, and Creutzfeldt-Jakob disease. The same research group also reported a decrease in olfactory and TASR expression in the dorsolateral prefrontal cortex in chronic schizophrenia [20]. There was a report that ORs are expressed in mouse mesencephalic dopaminergic (mDA) neurons. After screening ligands on Olf287, carvone enantiomers were identified as agonists of Olf287 and able to increase intracellular  $Ca^{2+}$  in solitary mDA neurons. The ORs were found to be expressed in human Substantia Nigra and downregulated in PD postmortem brains [21].

Gong *et al.* [22] found that some ORs are differentially expressed in the sciatic nerve and dorsal root ganglia after sciatic nerve injury in rats. The expression and expression profile of several ORs in the sciatic nerve were verified, and they also observed that the expression of some ORs in primary cultures of Schwann cells was upregulated under  $H_2O_2$  stimulation [22]. Recently, there has been a report that Olf544 is expressed in mouse brain and heart, as well as nose, adipose tissue, and spleen [23].

### Heart-related system

In 1995, Drutel *et al.* [9] reported the expression of OLI (Olr1654) in the developing rat heart and suggested its involvement in cardiac morphogenesis, where the OR was hardly detectable at the adult stages. As mentioned above, murine PSGRs (Olf78 and Olf59) are expressed in the hearts of mouse [16] and rat [15], in addition to prostate. In 2007, Zhang *et al.* [24] reported that ORs (including OR10G4) are expressed in human heart and various organs using reverse transcription-polymerase chain reaction (RT-PCR) and microarray.

Recently, OR10J5 was found to be expressed in human

aorta, coronary artery, and human umbilical vein endothelial cells [25]. In addition, Matrigel plug assay showed that lral enhances angiogenesis *in vivo*. These results let us surmise the physiological role of OR10J5 in angiogenesis.

Chang *et al.* [26] showed that a short-chain fatty acid OR, Olf78, is highly and selectively expressed in oxygen-sensitive glomus cells of the carotid body (CB). The CB is a chemosensory organ that monitors blood oxygen to control breath [27]. As lactate eventually activates Olf78 in heterologous expression experiments, Chang *et al.* [26] proposed that Olf78 may act as a hypoxia sensor in the breathing circuit, in addition to its role in olfaction. The next year, Zhou *et al.* [28] applied the single-cell RNA-Seq method to eliminate the contamination of genes derived from other cell types present in the CB to analyze the expression of ORs in CB glomus cells and identified Olf78 as the most highly abundant OR in the CB in mice. More recently, Jovancevic *et al.* [29] reported that OR51E1 activation by the application of cognate agonists induces a negative chronotropic effect in human stem cell-derived cardiomyocytes and also provokes negative inotropic activity in cardiac trabeculae and slice preparations of human explanted ventricles. These results imply that some ORs that are expressed in heart may be therapeutic targets for the metabolic regulation of cardiac function. A mouse OR—Olf544, an ortholog of the OR52 family (OR52K1)—was reported to be expressed in the heart of mice [23]. The unexpected cardiac expression of ORs may regulate cell growth and morphogenesis.

General studies on OR expression show that ORs are expressed throughout systems and have diverse functions throughout the body of animals (Table 1) [7-11, 13-26, 28-73].

## System-Wide Expression of Olfactory Receptors

### Testis

In 1992, ectopic expression of OR-like proteins in dog testis was reported after the OR multigene family was originally isolated from rat olfactory epithelium [7]. Subsequently, many experiments were performed to trace the presence of ORs in animal testicular tissues for the investigation of OR function in sperm-oocyte chemotaxis, where it was thought that ORs should have chemotactic roles during fertilization. Moreover, ORs are also thought to be involved in chemotaxis [74] and cytokinesis [75], and dozens of ORs were found to be expressed in the testis of mammals using deep sequencing [32]. Several tens of ORs were expressed in testicular tissues of mammals, and their roles in sperm-oocyte chemotaxis were investigated. Very

recently, Milardi *et al.* [33] suggested that the presence of ORs, such as OR4S1, OR4C13, and OR111, on mature cells might be related to acrosome activity and sperm motility after

a proteomic analysis. Itakura *et al.* [34] also identified  $G\alpha_{olf}$  and several kinds of ORs from rat placenta using RT-PCR, western blotting, and immunochemical methods. They

**Table 1.** Selected list of olfactory receptors expressed in system-wide tissues

Tissue	Organism	Representative OR(s)	Remarks (methods, DB, ligand)	Reference
Brain	Rat	RA1c (Olr59)	PCR, NB, ISH	[13]
	Mouse	MOL2.3 (RA1c, Olfr78)	PCR, $\beta$ -gal staining, ISH	[14]
	Mouse	MOL2.3 (Olf78)	$\beta$ -gal staining, ISH	[16]
	Mouse	M71 (olfr151), C6 (Olf49), (OR3)	Cerebral cortex	[17]
	Mouse	Olf110	AD, PSP, CJD, RT-PCR, cerebral cortex	[19]
	Human	ORs ( $\downarrow$ )	Frontal cortex, PD, microarray	[18]
	Mouse	Olf287, etc.	mDA cell, RT-PCR, ISH, FANTOM5	[21]
	Human	OR51E1, OR52L1 ( $\downarrow$ )	DPC, schizophrenia, PCR	[20]
	Mouse	Olf544	RT-PCR, qPCR	[23]
Heart-related	Rat	OL1 (Olr1654)	PCR, ISH	[9]
	Mouse, rat	PSGR homolog: RA1c (Olf78, Olf59)	NB	[15]
	Mouse	MOL2.3 (Olf78)	$\beta$ -gal staining, ISH	[16]
	Human	ORs	Microarray	[24]
	Mouse	Olf78	RT-PCR, $\beta$ -gal staining, luciferase assay	[30]
	Human	OR10J5	Human aorta, coronary artery, HUVEC	[25]
	Mouse	Olf78	Carotid body, immunostaining	[26]
	Rat	ORs	Sciatic nerve, dorsal root ganglia	[22]
	Mouse	Olf78	Carotid body, single-cell RNA-Seq	[28]
Human	OR51E1, OR2W3, OR51E2, etc.	TRXM analysis, RT-PCR, WB	[29]	
Mouse	Olf544	RT-PCR, qPCR	[23]	
Testis	Dog	HGMP07J/I (OR10J1), DTMT	PCR, NB	[7]
	Dog	DTMT	Antibody	[8]
	Rat	OL1 (Olr1654)	PCR, ISH	[9]
	Rat	OR2	Antibody	[31]
	Human	OR4N4, OR2W3, OR3A2, OR10J1, etc.	RNA-Seq	[32]
	Human	OR4S1, OR111, OR4C13	Testicular tissues, MS, WB, CA	[33]
Placenta	Rat	ORs	RT-PCR, WB, IH	[34]
Development	Mouse	Olf603	Embryonic tissue, IF	[35]
Erythroid	Human	HPFH1OR (OR52A1)	RT-PCR, erythroid (blood), all stages	[11]
Tongue	Rat	ORs	RT-PCR	[36]
	Human	ORs	RT-PCR	[37]
	Human	ORs	RT-PCR	[38]
	Human, mouse	ORs	RT-PCR, bioinformatics	[39]
GI tract	Mouse	PSGR (Olf78)	NB	[15]
	Human	4 ORs	RT-PCR, IH, amperometry	[40]
	Human	OR51E1	Microarray, RT-PCR, IH	[41]
	Pig	OR51E1	RT-qPCR, IH	[42]
	Human, mouse	OR1A1, OR1G1, OR51E1, Olf43	RT-qPCR, siRNA	[43]
Pancreas	Rat	OL2 (Olr857)	PCR	[10]
	Rat	Olr984 ( $\downarrow$ )	RT-PCR	[44]
	Rat	Orls ( $\downarrow$ )	Microarray, bioinformatics	[45]
	Mouse	Olf544	IC, IF, RT-PCR, RNA-Seq, WB	[46]
Spleen	Rat	OL2 (Olr857)	MIN6 cell line	[10]
	Rat	OdRs	Degenerate primers	[47]
	Human, mouse, rat	OR51E2, Olf78, Olf59	NB	[15]
Liver	Human, mouse, rat	OR51E2, Olf78, Olf59	Hepatocyte	[15]
	Human	OR1A1	PCR, IB, IH	[48]
	Human, mouse	OR10J5, Olf16	PCR, WB, siRNA	[49]
	Mouse	Olf544	RT-PCR, qPCR	[23]

Table 1. Continued

Tissue	Organism	Representative OR(s)	Remarks (methods, DB, ligand)	Reference
Kidney	Human	ORs	Microarray	[24]
	Mouse	Olf78, Olf90, Olf1373, Olf1392, Olf1393	RT-PCR, WB, IH	[50]
	Mouse	Olf78	RT-PCR, $\beta$ -gal staining, luciferase assay	[30]
	Human	OR51E1, OR11H7	RT-PCR, WB, IC, IH	[51]
	Mouse	Olf1393	RT-PCR, IH, IB	[52]
Lung	Mouse	MOL2.3 (Olf78)	$\beta$ -gal staining, ISH	[16]
	Human	OR51E1	RT-PCR, IH	[53]
	Human	OR2J3	NSCLC	[54]
Skin	Human	OR2AT4	Calcium imaging, wound scratch assay	[55]
	Human	OR51E2	RT-PCR, WB, IC	[56]
Muscle	Mouse	Olf16 (MOR23)	RT-PCR, IB	[57]
	Mouse	Olf16 (MOR23)	RT-PCR, IB	[58]
	Human	OR51E2	RT-PCR, CRISPR-Cas9, WB	[59]
	Human	OR51E1, OR11H7	RT-PCR, WB, IC, IH	[51]
Prostate	Human, mouse, rat	OR51E2, Olf78, Olf59	NB	[15]
	Human	OR51E2	RT-PCR, IH	[60]
Cancers	Human	PSGR (OR51E2)	CaP, RT-PCR	[61]
	Human	PSGR	CaP, RT-PCR	[62]
	Human	PSGR2 (OR51E1)	CaP, RT-PCR, NB	[63]
	Human	OR51E2	CaP, RT-PCR, IH	[60]
	Human	OR1D2 (OR17-4)	CaP, RT-PCR, CLSM, FACS	[64]
	Human	PSGR	LNCaP, RT-PCR, calcium imaging	[65]
	Human	PSGR	LNCaP, MS, IB, calcium imaging	[66]
	Human	OR51E1	SI-NEC	[67]
	Human	OR1A2	Hepatocarcinoma	[68]
	Human	OR51B5	Leukemia	[69]
	Human	OR7C1	CRC-CIC	[70]
	Human	OR3A4	Gastric cancer	[71]
	Human	OR2AT4	Myelogenous leukemia	[72]
	Human	ORs	Leukemia	[73]
	Human	OR2J3	NSCLC	[54]

OR, olfactory receptors; NB, northern blotting; ISH, *in situ* hybridization;  $\beta$ -gal staining,  $\beta$ -galactosidase staining; AD, Alzheimer disease; PSP, progressive supranuclear palsy; CJD, Creutzfeldt-Jakob disease; RT-PCR, reverse transcription polymerase chain reaction; PD, Parkinson disease; mDA, mesencephalic dopaminergic; DPC, dorsolateral prefrontal cortex; qPCR, quantitative polymerase chain reaction; PSGR, prostate-specific G protein-coupled receptor; HUVEC, human umbilical vein endothelial cells; RNA-Seq, RNA sequencing; WB, western blotting; MS, mass spectrometry; CA, confocal analysis; IH, immunohistochemistry; IF, immunofluorescence; GI, gastrointestinal; RT-qPCR, reverse transcription quantitative polymerase chain reaction; IC, immunocytochemistry; IB, immunoblotting; NSCLC, non-small-cell lung cancer; CaP, carcinoma of prostate; CLSM, confocal laser scanning microscopy; FACS, fluorescence-activated cell sorting; LNCaP, lymph node carcinoma of the prostate; SI-NEC, small intestine neuroendocrine carcinoma; CRC-CIC, colorectal cancer cancer-initiating cell.

suggested that ORs may recognize some small molecules in the placenta as environmental signals and mediate the growth of the fetus.

### Tongue and gastrointestinal tract

Since three transcripts of OR were identified in rat taste organs and reproductive tissues using an RT-PCR strategy [36], ORs, as well as TASRs, were isolated in the tongue of human [37, 38] and mouse [39]. In mouse, northern analysis revealed that the homolog of PSGR (Olf78), an OR, was predominantly expressed in colon [15]. In the case of human, the expression of four ORs, including OR73

(OR5D18), hOR17-7/11 (OR1A1), OR1G1, and hOR17-210 (OR1E3), in the enterochromaffin cells of human gut was confirmed [40]. Amperometric detection using  $\text{Ca}^{2+}$  imaging and immunostaining studies revealed that the ligands of those ORs consequently caused serotonin release. Another human OR, OR51E1, was identified from enterochromaffin cells in the jejunum and ileum of neuroendocrine carcinomas [41]. In 2015, the porcine homolog of OR51E1 was also found to be expressed along the gastrointestinal (GI) tract of pigs and modulated by intestinal microbiota [42]. Recently, human ORs (OR1A1, OR1G1, and OR51E1) and the mouse homolog of OR1A1 (Olf43) were found to

participate in glucose homeostasis during meal ingestion by inducing the secretion of gut peptides [43].

According to recent reports, ORs are expressed in non-olfactory tissues, including brain, heart, testis, tongue, GI tract, pancreas [46], spleen [15], liver [23], kidney [52], lung [53], skin [56], muscle [58], and diaphragm [30], and even cancerous tissues, such as prostate [64] and lung cancer cells [54] (Table 1).

In human primary keratinocytes, OR2AT4 is expressed and induces wound-healing processes by increasing cell proliferation and migration and regeneration of keratinocyte monolayers [55]. In this mechanism, the activation of OR2AT4 induces the phosphorylation of extracellular signal-regulated kinases (Erk1/2) and p38 mitogen-activated protein kinase (MAPK) by a cAMP-dependent pathway. This type of investigation should be applied in various fields of OR research. Another human OR, OR51E2, was detected in human epidermal melanocytes at the transcript and protein level, and its stimulation with its cognate ligand,  $\beta$ -ionone, significantly slows melanocyte proliferation [56]. Additional results also demonstrated that OR51E2 activation elevates cytosolic  $Ca^{2+}$  and cAMP using RNA silencing and receptor antagonists. These findings suggest that OR signaling can influence melanocyte homeostasis. We surmise that ORs play a pivotal role in maintaining organismal homeostasis according to results above.

## Olfactory Receptors in Cancer

After an OR was found to be overexpressed in human cancerous prostate and identified as a PSGR (OR51E2) [61], PSGR (OR51E2) [62], and PSGR2 (OR51E1) [63] were proven to be differentially expressed between prostate cells in malignant and benign tissues. In 2009, Neuhaus *et al.* [60] identified androstenone derivatives as ligands for recombinant OR51E2. Activation of endogenous OR51E2 by  $\beta$ -ionone resulted in the inhibition of prostate cancer cell proliferation. The mechanism of p38, including its trigger (Pky2) and a tumor suppressor protein (N-myc downstream regulated gene 1), was suggested later [65, 66]. Also, the expression of OR51E1 in microdissected small intestine neuroendocrine carcinoma cells was found to be higher than in adjacent microenvironment cells [67].

In the case of ORs other than PSGRs, mRNA overexpression of OR1D2 was confirmed in LNCap prostate carcinoma cells compared with non-prostate-derived cell lines using RT-PCR, confocal laser scanning microscopy, and fluorescence-activated cell sorting [64].

Recently, a kind of terpene, the major constituent of essential oils, was observed as an effector in cellular proliferation in hepatocarcinoma cells [68]. A human OR,

OR1A2, was detected at the mRNA and protein levels and demonstrated its potential involvement in (-)-citronellal-induced calcium signaling in the Huh7hepatocellular carcinoma cell line. Massberg *et al.* reported that the activation of OR1A2 results in the phosphorylation of p38 MAPK and reduces cell proliferation. More recently, Kalbe *et al.* [54] reported that a human OR, OR2J3, in non-small-cell lung cancer (NSCLC) cells induces apoptosis and inhibits cell proliferation and migration with its cognate ligand, helional. NSCLC is known to be resistant to common chemotherapy, and approaches via OR pathways could be helpful for the treatment of NSCLC with the precedence of further in-depth studies. Along with the cases mentioned above, diverse types of cancer appear to have various ORs, according to their occurring tissues (Table 1).

## Perspectives

Regardless of their discovery history, it would be better to have an extended view of ORs to get objective analysis results. Besides OR expression, its transportation to the cytoplasmic membrane of a cell is indispensable for correct function. Recently, Sharma *et al.* [76] reported a study on the relation between OR protein trafficking and OR transcriptional regulation using an *Rtp1* and *Rtp2* double-knockout mouse system; thus, a more sophisticated study of OR trafficking should be complemented.

In case of the cancer research, some identified ORs could serve as potential therapeutic targets for cancer diagnosis and its therapeutic applications. All biological phenomena, including cancers, are composite networks. Although much research has been done and is ongoing with regard to ORs, the relationship of the results is not clear. However, it has been proven that molecular functions and physiological changes in various parts could be caused by OR ligands in animal system. Therefore, the future task will be to identify at a holistic level the extent to which OR is involved in overall physiological phenomena *in vivo*. More in-depth and comprehensive studies on an OR production and delivery system from the viewpoint of chemosensory machinery are needed for the research of animal physiology and the application of OR-related mechanisms. Another additional point to consider is that the OR family is one of the most dynamic gene populations among individuals [77]. In order to discover new facts about the function of ORs, environmental issues, as well as internal factors, must be considered. The report that bats have a unique and diverse OR gene repertoire deserves consideration [78].

To advance the study of the various functions and roles of ORs, the modularity mechanism that acts between the receptor and its wide-ranged ligands should be addressed.

**ORCID:** S. June Oh: <http://orcid.org/0000-0001-7607-870X>

## References

- Vassilatis DK, Hohmann JG, Zeng H, Li F, Ranchalis JE, Mortrud MT, *et al.* The G protein-coupled receptor repertoires of human and mouse. *Proc Natl Acad Sci U S A* 2003;100:4903-4908.
- Fredriksson R, Lagerström MC, Lundin LG, Schiöth HB. The G-protein-coupled receptors in the human genome form five main families: phylogenetic analysis, paralogon groups, and fingerprints. *Mol Pharmacol* 2003;63:1256-1272.
- Downes GB, Gautam N. The G protein subunit gene families. *Genomics* 1999;62:544-552.
- Syrovatkina V, Alegre KO, Dey R, Huang XY. Regulation, signaling, and physiological functions of G-proteins. *J Mol Biol* 2016;428:3850-3868.
- Buck L, Axel R. A novel multigene family may encode odorant receptors: a molecular basis for odor recognition. *Cell* 1991; 65:175-187.
- Jones DT, Reed RR. Golf: an olfactory neuron specific-G protein involved in odorant signal transduction. *Science* 1989;244: 790-795.
- Parmentier M, Libert F, Schurmans S, Schiffmann S, Lefort A, Eggerickx D, *et al.* Expression of members of the putative olfactory receptor gene family in mammalian germ cells. *Nature* 1992;355:453-455.
- Vanderhaeghen P, Schurmans S, Vassart G, Parmentier M. Olfactory receptors are displayed on dog mature sperm cells. *J Cell Biol* 1993;123(6 Pt 1):1441-1452.
- Drutel G, Arrang JM, Diaz J, Wisnewsky C, Schwartz K, Schwartz JC. Cloning of OL1, a putative olfactory receptor and its expression in the developing rat heart. *Receptors Channels* 1995;3:33-40.
- Blache P, Gros L, Salazar G, Bataille D. Cloning and tissue distribution of a new rat olfactory receptor-like (OL2). *Biochem Biophys Res Commun* 1998;242:669-672.
- Feingold EA, Penny LA, Nienhuis AW, Forget BG. An olfactory receptor gene is located in the extended human beta-globin gene cluster and is expressed in erythroid cells. *Genomics* 1999; 61:15-23.
- Feldmesser E, Olender T, Khen M, Yanai I, Ophir R, Lancet D. Widespread ectopic expression of olfactory receptor genes. *BMC Genomics* 2006;7:121.
- Raming K, Konzelmann S, Breer H. Identification of a novel G-protein coupled receptor expressed in distinct brain regions and a defined olfactory zone. *Receptors Channels* 1998;6:141-151.
- Conzelmann S, Levai O, Bode B, Eisel U, Raming K, Breer H, *et al.* A novel brain receptor is expressed in a distinct population of olfactory sensory neurons. *Eur J Neurosci* 2000;12: 3926-3934.
- Yuan TT, Toy P, McClary JA, Lin RJ, Miyamoto NG, Kretschmer PJ. Cloning and genetic characterization of an evolutionarily conserved human olfactory receptor that is differentially expressed across species. *Gene* 2001;278:41-51.
- Weber M, Pehl U, Breer H, Strotmann J. Olfactory receptor expressed in ganglia of the autonomic nervous system. *J Neurosci Res* 2002;68:176-184.
- Otaki JM, Yamamoto H, Firestein S. Odorant receptor expression in the mouse cerebral cortex. *J Neurobiol* 2004;58: 315-327.
- Garcia-Esparcia P, Schlüter A, Carmona M, Moreno J, Ansoleaga B, Torrejón-Escribano B, *et al.* Functional genomics reveals dysregulation of cortical olfactory receptors in Parkinson disease: novel putative chemoreceptors in the human brain. *J Neuropathol Exp Neurol* 2013;72:524-539.
- Ansoleaga B, Garcia-Esparcia P, Llorens F, Moreno J, Aso E, Ferrer I. Dysregulation of brain olfactory and taste receptors in AD, PSP and CJD, and AD-related model. *Neuroscience* 2013; 248:369-382.
- Ansoleaga B, Garcia-Esparcia P, Pinacho R, Haro JM, Ramos B, Ferrer I. Decrease in olfactory and taste receptor expression in the dorsolateral prefrontal cortex in chronic schizophrenia. *J Psychiatr Res* 2015;60:109-116.
- Grison A, Zucchelli S, Urzì A, Zamparo I, Lazarevic D, Pascarella G, *et al.* Mesencephalic dopaminergic neurons express a repertoire of olfactory receptors and respond to odorant-like molecules. *BMC Genomics* 2014;15:729.
- Gong L, Chen Q, Gu X, Li S. Expression and identification of olfactory receptors in sciatic nerve and dorsal root ganglia of rats. *Neurosci Lett* 2015;600:171-175.
- Wu C, Hwang SH, Jia Y, Choi J, Kim YJ, Choi D, *et al.* Olfactory receptor 544 reduces adiposity by steering fuel preference toward fats. *J Clin Invest* 2017;127:4118-4123.
- Zhang X, De la Cruz O, Pinto JM, Nicolae D, Firestein S, Gilad Y. Characterizing the expression of the human olfactory receptor gene family using a novel DNA microarray. *Genome Biol* 2007;8:R86.
- Kim SH, Yoon YC, Lee AS, Kang N, Koo J, Rhyu MR, *et al.* Expression of human olfactory receptor 10J5 in heart aorta, coronary artery, and endothelial cells and its functional role in angiogenesis. *Biochem Biophys Res Commun* 2015;460:404-408.
- Chang AJ, Ortega FE, Riegler J, Madison DV, Krasnow MA. Oxygen regulation of breathing through an olfactory receptor activated by lactate. *Nature* 2015;527:240-244.
- Kumar P, Prabhakar NR. Peripheral chemoreceptors: function and plasticity of the carotid body. *Compr Physiol* 2012;2:141-219.
- Zhou T, Chien MS, Kaleem S, Matsunami H. Single cell transcriptome analysis of mouse carotid body glomus cells. *J Physiol* 2016;594:4225-4251.
- Jovancevic N, Dendorfer A, Matzkies M, Kovarova M, Heckmann JC, Osterloh M, *et al.* Medium-chain fatty acids modulate myocardial function via a cardiac odorant receptor. *Basic Res Cardiol* 2017;112:13.
- Pluznick JL, Protzko RJ, Gevorgyan H, Peterlin Z, Sipos A, Han J, *et al.* Olfactory receptor responding to gut microbiota-derived signals plays a role in renin secretion and blood pressure regulation. *Proc Natl Acad Sci U S A* 2013;110:4410-4415.
- Walensky LD, Roskams AJ, Lefkowitz RJ, Snyder SH, Ronnett GV. Odorant receptors and desensitization proteins colocalize in mammalian sperm. *Mol Med* 1995;1:130-141.
- Flegel C, Manteniotis S, Osthold S, Hatt H, Gisselmann G.

- Expression profile of ectopic olfactory receptors determined by deep sequencing. *PLoS One* 2013;8:e55368.
33. Milardi D, Colussi C, Grande G, Vincenzoni F, Pierconti F, Mancini F, *et al.* Olfactory receptors in semen and in the male tract: from proteome to proteins. *Front Endocrinol (Lausanne)* 2017;8:379.
  34. Itakura S, Ohno K, Ueki T, Sato K, Kanayama N. Expression of Golf in the rat placenta: possible implication in olfactory receptor transduction. *Placenta* 2006;27:103-108.
  35. Baker NL, Miller KA, Newgreen DF, Farlie PG. Olfr603, an orphan olfactory receptor, is expressed in multiple specific embryonic tissues. *Gene Expr Patterns* 2015;19:30-35.
  36. Thomas MB, Haines SL, Akeson RA. Chemoreceptors expressed in taste, olfactory and male reproductive tissues. *Gene* 1996;178:1-5.
  37. Gaudin JC, Breuils L, Haertlé T. New GPCRs from a human lingual cDNA library. *Chem Senses* 2001;26:1157-1166.
  38. Durzyński L, Gaudin JC, Myga M, Szydlowski J, Goździcka-Józefiak A, Haertlé T. Olfactory-like receptor cDNAs are present in human lingual cDNA libraries. *Biochem Biophys Res Commun* 2005;333:264-272.
  39. Gaudin JC, Breuils L, Haertlé T. Mouse orthologs of human olfactory-like receptors expressed in the tongue. *Gene* 2006;381:42-48.
  40. Braun T, Volland P, Kunz L, Prinz C, Gratzl M. Enterochromaffin cells of the human gut: sensors for spices and odorants. *Gastroenterology* 2007;132:1890-1901.
  41. Leja J, Essaghir A, Essand M, Wester K, Oberg K, Tötterman TH, *et al.* Novel markers for enterochromaffin cells and gastrointestinal neuroendocrine carcinomas. *Mod Pathol* 2009;22:261-272.
  42. Priori D, Colombo M, Clavanzani P, Jansman AJ, Lallès JP, Trevisi P, *et al.* The olfactory receptor OR51E1 is present along the gastrointestinal tract of pigs, co-localizes with enteroendocrine cells and is modulated by intestinal microbiota. *PLoS One* 2015;10:e0129501.
  43. Kim KS, Lee IS, Kim KH, Park J, Kim Y, Choi JH, *et al.* Activation of intestinal olfactory receptor stimulates glucagon-like peptide-1 secretion in enteroendocrine cells and attenuates hyperglycemia in type 2 diabetic mice. *Sci Rep* 2017;7:13978.
  44. Choi SA, Suh HJ, Yun JW, Choi JW. Differential gene expression in pancreatic tissues of streptozocin-induced diabetic rats and genetically-diabetic mice in response to hypoglycemic dipeptide cyclo (His-Pro) treatment. *Mol Biol Rep* 2012;39:8821-8835.
  45. Ng SF, Lin RC, Maloney CA, Youngson NA, Owens JA, Morris MJ. Paternal high-fat diet consumption induces common changes in the transcriptomes of retroperitoneal adipose and pancreatic islet tissues in female rat offspring. *FASEB J* 2014;28:1830-1841.
  46. Kang N, Bahk YY, Lee N, Jae Y, Cho YH, Ku CR, *et al.* Olfactory receptor Olfr544 responding to azelaic acid regulates glucagon secretion in alpha-cells of mouse pancreatic islets. *Biochem Biophys Res Commun* 2015;460:616-621.
  47. Walensky LD, Ruat M, Bakin RE, Blackshaw S, Ronnett GV, Snyder SH. Two novel odorant receptor families expressed in spermatids undergo 5'-splicing. *J Biol Chem* 1998;273:9378-9387.
  48. Wu C, Jia Y, Lee JH, Kim Y, Sekharan S, Batista VS, *et al.* Activation of OR1A1 suppresses PPAR-gamma expression by inducing HES-1 in cultured hepatocytes. *Int J Biochem Cell Biol* 2015;64:75-80.
  49. Tong T, Ryu SE, Min Y, de March CA, Bushdid C, Golebiowski J, *et al.* Olfactory receptor 10J5 responding to alpha-cedrene regulates hepatic steatosis via the cAMP-PKA pathway. *Sci Rep* 2017;7:9471.
  50. Pluznick JL, Zou DJ, Zhang X, Yan Q, Rodriguez-Gil DJ, Eisner C, *et al.* Functional expression of the olfactory signaling system in the kidney. *Proc Natl Acad Sci U S A* 2009;106:2059-2064.
  51. Kalbe B, Schlimm M, Wojcik S, Philippou S, Maßberg D, Jansen F, *et al.* Olfactory signaling components and olfactory receptors are expressed in tubule cells of the human kidney. *Arch Biochem Biophys* 2016;610:8-15.
  52. Shepard BD, Cheval L, Peterlin Z, Firestein S, Koepsell H, Doucet A, *et al.* A renal olfactory receptor aids in kidney glucose handling. *Sci Rep* 2016;6:35215.
  53. Giandomenico V, Cui T, Grimelius L, Öberg K, Pelosi G, Tsolakis AV. Olfactory receptor 51E1 as a novel target for diagnosis in somatostatin receptor-negative lung carcinoids. *J Mol Endocrinol* 2013;51:277-286.
  54. Kalbe B, Schulz VM, Schlimm M, Philippou S, Jovancevic N, Jansen F, *et al.* Helional-induced activation of human olfactory receptor 2J3 promotes apoptosis and inhibits proliferation in a non-small-cell lung cancer cell line. *Eur J Cell Biol* 2017;96:34-46.
  55. Busse D, Kudella P, Gruning NM, Gisselmann G, Ständer S, Luger T, *et al.* A synthetic sandalwood odorant induces wound-healing processes in human keratinocytes via the olfactory receptor OR2AT4. *J Invest Dermatol* 2014;134:2823-2832.
  56. Gelis L, Jovancevic N, Veitinger S, Mandal B, Arndt HD, Neuhaus EM, *et al.* Functional Characterization of the Odorant Receptor 51E2 in Human Melanocytes. *J Biol Chem* 2016;291:17772-17786.
  57. Griffin CA, Kafadar KA, Pavlath GK. MOR23 promotes muscle regeneration and regulates cell adhesion and migration. *Dev Cell* 2009;17:649-661.
  58. Pichavant C, Burkholder TJ, Pavlath GK. Decrease of myofiber branching via muscle-specific expression of the olfactory receptor mOR23 in dystrophic muscle leads to protection against mechanical stress. *Skelet Muscle* 2016;6:2.
  59. Aisenberg WH, Huang J, Zhu W, Rajkumar P, Cruz R, Santhanam L, *et al.* Defining an olfactory receptor function in airway smooth muscle cells. *Sci Rep* 2016;6:38231.
  60. Neuhaus EM, Zhang W, Gelis L, Deng Y, Noldus J, Hatt H. Activation of an olfactory receptor inhibits proliferation of prostate cancer cells. *J Biol Chem* 2009;284:16218-16225.
  61. Xu LL, Stackhouse BG, Florence K, Zhang W, Shanmugam N, Sesterhenn IA, *et al.* PSGR, a novel prostate-specific gene with homology to a G protein-coupled receptor, is overexpressed in prostate cancer. *Cancer Res* 2000;60:6568-6572.
  62. Xu LL, Sun C, Petrovics G, Makarem M, Furusato B, Zhang W,

- et al.* Quantitative expression profile of PSGR in prostate cancer. *Prostate Cancer Prostatic Dis* 2006;9:56-61.
63. Weng J, Wang J, Hu X, Wang F, Ittmann M, Liu M. PSGR2, a novel G-protein coupled receptor, is overexpressed in human prostate cancer. *Int J Cancer* 2006;118:1471-1480.
64. Sturzu A, Sheikh S, Echner H, Nägele T, Deeg M, Schwentner C, *et al.* Novel bourgeonal fragrance conjugates for the detection of prostate cancer. *Invest New Drugs* 2013;31:1151-1157.
65. Sanz G, Leray I, Dewaele A, Sobilo J, Lerondel S, Bouet S, *et al.* Promotion of cancer cell invasiveness and metastasis emergence caused by olfactory receptor stimulation. *PLoS One* 2014;9:e85110.
66. Wiese H, Gelis L, Wiese S, Reichenbach C, Jovancevic N, Osterloh M, *et al.* Quantitative phosphoproteomics reveals the protein tyrosine kinase Pyk2 as a central effector of olfactory receptor signaling in prostate cancer cells. *Biochim Biophys Acta* 2015;1854:632-640.
67. Cui T, Tsolakis AV, Li SC, Cunningham JL, Lind T, Öberg K, *et al.* Olfactory receptor 51E1 protein as a potential novel tissue biomarker for small intestine neuroendocrine carcinomas. *Eur J Endocrinol* 2013;168:253-261.
68. Massberg D, Simon A, Haussinger D, Keitel V, Gisselmann G, Conrad H, *et al.* Monoterpene (-)-citronellal affects hepatocarcinoma cell signaling via an olfactory receptor. *Arch Biochem Biophys* 2015;566:100-109.
69. Manteniotis S, Wojcik S, Göthert JR, Dürig J, Dührsen U, Gisselmann G, *et al.* Deorphanization and characterization of the ectopically expressed olfactory receptor OR51B5 in myelogenous leukemia cells. *Cell Death Discov* 2016;2:16010.
70. Morita R, Hirohashi Y, Torigoe T, Ito-Inoda S, Takahashi A, Mariya T, *et al.* Olfactory receptor family 7 subfamily C member 1 is a novel marker of colon cancer-initiating cells and is a potent target of immunotherapy. *Clin Cancer Res* 2016;22:3298-3309.
71. Guo X, Yang Z, Zhi Q, Wang D, Guo L, Li G, *et al.* Long non-coding RNA OR3A4 promotes metastasis and tumorigenicity in gastric cancer. *Oncotarget* 2016;7:30276-30294.
72. Manteniotis S, Wojcik S, Brauhoff P, Möllmann M2, Petersen L, Göthert JR, *et al.* Functional characterization of the ectopically expressed olfactory receptor 2AT4 in human myelogenous leukemia. *Cell Death Discov* 2016;2:15070.
73. Jung H, Chae YC, Kim JY, Jeong OS, Kook H, Seo SB. Regulatory role of G9a and LSD1 in the transcription of olfactory receptors during leukaemia cell differentiation. *Sci Rep* 2017;7:46182.
74. Spehr M, Gisselmann G, Poplawski A, Riffell JA, Wetzel CH, Zimmer RK, *et al.* Identification of a testicular odorant receptor mediating human sperm chemotaxis. *Science* 2003;299:2054-2058.
75. Zhang X, Bedigian AV, Wang W, Eggert US. G protein-coupled receptors participate in cytokinesis. *Cytoskeleton (Hoboken)* 2012;69:810-818.
76. Sharma R, Ishimaru Y, Davison I, Ikegami K, Chien MS, You H, *et al.* Olfactory receptor accessory proteins play crucial roles in receptor function and gene choice. *Elife* 2017;6:e21895.
77. Olender T, Waszak SM, Viavant M, Khen M, Ben-Asher E, Reyes A, *et al.* Personal receptor repertoires: olfaction as a model. *BMC Genomics* 2012;13:414.
78. Hayden S, Bekaert M, Goodbla A, Murphy WJ, Dávalos LM, Teeling EC. A cluster of olfactory receptor genes linked to frugivory in bats. *Mol Biol Evol* 2014;31:917-927.