



Dental stem cells as a cell source for tissue engineering

Bong-Wook Park, D.D.S., M.S.D., Ph.D.^{1,2}

Section Editor of JKAOMS

¹Department of Dentistry, Gyeongsang National University School of Medicine, Jinju,

²Department of Oral and Maxillofacial Surgery, Changwon Gyeongsang National University Hospital, Changwon, Korea

Dental stem cells (DSCs) have been isolated from dental follicles, pulps, root apical papillae, and periodontal ligaments of extracted teeth, and all have exhibited multipotential, self-renewal, and mesenchymal stem cell (MSC) characteristics^{1,2}. However, DSCs from different sources vary in their differentiation properties: stem cells from dental follicles or root apical papillae of wisdom teeth showed superior osteogenic differentiation potential, whereas dental pulp stem cells demonstrated remarkably enhanced *in vitro* and *in vivo* neurogenic differentiation ability¹⁻³. Importantly, a new tissue cryopreservation protocol has developed for use as an autologous stem cell source⁴. In previous reports, MSCs were successfully isolated from long-term cryopreserved dental tissues (follicles, pulps, and root apical papillae)^{4,5}. MSCs from cryopreserved dental tissues showed identical characteristics to those from fresh dental tissues, including stemness, *in vitro* differentiation potential, and cell proliferation rate^{4,5}. This tissue cryopreservation method makes it possible to safely store dental tissues after tooth extraction for a long time, allowing them to be used as an autologous stem cell resource. Cells from autologous source may reduce unexpected side effects, such as various immune reactions, when they were *in vivo* transplanted.

In one previous report, MSCs from cryopreserved and fresh dental follicles showed excellent osteogenic regeneration potential and immunomodulatory properties through

suppression of the T-helper cell-mediated adaptive immune response, when they were transplanted *in vivo* transplanted into mandibular bone defects³. In addition, DSCs from follicles or pulps of extracted wisdom teeth successfully differentiated into cardiomyocyte- and hepatocyte-like cells *in vitro*^{5,6}. More importantly, dental pulp-derived stem cells (DPSCs) were distinguished as one of the most powerful cell sources for neurogenic differentiation *in vitro* and *in vivo*^{2,7,8}. Previously, DPSCs were shown to exhibit remarkable neurogenic differentiation potential, in terms of expression of neuron-specific markers and having higher Na⁺ and K⁺ current with the expression of synaptic markers². According to one hypothesis, the origin of dental pulp from the neural crest allows it to more easily differentiate into neuronal cells, compared to other cell lines⁷. Moreover, because of their immunomodulatory effect, DPSCs can activate microglia/astroglia in the host microenvironment, which will enhance the Wallerian degeneration and macrophage invasion, accelerating peripheral nerve regeneration². Interestingly, *in vivo* transplantation of DPSCs to treat sciatic nerve defects in experimental rats increases motor nerve function with histologically abundant regeneration of axonal fibers⁸. Moreover, recent work revealed that DPSCs exhibit higher cholinergic neuronal differentiation potential with acetylcholine production *in vitro* (unpublished data).

Taken together, stem cells from dental tissue, including pulps, apical papillae, and follicles, are shown to be excellent sources for tissue regeneration, with immunomodulatory properties. Stem cells from root apical papillae and follicles demonstrated superior osteogenic differentiation and regeneration potential *in vitro* and *in vivo*. Whereas stem cells from dental pulp revealed excellent *in vitro* neurogenic differentiation and *in vivo* peripheral nerve regeneration potential. In addition, the new method for long-term cryopreservation of dental tissues after tooth extraction can safely preserve them for use as an autologous stem cell source for patients in need,

Bong-Wook Park

Department of Oral and Maxillofacial Surgery, Changwon Gyeongsang National University Hospital, 11 Samjeongja-ro, Seongsan-gu, Changwon 51472, Korea

TEL: +82-55-241-3885 FAX: +82-55-214-3265

E-mail: parkbw@gnu.ac.kr

ORCID: <https://orcid.org/0000-0002-2699-9188>

© This is an open-access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>), which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Copyright © 2018 The Korean Association of Oral and Maxillofacial Surgeons. All rights reserved.

which would create a new business model in dentistry.

Conflict of Interest

No potential conflict of interest relevant to this article was reported.

References

1. Song JH, Park BW, Byun JH, Kang EJ, Rho GJ, Shin SH, et al. Isolation and characterization of human dental tissue-derived stem cells in the impacted wisdom teeth: comparison of dental follicle, dental pulp, and root apical papilla-derived cells. *J Korean Assoc Oral Maxillofac Surg* 2010;36:186-96.
2. Ullah I, Subbarao RB, Kim EJ, Bharti D, Jang SJ, Park JS, et al. In vitro comparative analysis of human dental stem cells from a single donor and its neuronal differentiation potential evaluated by electrophysiology. *Life Sci* 2016;154:39-51.
3. Kang YH, Lee HJ, Jang SJ, Byun JH, Lee JS, Lee HC, et al. Immunomodulatory properties and in vivo osteogenesis of human dental stem cells from fresh and cryopreserved dental follicles. *Differentiation* 2015;90:48-58.
4. Park BW, Jang SJ, Byun JH, Kang YH, Choi MJ, Park WU, et al. Cryopreservation of human dental follicle tissue for use as a resource of autologous mesenchymal stem cells. *J Tissue Eng Regen Med* 2017;11:489-500.
5. Han YJ, Kang YH, Shivakumar SB, Bharti D, Son YB, Choi YH, et al. Stem cells from cryopreserved human dental pulp tissues sequentially differentiate into definitive endoderm and hepatocyte-like cells in vitro. *Int J Med Sci* 2017;14:1418-29.
6. Sung IY, Son HN, Ullah I, Bharti D, Park JM, Cho YC, et al. Cardiomyogenic differentiation of human dental follicle-derived stem cells by suberoylanilide hydroxamic acid and their in vivo homing property. *Int J Med Sci* 2016;13:841-52.
7. Huang GT, Gronthos S, Shi S. Mesenchymal stem cells derived from dental tissues vs. those from other sources: their biology and role in regenerative medicine. *J Dent Res* 2009;88:792-806.
8. Ullah I, Park JM, Kang YH, Byun JH, Kim DG, Kim JH, et al. Transplantation of human dental pulp-derived stem cells or differentiated neuronal cells from human dental pulp-derived stem cells identically enhances regeneration of the injured peripheral nerve. *Stem Cells Dev* 2017;26:1247-57.