

# Establishment of a $T_m$ -shift Method for Detection of Cat-Derived Hookworms

Yeqi Fu, Yunqiu Liu, Asmaa M.I. Abuzeid, Yue Huang, Xue Zhou, Long He, Qi Zhao, Xiu Li, Jumei Liu, Rongkun Ran, Guoqing Li\*

Guangdong Provincial Zoonosis Prevention and Control Key Laboratory, College of Veterinary Medicine, South China Agricultural University, Guangzhou 510542, P.R. China

**Abstract:** Melting temperature shift ( $T_m$ -shift) is a new detection method that analyze the melting curve on real-time PCR thermocycler using SYBR Green I fluorescent dye. To establish a  $T_m$ -shift method for the detection of *Ancylostoma ceylanicum* and *A. tubaeforme* in cats, specific primers, with GC tail of unequal length attached to their 5' end, were designed based on 2 SNP loci (ITS101 and ITS296) of the internal transcribed spacer 1 (ITS1) sequences. The standard curve of  $T_m$ -shift was established using the standard plasmids of *A. ceylanicum* (AceP) and *A. tubaeforme* (AtuP). The  $T_m$ -shift method stability, sensitivity, and accuracy were tested with reference to the standard curve, and clinical fecal samples were also examined. The results demonstrated that the 2 sets of primers based on the 2 SNPs could accurately distinguish between *A. ceylanicum* and *A. tubaeforme*. The coefficient of variation (CV) of  $T_m$ - values of AceP and AtuP was 0.07% and 0.06% in ITS101 and was 0.06% and 0.08% in ITS296, respectively. The minimum detectable DNA concentration was  $5.22 \times 10^{-6}$  and  $5.28 \times 10^{-6}$  ng/ $\mu$ l samples of AceP and AtuP, respectively. The accuracy of  $T_m$ -shift method reached 100% based on examination of 10 hookworm DNA samples with known species. In the clinical detection of hookworm in 69 stray cat fecal sample, the  $T_m$ -shift detection results were consistent with the microscopic examination and successfully differentiated between the 2-hookworm species. In conclusion, the developed method is a rapid, sensitive and accurate technique and can provide a promising tool for clinical detection and epidemiological investigation of cat-derived hookworms.

**Key words:** *Anoylostoma ceylanicum*, *Anoylostoma tubaeforme*, cat,  $T_m$ -shift, SNP, ITS1

## INTRODUCTION

Hookworms are common parasitic nematodes in the digestive tract of human beings, dogs and cats, which can cause a serious hazard to the host. The main hookworms infecting cats include *Ancylostoma ceylanicum*, *A. tubaeforme*, *A. braziliense*, and *Uncinaria stenocephala*. The recent investigations have shown that the cat's hookworm infection in China is mainly caused by *A. ceylanicum* and *A. tubaeforme* [1,2]. *A. ceylanicum* is globally distributed among dogs and cats, however it is particularly severe in Asia, especially in Southeast Asia and the Pacific. Recently, *A. ceylanicum* has been recorded in many South-east Asian countries, such as China [3,4], Japan [5], Malaysia [6], Laos [7], Thailand [8]. *A. tubaeforme* mainly infects cats,

but it is rarely reported in canines. According to the previous surveys, *A. tubaeforme* has been reported in cats in Australia [9,10], Brazil [11], Italy [12], Qatar [13], Spain [14], the United States [15,16] and China [2]. Currently, *A. ceylanicum* has become the second largest species of hookworm in Asia [17], that can feed on blood and develop into adults in the human body, causing anemia, abdominal pain, diarrhea, and even death. As the number of cat owners increases, it is important to establish a rapid and accurate method that can distinguish *A. ceylanicum* from *A. tubaeforme* to enhance public health and safety.

The fecal examination is a classical method for hookworm detection, but it fails to identify hookworm species. Therefore, it is difficult to judge whether the infection is caused by *A. ceylanicum*. Presently, many molecular detection techniques have been applied to identify hookworms, such as specific polymerase chain reaction (PCR), multiplex PCR, restriction fragment length polymorphism (RFLP) and high-resolution melting (HRM) analysis [18-20]. All the former techniques have certain limitations, some can only detect a small number of

•Received 19 November 2018, revised 25 January 2019, accepted 6 February 2019.

\*Corresponding author (gqli@scau.edu.cn)

© 2019, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

samples at the same time, and others are complicated and expensive to operate.

Melting temperature shift ( $T_m$ -shift) method based on single nucleotide polymorphism (SNP) is a new molecular detection technique, which is high-throughput, accurate and inexpensive. This method utilizes 2 allele-specific primers, each of which attach to one of the two SNP loci at its 3' end, a reverse primer that amplifies both alleles, and a fluorescent dye, SYBR® Green I (Molecular Probes, Eugene, Oregon, USA). Either one or both allele specific primer(s) amplify according to the sample genotype. GC rich tails of unequal length are attached to the 5' end of the allele-specific primers, so that the PCR product has a different melting temperature depending on which of the 2 primers is used. Therefore, genotypes can be distinguished by checking the melting curve on a real-time PCR instrument [21,22]. The  $T_m$ -shift method has been applied in many fields, such as medicine, microbiology, and aquatic biology [23-25]. In our laboratory, Pan et al. [26] established a  $T_m$ -shift genotyping method for *Giardia lamblia* assemblages A and F in cats, which pioneered a new genotyping method for veterinary parasitic protozoa. In addition, Fu et al. [27] developed a  $T_m$ -shift detection method for *A. ceylanicum* and *A. caninum* in dogs and applied it for the first time to identify veterinary parasitic nematode.

The present study aimed to develop and evaluate a  $T_m$ -shift method for the identification of *A. ceylanicum* and *A. tubaeforme* in cats, to provide a fast and accurate technique for the clinical detection and zoonotic risk assessment of cat-derived hookworms.

## MATERIALS AND METHODS

### Source of samples

Adult *A. ceylanicum* and *A. tubaeforme* samples were isolated and identified by Liu et al. [4] and Shi et al. [2], and preserved in 50% ethyl alcohol in our laboratory. Fecal samples were collected from stray cats at the Animal Rescue Station in Shaoguan city of Guangdong province in China and stored at 4°C.

### Genomic DNA extraction

The preserved adult hookworms were repeatedly washed with double-distilled water (ddH<sub>2</sub>O). After that, the genomic DNA was extracted using the Wizard® SV Genomic DNA Purification System (Promega, Guangzhou, China) according to the manufacturer's instructions, then stored at -20°C.

### PCR amplification of ITS1 sequence

The extracted DNA was subjected to polymerase chain reaction targeting *A. ceylanicum* and *A. tubaeforme* internal transcribed spacer 1 (ITS1) sequences using a forward primer AF (5'-CTTGTCTCGGGAAGGTTGG-3') and a reverse primer AR (5'-TTCACCACTCTAAGCGTCT-3') previously designed by Liu et al. [4]. The target amplification fragment length was 404 bp. PCR was carried out in a final volume of 25 µl, containing 12.5 µl Premix Ex-Taq polymerase (TaKaRa, Dalian, China), 9.5 µl ddH<sub>2</sub>O, 0.5 µl of each primer AF/AR (50 µmol/L), and 2 µl DNA template. The utilized PCR program was as follows: 94°C for 5 min, 35 × (94°C for 30 sec, 61.5°C for 30 sec and 72°C for 45 sec), and 72°C for 7 min. The PCR products were assessed by electrophoresis in agarose gel (1.5%) with ethidium bromide stain (0.2 mg/ml) and visualized on a UV transilluminator. The amplified fragment was recovered from the gel following the protocol of the gel recovery kit (Omega, Guangzhou, China) and kept at -20°C until further use.

### Preparation of standard plasmids

The purified PCR products were cloned in *Escherichia coli* and connected with pMD18-T vector (TaKaRa), then transformed into DH5 Competent Cells (TaKaRa). Positive clones were screened by bacterial PCR and sent to Sangon Biotech (Shanghai, China) for sequencing. The Plasmid Kit (Omega) was used to extract the plasmid DNAs. The plasmids harboring *A. ceylanicum* and *A. tubaeforme* ITS1 sequences were termed as AceP and AtuP. The plasmids (1 µL) were analyzed by the ultramicro-spectrophotometer to evaluate their purity and concentration, where A260/A280 value and optimal concentration should be 1.8-2.0 and 50 ng/µL, respectively.

### Establishment of $T_m$ -shift method based on SNP

Based on the 2 SNPs (ITS101 and ITS296), 2 sets of  $T_m$ -shift primers were designed using Primer Premier 5.0 software, referencing to the 2 sequences (MG733992, KP072069) downloaded from GenBank. The primer sequences and predicted amplification fragment length are demonstrated in Table 1. The above-mentioned primers were synthesized by Sangon Biotech (Shanghai, China). The synthesized primers were prepared in a template liquid at 100 pmol/µl and stored at -20°C. Then they were re-diluted to a final working concentration of 10 pmol/µl and stored at -20°C for use after partial shipments. PCR amplification and  $T_m$ -shift reaction were performed once in Rotor-Gene Q. The qPCR- $T_m$ -shift reaction mixture had the

**Table 1.** Primers for  $T_m$ -shift method based on 2 SNPs

Primer	Nucleotide sequence (5'-3')	Product length (bp)
ITS101AF (Atu)	<u>gattaccg</u> GGCGGCAGTGATTGCTGTA	
ITS101GF (Ace)	gcgggcagggcggg GGCGGCAGTGATTGCTGTG	106
ITS101R	GCCTAATGCTCAACCACCAACA	
ITS296TF (Ace)	<u>gattaccg</u> TTTGCAGAATCGTGACTTT	
ITS296AF (Atu)	gcgggcagggcggg TTTGCAGAATCGTGACTTA	127
ITS296R	TTCACCACTCTAAGCGTCT	

GC tails are underlined.

final volume of 20  $\mu$ l, containing 2  $\times$  SYBR<sup>®</sup> Premix Ex *Taq* TM II (10  $\mu$ l), long-tail primer (0.4  $\mu$ l), short-tail primer (0.4  $\mu$ l), common reverse primer (0.8  $\mu$ l), ddH<sub>2</sub>O (7.4  $\mu$ l), and plasmid (1.0  $\mu$ l). The cycling program was as follows: 95°C for 5 min, 40  $\times$  (95°C for 10 sec and 63°C for 30 sec). The melting process ranged from 70 to 95°C at the rate of 0.5°C/sec.

#### Stability, sensitivity and accuracy test

The stability of the established  $T_m$ -shift method was detected by repeating the experiments using 2 standard plasmids (AceP and AtuP). Intra-assay was checked 7 times, while inter-assay was checked 3 times for each plasmid. To evaluate the  $T_m$ -shift method sensitivity, AceP and AtuP were diluted 10 times and examined following the concentration of 1:10-1:10<sup>8</sup>. To assess the  $T_m$ -shift method accuracy, 10 samples with known hookworm species were examined. The qPCR- $T_m$ -shift reaction mixture and cycling program were identical to the mentioned above.

#### Clinical detection

Sixty-nine stray cat fecal samples were analyzed by zinc sulfate flotation technique and  $T_m$ -shift method. All fecal samples were suspended in ddH<sub>2</sub>O, heated for 5 cycles at 100°C for 5 min and immediately frozen at -80°C for 5 min. Genomic DNAs were extracted from the fecal samples using Stool DNA extraction kit (OMEGA, Guangzhou, China) as indicated by the manufacturer's protocols. The qPCR- $T_m$ -shift reaction mixture and cycling program were the same as the mentioned above. All results were further confirmed by DNA sequencing.

## RESULTS

#### Amplification of ITS1 fragment and preparation of standard plasmids

The amplification of the 2-hookworm species ITS1 sequences produced fragments with the length of 404 bp (Fig. 1), which agreed with the expected fragment length. The generat-


ed sequence data were submitted to GenBank (accession number: KF279132, JO812691). BLAST analysis showed the highest similarity (100%) with *A. ceylanicum* from Japan (LC036567) and the highest similarity (100%) with *A. tubaeforme* from Guangzhou (MG865903). Thus, 2 hookworms were identified as *A. ceylanicum* and *A. tubaeforme*. The A260/A280 values of positive plasmids of *A. ceylanicum* (AceP) and *A. tubaeforme* (AtuP) were between 1.8 and 2.0. In addition, the concentration of AceP and AtuP was 53.3 and 50.3 ng/ $\mu$ l, respectively, which was consistent with the optimal criteria.

#### Detection of qPCR- $T_m$ -shift


The  $T_m$ -shift standard curves based on the 2 SNPs (ITS101 and ITS296) for standard plasmid AceP and AtuP are revealed in Fig. 2. The result showed that the established  $T_m$ -shift method could distinguish *A. ceylanicum* (AceP) from *A. tubaeforme* (AtuP). The analysis of the standard curves by Rotor-Gene Q series software demonstrated that in the ITS101 primer, the  $T_m$  values of AceP and AtuP were 88.25°C and 86.50°C, respectively. In the ITS296 primer, the  $T_m$  values of AceP and AtuP were 85.00°C and 86.60°C, respectively.

#### Stability, sensitivity and accuracy

The stability test results are presented in Table 2. In the prim-



**Fig. 1.** PCR amplification of the ITS1 sequence of 2 hookworms. M, DL-500 DNA marker; 1, Positive control; 2, *A. ceylanicum*; 3, *A. tubaeforme*; 4, Negative control.



**Fig. 2.** Standard curves of  $T_m$ -shift for 2 hookworm standard plasmids based on ITS101 (A) and ITS296 (B). AceP: *A. ceylanicum* standard plasmid; AtuP: *A. tubaeforme* standard plasmid; Neg: negative control.

**Table 2.** Stability of  $T_m$ -shift method

Repeat	ITS101 ( $T_m$ ) (°C)		ITS296 ( $T_m$ ) (°C)	
	AtuP	AceP	AtuP	AceP
First (n=7)	86.49±0.09	88.25±0.10	86.63±0.07	85.01±0.09
Second (n=7)	86.47±0.07	88.21±0.06	86.60±0.10	85.04±0.06
Third (n=7)	86.44±0.09	88.26±0.11	86.59±0.11	85.00±0.10
Average (n=21)	86.47	88.24	85.00	86.60
CV (%)	0.06	0.07	0.08	0.06

er ITS101, the coefficient of variation (CV) of AceP and AtuP melting temperature ( $T_m$ ) was 0.07% and 0.06%, respectively. In primer ITS296, the CV of AceP and AtuP melting temperature was 0.06% and 0.08%, respectively. The sensitivity test results indicated that the  $T_m$ -shift method could still differentiate between *A. ceylanicum* (AceP) and *A. tubaeforme* (AtuP), when they were diluted to  $1:10^7$  ( $5.22 \times 10^{-6}$  ng/ $\mu$ l and  $5.28 \times 10^{-6}$  ng/ $\mu$ l, respectively) (Table 3). The results of examination of 10 DNA samples by  $T_m$ -shift method were identical to their known hookworm species with 100% accuracy (Table 4).

**Clinical detection**


Twelve out of 69 cat fecal samples were positive for hookworm by the fecal flotation and  $T_m$ -shift methods. Using  $T_m$ -shift method, 8 and 4 samples were identified as *A. ceylanicum*

**Table 3.** Sensitivity of  $T_m$ -shift method

Dilution	ITS101 ( $T_m$ ) (°C)		ITS296 ( $T_m$ ) (°C)	
	AtuP	AceP	AtuP	AceP
1: $10^1$	86.50	88.25	86.60	85.00
1: $10^2$	86.50	88.25	86.65	85.00
1: $10^3$	86.50	88.35	86.60	84.90
1: $10^4$	86.50	88.35	86.60	85.00
1: $10^5$	86.40	88.15	86.50	85.00
1: $10^6$	86.40	88.25	86.60	85.10
1: $10^7$	86.40	88.25	86.60	85.10
1: $10^8$	-	-	-	-

-, Not detected.

and *A. tubaeforme*, respectively. The melting curves and detection results are shown in Fig. 3. All detection results were identical to the sequencing results.



**Fig. 3.** Melting curve of  $T_m$ -shift based on ITS101 (A) and ITS296 (B) and detection results for 12 hookworm -positive samples.

**Table 4.** The accuracy of  $T_m$ -shift method

Sample	Species	ITS101 ( $T_m$ ) (°C)	ITS296 ( $T_m$ ) (°C)	GenBank
M6	<i>A. ceylanicum</i>	88.25	85.00	KF279134
M55	<i>A. ceylanicum</i>	88.25	85.00	KF279135
M58	<i>A. ceylanicum</i>	88.25	85.00	KF279136
M60	<i>A. ceylanicum</i>	88.35	85.00	KF279137
M76	<i>A. ceylanicum</i>	88.25	85.10	KF279138
C1	<i>A. tubaeforme</i>	86.50	86.60	MG904956
C2	<i>A. tubaeforme</i>	86.50	86.60	MG904957
C3	<i>A. tubaeforme</i>	86.40	86.60	MG904958
C4	<i>A. tubaeforme</i>	86.50	86.60	MG904959
C5	<i>A. tubaeforme</i>	86.40	86.50	MG904960

Sample M6, M55, M58, M60, M76 were isolated and identified by Liu et al. [4]. The sequences of sample C1, C2, C3, C4, and C5 have been uploaded to GenBank.

## DISCUSSION

*Ancylostoma ceylanicum* and *A. tubaeforme* are the 2 most common parasitic nematodes in cats. The former is the only animal-derived hookworm that can cause severe infection in humans. In Malaysia, the infection rate of *A. ceylanicum* is as high as 23.4%, which has become the second largest hook-

worm in local people [28]. In Laos, one-third of human hookworm infections are caused by *A. ceylanicum* [29]. Also, there are reports of human infection with *A. ceylanicum* in Taiwan [30]. In recent years, many cases of cat infection with *A. tubaeforme* have been detected in Guangzhou, China [2]. With the economic development and life quality improvement, more and more people are raising cats. Hence, it is of great significance to establish a  $T_m$ -shift detection method for cat-derived *A. ceylanicum* and *A. tubaeforme* for the prevention and control of zoonotic hookworm disease.

Common molecular biological methods for hookworm species identification include polymerase chain reaction (PCR), multiplex PCR, restriction fragment length polymorphism (RFLP), and high-resolution melting curve (HRM) [18-20]. Although the first 3 methods are specific and sensitive, they require gel electrophoresis after PCR, which makes the samples can be easily contaminated. In addition, the long-term use of ethidium bromide may seriously endanger the health of the operator. As a new molecular detection method,  $T_m$ -shift method is simple and quick to operate compared with the above-mentioned methods. The  $T_m$ -shift method is completed at one time on a closed fluorescent PCR instrument

minimizing the contamination risk. As well as, it can examine 36 or 72 samples at the same time. Compared with HRM, the  $T_m$ -shift method does not require a relatively complicated HRM program. The used fluorescent dye in  $T_m$ -shift method is the ordinary SYBR Green I fluorescent dye, which is less expensive than the saturated dye required for HRM. Furthermore, the  $T_m$ -shift method can complete the amplification of the target fragment by using a 2-step method instead of 3 steps in HRM, which greatly saves the experiment time and improves the detection efficiency.

Even though the  $T_m$ -shift detection method has clear benefits over the other methods, it has some limitations during application. For example, the  $T_m$ -shift method has higher requirements for the design of 2 specific primers. The  $T_m$  value of the specific primer should be between 59°C and 62°C, and the primer length should be controlled in 15-22 bp. For a higher *Taq* polymerase amplification efficiency, the difference of  $T_m$  value between the reverse and specific primer (before adding sequences at the 5' end) should be ranged from 4 to 5°C. In our study during the establishment of  $T_m$ -shift detection methods for cat-derived *A. ceylanicum* and *A. tubaeforme*, the primers were optimized where the short chain "gcggcg" was replaced with "gattaccg". The results showed that this operation increased the  $T_m$  value of 2 melting curves to some extent. The difference between the peak of the melting curve for the 2 hookworms was 1.75°C at ITS101, while it was 1.60°C at ITS296. The established  $T_m$ -shift detection method was applied to examine 69 stray cat fecal samples in Shaoguan, Guangdong. The examination results of clinical fecal samples showed that 12 samples were positive for hookworm, from which 8 samples were *A. ceylanicum* and 4 samples were *A. tubaeforme*. The positive rate of  $T_m$ -shift method was consistent with the microscopic examination results, and this method could distinguish *A. ceylanicum* from *A. tubaeforme*. These results indicated that the designed  $T_m$ -shift method is rapid, accurate, efficient and sensitive.

It could be concluded that the developed  $T_m$ -shift method based on the 2 SNPs loci of ITS1 sequence can successfully identify *A. ceylanicum* and *A. tubaeforme* species. As well as, it can provide a suitable technical mean for clinical diagnosis and zoonotic risk judgment of cat-derived hookworms.

## ACKNOWLEDGMENT

This work was supported by a grant from National Natural

Science Foundation of China (grant no. 31672541) and the Science and Technology Planning Project of Guangdong Province, China (grant no. 2014A020214005).

## CONFLICTS OF INTEREST

The authors declare that they have no conflicts of interest.

## REFERENCES

- Hu W, Wu S, Yu XG, Tan LP, Song MR, Abdulahi AY, Wang Z, Jiang B, Li GQ. Levels of *Ancylostoma* infections and phylogenetic analysis of cox 1 gene of *A. ceylanicum* in stray cat faecal samples from Guangzhou, China. *J Helminthol* 2015; 90: 392-397.
- Shi XL, Fu YQ, Abdullahi AY, Wang MW, Yang F, Yu XG, Pan WD, Yan XX, Hang JX, Zhang P, Li GQ. The mitochondrial genome of *Ancylostoma tubaeforme* from cats in China. *J Helminthol* 2018; 92: 22-33.
- Chen J, Xu MJ, Zhou DH, Song HQ, Wang CR, Zhu XQ. Canine and feline parasitic zoonoses in China. *Parasite Vectors* 2012; 5: e152.
- Liu YJ, Zheng GC, Zhang P, Alsarakibi M, Zhang XH, Li YW, Liu T, Ren SN, Chen ZX, Liu YL, Li SJ, Li GQ. Molecular identification of hookworms in stray and shelter dogs from Guangzhou city, China using ITS sequences. *J Helminthol* 2015; 89: 196-202.
- Kaya D, Yoshikawa M, Nakatani T, Tomo-oka F, Fujimoto Y, Ishida K, Fujinaga Y, Aihara Y, Nagamatsu S, Matsuo E, Tokoro M, Ouji Y, Kikuchi E. *Ancylostoma ceylanicum* hookworm infection in Japanese traveler who presented chronic diarrhea after return from Lao People's Democratic Republic. *Parasitol Int* 2016; 65: 737-740.
- Ngui R, Ching LS, Kai TT, Roslan MA, Lim YA. Molecular identification of human hookworm infections in economically disadvantaged communities in Peninsular Malaysia. *Am J Trop Med Hyg* 2012; 86: 837-842.
- Conlan JV, Khamlome B, Vongxay K, Elliot A, Pallant L, Sripa B, Blacksell SD, Fenwick S, Thompson RC. Soil-Transmitted Helminthiasis in Laos: a community-wide cross-sectional study of humans and dogs in a mass drug administration environment. *Am J Trop Med Hyg* 2012; 86: 624-634.
- Traub RJ, Inpankaew T, Sutthikornchai C, Sukthana Y, Thompson RC. PCR-based coprodiagnostic tools reveal dogs as reservoirs of zoonotic ancylostomiasis caused by *Ancylostoma ceylanicum* in temple communities in Bangkok. *Vet Parasitol* 2008; 155: 67-73.
- Wilson-Hanson SL, Prescott CW. A survey for parasites in cats. *Aust Vet J* 1982; 59: 194.
- Meloni BP, Thompson RC, Hopkins RM, Reynoldson JA, Gracey M. The prevalence of *Giardia* and other intestinal parasites in children, dogs and cats from aboriginal communities in the Kimberley. *Med J Aust* 1993; 158: 157-159.

11. Labarthe N, Serrão ML, Ferreira AM, Almeida NK, Guerrero J. A survey of gastrointestinal helminths in cats of the metropolitan region of Rio de Janeiro, Brazil. *Vet Parasitol* 2004; 123: 133-139.
12. Riggio F, Mannella R, Ariti G, Perrucci S. Intestinal and lung parasites in owned dogs and cats from central Italy. *Vet Parasitol* 2013; 193: 78-84.
13. Abu-Madi MA, Behnke JM, Prabhaker KS, Al-Ibrahim R, Lewis JW. Intestinal helminths of feral cat populations from urban and suburban districts of Qatar. *Vet Parasitol* 2010; 168: 284-292.
14. Calvete C, Lucientes J, Castillo JA, Estrada R, Gracia MJ, Peribáñez MA, Ferrer M. Gastrointestinal helminth parasites in stray cats from the mid-Ebro Valley, Spain. *Vet Parasitol* 1998; 75: 235-240.
15. Anderson TC, Foster GW, Forrester DJ. Hookworms of feral cats in Florida. *Vet Parasitol* 2003; 115: 19-24.
16. Gates MC, Nolan TJ. Endoparasite prevalence and recurrence across different age groups of dogs and cats. *Vet Parasitol* 2009; 166: 153-158.
17. Traub RJ. *Ancylostoma ceylanicum*, a re-emerging but neglected parasitic zoonosis. *Int J Parasitol* 2013; 43: 1009-1015.
18. Hu W, Wu S, Yu X, Abdullahi AY, Song M, Tan L, Wang Z, Jiang B, Li G. A multiplex PCR for simultaneous detection of three zoonotic parasites *Ancylostoma ceylanicum*, *A. caninum*, and *Giardia lamblia* Assemblage A. *BioMed Res Int* 2015; 2015: e406168.
19. Liu Y, Zheng G, Alsarakibi M, Zhang X, Hu W, Lu P, Lin L, Tan L, Luo Q, Li G. Molecular identification of *Ancylostoma caninum* isolated from cats in Southern China based on complete ITS sequence. *BioMed Res Int* 2013; 2013: e868050.
20. Ngui R, Lim YA, Chua KH. Rapid detection and identification of human hookworm infections through high resolution melting (HRM) analysis. *PLoS One* 2012; 7: e41996.
21. Germer S, Higuchi R. Single-tube genotyping without oligonucleotide probes. *Genome Res* 1999; 9: 72-78.
22. Wang J, Chuang K, Ahluwalia M, Patel S, Umblas N, Mirel D, Higuchi R, Germer S. High-throughput SNP genotyping by single-tube PCR with *T<sub>m</sub>*-shift primers. *Biotechniques* 2005; 39: 885-893.
23. Huang Y, Nie S, Zhou S, Li K, Sun J, Zhao J, Fei B, Wang Z, Ye H, Hong Q, Duan S. PPARD rs2016520 polymorphism and circulating lipid levels connect with brain diseases in Han Chinese and suggest sex-dependent effects. *Biomed Pharmacother* 2015; 70: 7-11.
24. Derzelle S, Mendy C, Laroche S, Madani N. Use of high-resolution melting and melting temperature-shift assays for specific detection and identification of *Bacillus anthracis* based on single nucleotide discrimination. *J Microbiol Methods* 2011; 87: 195-201.
25. Ahn JJ, Kim Y, Hong JY, Kim GW, Kim SY, Hwang SY. Probe-based fluorescence melting curve analysis for differentiating *Larimichthys polyactis* and *Larimichthys crocea*. *Food Anal Methods* 2016; 9: 2036-2041.
26. Pan W, Fu Y, Abdullahi AY, Wang M, Shi X, Yang F, Yu X, Yan X, Zhang P, Hang J, Li GQ. Development of *T<sub>m</sub>*-shift genotyping method for detection of cat-derived *Giardia lamblia*. *Parasitol Res* 2017; 116: 1151-1157.
27. Fu Y, Wang M, Yan X, Abdullahi AY, Hang J, Zhang P, Huang Y, Liu Y, Sun Y, Ran R, Li GQ. *T<sub>m</sub>*-Shift detection of dog-derived *Ancylostoma ceylanicum* and *A. caninum*. *BioMed Res Int* 2018; 2018: e7617094.
28. Ngui R, Lim YA, Traub R, Mahmud R, Mistam MS. Epidemiological and genetic data supporting the transmission of *Ancylostoma ceylanicum* among human and domestic animals. *PLoS Negl Trop Dis* 2012; 6: e1522.
29. Conlan JV, Sripa B, Attwood S, Newton PN. A review of parasitic zoonoses in a changing Southeast Asia. *Vet Parasitol* 2011; 182: 22-40.
30. Hsu YC, Lin JT. Images in clinical medicine. Intestinal infestation with *Ancylostoma ceylanicum*. *New Engl J Med* 2012; 366: e20.