

Molecular Prevalence of *Cryptosporidium* spp. in Breeding Kennel Dogs

Naoyuki Itoh*, Hazuki Tanaka, Yuko Iijima, Satoshi Kameshima, Yuya Kimura

Laboratory of Small Animal Internal Medicine, School of Veterinary Medicine, Kitasato University, Towada, Aomori, Japan

Abstract: *Cryptosporidium* is a common intestinal protozoan that can lead to diarrhea in humans and dogs. The predominant species of infection are *C. hominis* and *C. parvum* in humans, and *C. canis* in dogs. However, *C. canis* can infect immunocompromised humans. Considering the close contact with humans, dogs have the potential to be reservoirs for human cryptosporidiosis. Breeding kennels are the major supply source of puppies for pet shops. The present study is to determine the molecular prevalence and characteristics of *Cryptosporidium* spp. found in breeding kennel dogs. A total of 314 fecal samples were collected from young and adult dogs kept in 5 breeding kennels. A polymerase chain reaction targeting the small subunit rRNA gene was employed for the detection of *Cryptosporidium* spp. To determine the species, the DNA sequences were compared to GenBank data. Overall, 21.0% of the fecal samples were positive for *Cryptosporidium* spp. infection. *Cryptosporidium* spp. was detected in all 5 facilities. A sequencing analysis demonstrated that all isolates shared 99-100% similarity with *C. canis*. The results suggest that *Cryptosporidium* spp. infection is present at a high-level in breeding kennel dogs. However, because dominant species in this survey was *C. canis*, the importance of breeding kennel dogs as reservoirs for *Cryptosporidium* spp. transmission to humans is likely to be low in Japan.

Key words: *Cryptosporidium* spp., *Cryptosporidium canis*, dog, breeding kennel

Cryptosporidium is a common intestinal protozoan in many mammals, including humans and dogs, with a recognized potential to cause pathogen digestive tract obstruction in hosts [1-3]. The genus *Cryptosporidium* comprises more than 20 different species and at least 30 genotypes [3-5]. It was previously thought that each *Cryptosporidium* spp. had a narrow host range and was adapted to a specific animal or group of closely related animals [5]. For example, *C. hominis* and *C. canis* are apparently adapted to humans and dogs, respectively [3,4]. However, recently, it was demonstrated that some *Cryptosporidium* spp., such as *C. parvum* and *C. muris*, possess a broad range of hosts beyond the known animal species; therefore, they are zoonotic species [4,5]. Although there is no doubt that *C. hominis* and *C. parvum* predominantly infect humans and that *C. canis* predominantly infects dogs, *C. canis* and *C. parvum* have been isolated from both hosts [4,5]. Thus, *C. canis* can infect humans and dogs. Considering the close contact

dogs have with humans, dogs have the potential to be reservoirs for human cryptosporidiosis. Immunocompromised individuals, such as those infected with HIV, are at especially high risk for *C. canis* infection [2,5]. In Japan, the population of suspected immunocompromised individuals is growing because of an increasing elderly population and the increasing incidence of cancer.

The authors have previously determined the molecular prevalence and characteristics of the *Cryptosporidium* spp. in private household dogs and pet shop puppies and have demonstrated that pet shop puppies have a high-level prevalence of *C. canis* infection without exhibiting clinical signs [6]. Therefore, the present hypothesis that *Cryptosporidium* spp. will be detected at high-level prevalence in puppies/young dogs living in breeding kennel is logical, because the breeding kennels are the major source of puppies supplied to pet shops. Nevertheless, there is insufficient available data regarding the prevalence of *Cryptosporidium* spp. infection in breeding kennel dogs (young and adult); no such data exist in Japan [7-9].

The aim of the present study is to determine the molecular prevalence and characteristics of *Cryptosporidium* spp. in breeding kennel dogs, including an analysis of some factors such as dog age, so as to evaluate the risk of zoonotic transmission

•Received 26 December 2018, revised 27 February 2019, accepted 27 February 2019.

*Corresponding author (naoitoh@vmas.kitasato-u.ac.jp)

© 2019, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

from breeding kennel dogs to humans.

Between August 2014 and July 2017, a total of 314 fresh voided fecal samples were randomly collected on a single occasion from dogs (aged 2 month-11 years, 85 males and 229 females) kept in 5 breeding kennels (A-E), located in the prefectures of Miyagi (A; n=25), Niigata (B; n=48), Gunma (C; n=87), Shizuoka (D; n=31), and Aichi (E; n=123), in east Japan. All breeding kennel managers granted permission to include their dogs in the examination and donated fecal samples. The samples were collected immediately after natural defecation and were stored at 4°C prior to DNA extraction, which occurred within 3 days of collection.

The oocysts of *Cryptosporidium* spp. were separated using a sucrose gradient concentration method with a specific gravity of 1.26, and DNA extraction from oocysts was performed using a QIAamp DNA Mini Kit (QIAGEN GmbH, Hilden, Germany) according to the manufacturer's instructions. The DNA samples were stored at -20°C prior to analysis.

Two-step nested polymerase chain reaction (PCR) targeting the small subunit (SSU) rRNA gene was employed for the detection of *Cryptosporidium* spp. For the first step reaction, the forward primer (5'-TTCTAGAGCTAATACATGCG-3') and the reverse primer (5'-CCCATTTTCCTTCGAAACAGGA-3') were used to amplify a DNA fragment of approximately 1,325 bp. For the second step reaction, the forward primer (5'-GGAAGG GTTGTATTATTAGATAAAG-3') and the reverse primer (5'-AAGGAGTAAGGAACAACCTCCA-3') were used to amplify a fragment of approximately 826 bp [10].

For the first step reaction, the PCR mixture comprised 1×buffer containing 1.5 mM MgCl₂, 200 μM of each dNTP, 0.5 μM of each primer, 1.25 units of GoTaq DNA polymerase (Promega Corporation, Madison, Wisconsin, USA), and 2.0 μl of template DNA in a total reaction volume of 25 μl. For the second step reaction, the PCR mixture was the same as for the first step reaction, except the amplicons from the primary PCR were used as templates. The first-step PCR was performed as follows: after an initial denaturation step of 3 min at 95°C, 35 cycles, which each consist of 45 sec at 95°C, 45 sec at 59°C and 1 min at 72°C, were performed, followed by a final extension of 5 min at 72°C. The second-step PCR condition was performed as follows: after an initial denaturation step of 3 min at 95°C, 35 cycles, which each consist of 30 sec at 95°C, 1 min at 58°C and 1 min at 72°C, were performed, followed by a final extension of 5 min at 72°C.

All second-step PCR products were identified by electropho-

resis on 1.5% agarose gels. The specific DNA fragments (approximately 826 bp) were confirmed by alternative ethidium bromide staining and visualization under UV light using a transilluminator.

Second-step PCR amplicons of the predicted size were purified using a QIAquick Gel Extraction kit (QIAGEN GmbH, Hilden, Germany) and were sequenced with the second-step primer set. Sequences were analyzed by a commercial laboratory (FASMAC Co., Ltd., Kanagawa, Japan).

Sequence alignment and compilation were performed using the MEGA 6.06 program (www.megasoftware.net). To determine the species of *Cryptosporidium*, the DNA sequences were compared to GenBank sequences of *Cryptosporidium* spp. by BLAST searches (<http://www.ncbi.nlm.nih.gov/>). The similarity between the isolated and reference sequences was determined based on the degree of sequence identity.

The data were stratified according to age group (<1-year-old vs ≥1-year-old), fecal condition (normal vs soft/diarrhea) and the breeding kennel. Data were analyzed statistically using Fisher's exact probability test, with values of $P < 0.05$ considered significant.

Overall, of the 314 breeding kennel dogs, 66 animals (21.0%) were positive for *Cryptosporidium* spp. infection. According to age group, the prevalence of *Cryptosporidium* spp. in <1-year-old dogs (45.0%) was significantly ($P < 0.001$) higher than that in ≥1-year-old dogs (17.5%) (Table 1). No significant difference was observed in fecal condition (normal: 20.7% vs soft/diarrhea: 23.3%). *Cryptosporidium* spp. infection was determined in all breeding kennel facilities, ranging from

Table 1. Prevalence of *Cryptosporidium* species infection in breeding kennel dogs

	Examined No.	Positive No.	Prevalence (%)	P-value
Overall	314	66	21.00	-
Age				
<1-year-old	40	18	45.00	< 0.001
≥1-year-old	274	48	17.50	-
Fecal condition				
Normal	271	56	20.70	-
Soft/Diarrhea	43	10	23.30	NS
Breeding kennel				
A	25	2	8.00	-
B	48	10	20.80	NS
C	87	17	19.50	NS
D	31	5	16.10	NS
E	123	32	26.00	NS

NS, not significant.

8.0% to 26.0% prevalence, with no significant differences observed between kennels.

A sequencing analysis of the SSU rRNA gene fragments demonstrated that all 66 PCR-positive amplicons shared 99-100% similarity with the sequences of *C. canis* (accession numbers: KR999987, KR999984, JN543380, MF589918, KF516542 and EU754826) retrieved from the GenBank database.

The present study is the first report describing the molecular prevalence of *Cryptosporidium* spp. infection in breeding kennel dogs in Japan. The results of the study support the authors' hypothesis because *Cryptosporidium* spp. infection was detected at markedly high prevalence (45.0%) in young dogs in breeding kennels. In addition, the prevalence of infection was also high in adult dogs (17.5%). In contrast, previous recent studies have reported low prevalences of *Cryptosporidium* spp. infection in breeding kennel dogs, such as 5.0% in Italy [7], 0% in Belgium [8] and 7.0% in China [9]. The cause of low infection prevalence in those previous cases is not known. A significantly high-level detection of intestinal protozoans, such as *Giardia* and *Cystoisospora*, has been reported in breeding kennel dogs [8,11]. Additionally, a high prevalence of *Cryptosporidium* spp. infection has been detected in pet shop puppies [6], most of which were likely derived from breeding kennels. However, the authors cannot simply compare the results of the present study to those of previous studies due to differences in dog's ages, study regions, and examination methods. Therefore, the most important highlight here is the suggestion of high-level *Cryptosporidium* spp. infection in not only young dogs, but also adult animals kept in breeding kennels.

A high-level of *Cryptosporidium* spp. infection in younger dogs, including puppies, has previously been documented [6,12,13]. Although the role of immune responses against *Cryptosporidium* spp. infections is unknown in dogs, the immature immune system of young dogs is suspected to be a major factor for high-level infection. Because it is commonly accepted that both innate and adaptive immune responses are important for controlling *Cryptosporidium* spp. infections [2,14,15], an immature immune system is indicated in young dogs [16,17]. Additionally, the stressful situations presented in breeding kennels, such as frequent contact with other animals in a limited space, is also a significant factor for high prevalence due to the induction of immunosuppressive status [18]. Simultaneously, the stressful situations in breeding kennels are likely to be factors in the high prevalence of infection in adult dogs. In the present study, the prevalence of infection in

≥ 1-year-old male dogs (15.4%; 12/78 dogs) was roughly the same as that in females in the same age group (18.4%; 36/196 dogs). Lastly, sanitary management and environmental conditions were not evaluated here, but environmental contamination via *Cryptosporidium* oocysts is considered the most important factor for high prevalence in breeding kennel dogs. It has been demonstrated that the *Cryptosporidium* oocyst is resistant against common disinfectants and can survive in environments for duration of at least several months [2,19]; thus, it is very difficult to eliminate *Cryptosporidium* oocysts in breeding kennel facilities. In all breeding kennels, the above factors contribute to maintain a high-level of *Cryptosporidium* spp. infection, and frequent reinfection is suspected. As with results from related previous studies, there was no correlation between the detection of *Cryptosporidium* spp. in dogs and the fecal conditions [6,20]. The coinfection with other intestinal protozoa such as *Giardia* spp. is suggested as a factor of inducing clinical signs [1].

The sequencing data demonstrates the dominance of *C. canis* in breeding kennel dogs in Japan. It is easy to understand this result, because *C. canis* is a dog-adapted species [3,4]. Therefore, the importance of breeding kennel dogs as reservoirs for *Cryptosporidium* spp. transmission to humans is likely to be low in Japan, since the predominant species for human cryptosporidiosis are *C. hominis* and *C. parvum*. However, we should remember that immunocompromised humans have a potential risk for infection of *C. canis* [2,5].

CONFLICT OF INTEREST

On behalf of all authors, the corresponding author states that there is no conflict of interest.

REFERENCES

1. Scorza S, Tangtrongsup S. Update on the diagnosis and management of *Cryptosporidium* spp infections in dogs and cats. *Top Companion Anim Med* 2010; 25: 163-169.
2. Bouzid M, Hunter PR, Chalmers RM, Tyler KM. *Cryptosporidium* pathogenicity and virulence. *Clin Microbiol Rev* 2013; 26: 115-134.
3. Ryan U, Zahedi A, Paparini A. *Cryptosporidium* in humans and animals--a one health approach to prophylaxis. *Parasite Immunol* 2016; 38: 535-547.
4. Feng Y, Ryan UM, Xiao L. Genetic diversity and population structure of *Cryptosporidium*. *Trends Parasitol* 2018; 34: 997-1011.
5. Khan A, Shaik JS, Grigg ME. Genomics and molecular epidemi-

- ology of *Cryptosporidium* species. *Acta Trop* 2018; 184: 1-14.
6. Itoh N, Oohashi Y, Ichikawa-Seki M, Itagaki T, Ito Y, Saeki H, Kanai K, Chikazawa S, Hori Y, Hoshi F, Higuchi S. Molecular detection and characterization of *Cryptosporidium* species in household dogs, pet shop puppies, and dogs kept in a school of veterinary nursing in Japan. *Vet Parasitol* 2014; 200: 284-288.
 7. Giangaspero A, Iorio R, Paoletti B, Traversa D, Capelli G. Molecular evidence for *Cryptosporidium* infection in dogs in Central Italy. *Parasitol Res* 2006; 99: 297-299.
 8. Claerebout E, Casaert S, Dalemans AC, De Wilde N, Levecke B, Vercruyse J, Geurden T. *Giardia* and other intestinal parasites in different dog populations in Northern Belgium. *Vet Parasitol* 2009; 161: 41-46.
 9. Jian F, Qi M, He X, Wang R, Zhang S, Dong H, Zhang L. Occurrence and molecular characterization of *Cryptosporidium* in dogs Henna Province, China. *BMC Vet Res* 2014; 10: 26.
 10. Xiao L, Morgan UM, Limor J, Escalante A, Arrowood M, Shulaw W, Thompson RCA, Fayer R, Lal AA. Genetic diversity within *Cryptosporidium parvum* and related *Cryptosporidium* species. *Appl Environ Microbiol* 1999; 65: 3386-3391.
 11. Itoh N, Kanai K, Kimura Y, Chikazawa S, Hori Y, Hoshi F. Prevalence of intestinal parasites in breeding kennel dogs in Japan. *Parasitol Res* 2015; 114: 1221-1224.
 12. Hamnes IS, Gjerde BK, Robertson LJ. A longitudinal study on the occurrence of *Cryptosporidium* and *Giardia* in dogs during their first year life. *Acta Vet Scand* 2007; 49: 22.
 13. Uehlinger FD, Greenwood SJ, McClure JT, Conboy G, O'Handley R, Barkema HW. Zoonotic potential of *Giardia duodenalis* and *Cryptosporidium* spp. and prevalence of intestinal parasites in young dogs from different populations on Prince Edward Island, Canada. *Vet Parasitol* 2013; 196: 509-514.
 14. Petry F, Jakobi V, Tessema TS. Host immune response to *Cryptosporidium parvum* infection. *Exp Parasitol* 2010; 126: 304-309.
 15. McDonald V, Korbel DS, Barakat FM, Choudhry N, Petry F. Innate immune responses against *Cryptosporidium parvum* infection. *Parasite Immunol* 2013; 35: 55-64.
 16. Faldyna M, Levá L, Knöťigová P, Toman M. Lymphocyte subsets in peripheral blood of dogs--a flow cytometric study. *Vet Immunol Immunopathol* 2001; 82: 23-37.
 17. Day MJ. Immune system development in the dog and cat. *J Comp Pathol* 2007; 137 (suppl): 10-15.
 18. Powell ND, Allen RG, Hufnagle AR, Sheridan JF, Bailey MT. Stressor-induced alterations of adaptive immunity to vaccination and viral pathogens. *Immunol Allergy Clin N Am* 2011; 31: 69-79.
 19. Sunnotel O, Lowery CJ, Moore JE, Dooley JSG, Xiao L, Millar BC, Rooney PJ, Snelling WJ. *Cryptosporidium*. *Lett Appl Microbiol* 2006; 43: 7-16.
 20. Wang A, Ruch-Gallie R, Scorza V, Lin P, Lappin MR. Prevalence of *Giardia* and *Cryptosporidium* species in dog park attending dogs compared to non-dog park attending dogs in one region of Colorado. *Vet Parasitol* 2012; 184: 335-340.