

Production of Inflammatory Cytokines and Nitric Oxide by Human Mast Cells Incubated with *Toxoplasma gondii* Lysate

Eun-Ah Park¹, Ik-Hwan Han², Jung-Hyun Kim², Soon-Jung Park¹, Jae-Sook Ryu², Myoung-Hee Ahn^{2,*}

¹Department of Environmental Medical Biology and Institute of Tropical Medicine, Brain Korea 21 PLUS Project for Medical Science, Yonsei University College of Medicine, Seoul 03722, Korea; ²Department of Environmental Biology and Medical Parasitology, Hanyang University College of Medicine, Seoul 04763, Korea

Abstract: The roles of mast cells in allergic diseases and helminth infections are well known. However, the roles of mast cells in *T. gondii* infection is poorly understood. This study was focused on the production of pro-inflammatory cytokines (TNF- α , IL-4), chemokines (CXCL8, MCP-1) and nitric oxide (NO) by mast cells in response to soluble lysate of *T. gondii* tachyzoites. Production of CXCL8 (IL-8), MCP-1, TNF- α and IL-4 were measured by RT-PCR and ELISA. Western blot were used for detection of CXCR-1 and CXCR2. Our results showed that *T. gondii* lysates triggered mast cells to release CXCL8, MCP-1, TNF- α , IL-4 and to produce NO. This suggests that mast cells play an important role in inflammatory responses to *T. gondii*.

Key words: *Toxoplasma gondii*, mast cell, cytokine, nitric oxide

Toxoplasma gondii is an obligate intracellular protozoan parasite with a global distribution in humans and other mammals. Although this parasite most often causes subclinical infection, primary infection during pregnancy can induce fetal damage and abortion in both humans and animals. *T. gondii* is a well-known inducer of type 1 cytokines, and while these cytokines are required to survive infection, their over-production can be harmful and even lethal [1,2].

Mast cells are abundant in the skin, mucosa of the digestive and respiratory tracts and conjunctiva. These cells provoke IgE-mediated allergic responses and are involved in the control of helminth infections of the gastrointestinal tract. They play a key role in innate and adaptive immunity [3,4] and have been shown to induce acute inflammatory reactions in mice in response to intraperitoneal injection of *T. gondii* [5]. Mast cells also protect the host when *T. gondii* is introduced by the oral route [3]. Mediators produced by activated mast cells may reduce acute inflammation and stimulate parasite clearance in *T.*

gondii infections [6]. On the other hand, *T. gondii* itself inhibits mast cell degranulation by suppressing the mobilization of intracellular Ca²⁺ by phospholipase C [7]. So, the overall effect of mast cells on *T. gondii* infections is uncertain. In addition, it is not known whether *T. gondii* lysates can induce inflammatory responses in the mast cell leukaemia cell line, HMC-1. The aim of this study was to evaluate whether *T. gondii* lysates induce the activation of HMC-1 cells to release cytokines and nitric oxide.

T. gondii (RH strain) tachyzoites were maintained by intra-peritoneal infection of ICR mice (Osan, Korea). Tachyzoites were harvested from the mice, washed with PBS and centrifuged at low and high speeds (500 g and 3,000 g) for 5 min to remove peritoneal cells. To prepare *T. gondii* lysate, the tachyzoites were frozen and thawed 3 times and ultrasonicated at 100% amplitude, 0.75 cycles, 7-10 times, until the tachyzoites were completely disrupted. The sonicate was centrifuged at 10,000 g for 1 hr, and the supernatant was filtered through a 0.2 μ m membrane (Millipore, Bedford, Massachusetts, USA). Protein concentrations were determined with the Bradford assay (Bio-Rad, Hercules, California, USA).

Human leukemic mast cells (HMC-1) were grown in IMDM supplemented with 10% FBS at 37°C in a 5%-CO₂ incubator. To examine cytokine and iNOS mRNA expression, HMC-1

•Received 10 December 2018, revised 14 March 2019, accepted 19 March 2019.

*Corresponding author (mhahn@hanyang.ac.kr)

© 2019, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

cells (1.5×10^6 /well) were pretreated with IFN- γ (5 μ g/ml, ProSpec, East Brunswick New Jersey, USA) for 1 hr and then stimulated with an aliquot of *T. gondii* lysate (50 μ g/ml) or LPS (10 μ g/ml, Sigma, St Louis, Missouri, USA). As a positive control, the HMC-1 cells were incubated with PMA (100 nm/ml) and A23187 (10 μ m/ml). Total RNA was extracted from the cells using Trizol reagent (Invitrogen, Carlsbad, California, USA) and 200 μ l chloroform (Sigma Aldrich, St. Louis, Missouri, USA). cDNAs were synthesized and reverse-transcribed using RT Premix and PCR Premix (iNtRON Biotechnology, Seongnam, Korea). Primer sequences were as follows; β -actin (5'-CCA GAG CAA GAG AGG TAT CC-3' and 5'-CTG TGG TGG TGA AGC TGT AG-3'), human IL-8 (5'-GCC AAG AGA ATA TCC GAA CT-3' and 5'-AAA GTG CAA CCA CAT GTC CT-3'), MCP-1 (5'-TCT GTG CCT GCT GCT CAT AG-3' and 5'-GCT GCA GAT TCT TGG GTT GT-3'), IL-4 (5'-CCC TTT CGG CAA AAT CCT CC-3' and 5'-AGG AAT ACA GTG CAG CTT ACC A-3'), TNF- α (5'-ACT CTT CTG CCT GCT GCA CTT TGG-3' and 5'-GTT GAC CTT TGT CTG GTA GGA GAC GG-3'), iNOS (5'-AGC ATG AGC CCC TTC ATC AAT-3' and 5'-CTG TTT CAA CGA CCT CCG GG-3'), and GAPDH (5'-GTC AGT GGT GGA CCT GAC CT-3' and 5'-AGG GGT CTA CAT GGC AAC TG-3'). PCR conditions were initial DNA denaturation at 94°C for 5 min followed by 35 rounds of denaturation at 98°C for 15 sec, and annealing for 30s at 58°C for β -actin, GAPDH & iNOS, 56°C for IL-8 & IL-4, 55°C for MCP-1, and 62°C for TNF- α , and extension at 72°C for 35 sec.

To measure IL-8, MCP-1, TNF- α and IL-4 proteins, HMC-1 cells were pretreated with IFN- γ followed by exposure to *T. gondii* lysate or LPS for 18 hr or 24 hr. Supernatants were collected and stored at -20°C and cytokines were measured by ELISA (BD Biosciences, San Diego, California, USA).

Western blot analysis with monoclonal antibodies was used to detect expression of IL-8 receptor (CXCR1 and CXCR2) on HMC-1 cells which were pretreated with IFN- γ for 1 hr followed by *T. gondii* lysate or LPS. After incubation, the cells were harvested and lysed in PRO-PREP protein extraction solution (iNtRON Biotechnology, Seongnam, Korea). Equal amounts of protein were denatured and separated by SDS-PAGE on 10% polyacrylamide gels and transferred to Immune-Blot1 PVDF membranes (Bio-Rad, Quarry Bay, HongKong). The membranes were probed with anti-CXCR1 and anti-CXCR2 antibodies (1:1,000; Abcam, Cambridge, UK), or β -actin polyclonal antibody (1:3,000; Abcam, Cambridge, Massachusetts, USA) overnight at 4°C followed by goat anti-rabbit IgG poly-

clonal antibody (1:10,000; ADISAB-300-J, Enzo Life Sciences, New York, USA) for 1 hr at room temperature. The blots were visualized using Chemiluminescent Sensitive Plus HRP Microwell and/or Membrane Substrate (SurModics, Minneapolis, Minnesota, USA), and signals were measured with a Chemo-Doc1 (Bio-Rad).

For measuring nitric oxide, HMC-1 cells was pretreated with IFN- γ followed by *T. gondii* lysate or LPS for 3 hr, 6 hr, 18 hr or 24 hr and supernatants were collected. NO was determined as nitrite using Griess reagent [8].

Statistical analyses were performed with SPSS, version 21 (IBM, Chicago, Illinois, USA). The Mann-Whitney U test was used to compare results, and *P*-values < 0.05 were considered statistically significant. Data are expressed as means \pm SDs of three independent experiments.

In this experiment, up-regulation of IL-8 mRNA and increased secretion of IL-8 were observed in the *T. gondii*-treated HMC-1 cells 6 hr and 18 hr post inoculation (PI), respectively. However the IL-8 levels of HMC-1 cells stimulated with LPS as positive control were significantly higher than those of the of *T. gondii*-stimulated cells. Up-regulations of MCP-1 mRNA and protein were also observed in the HMC-1 cells exposed to *T. gondii* lysate or LPS. Similarly, enhanced TNF- α mRNA expression and increased TNF- α secretion were noted in supernatants of HMC-1 cells treated with *T. gondii* lysate. IL-4 mRNA was also increased by incubation with *T. gondii* lysate for 24 hr, and IL-4 protein increased in the supernatants of treated HMC-1 cells (Fig. 1A, B). To investigate the expression of chemokine receptors for IL-8, CXCR1 and CXCR2 production was analyzed by western blotting after treatment of HMC-1 cells with *T. gondii* lysate or LPS for 1 hr. Both CXCR1 and CXCR2 increased in response to *T. gondii* lysate (Fig. 1C).

The results in Fig. 2A show that iNOS mRNA increased in HMC-1 cells treated with *T. gondii* lysate for 3 hr, and then gradually disappeared. In LPS-treated cells, iNOS mRNA continued to increase between 3 hr and 24 hr of incubation although at a decreasing rate. Maximal NO production in HMC-1 cells treated with *T. gondii* lysate or LPS was observed after 3 hr of stimulation (Fig. 2B).

In *T. gondii* infections, immune cells including neutrophils, monocytes and macrophages are activated, and also these cells are targets of *T. gondii* [9,10]. *T. gondii* down-regulates type I immune responses, blocking production of pro-inflammatory cytokine including IFN- γ in macrophages and dendritic cells [11]. Mast cells are abundant in the intestinal mucosa and play




Fig. 1. Increased production of IL-8, MCP-1, TNF- α and IL-4 mRNAs and proteins by HMC-1 cells incubated with *T. gondii* lysate. HMC-1 cells were pretreated with IFN- γ for 1 hr and then stimulated with *T. gondii* lysate or LPS. mRNA expression of each cytokine (A) and cytokine level (B) were increased. CXCR1 and CXCR2 was measured by western blotting with monoclonal antibodies to each receptor after 1 hr of treatment with *T. gondii* lysate. PMA +A23187 was used as positive control. M, culture supernatant of HMC-1 alone; M*, culture supernatant of HMC-1 treated with IFN- γ ; TgL, *T. gondii* lysate. * $P < 0.05$ vs HMC-1 treated with IFN- γ .




Fig. 2. iNOS mRNA and nitric oxide (NO) productions were up-regulated by HMC-1 cells stimulated with *T. gondii* lysate. HMC-1 cells were pretreated with IFN- γ for 1 hr and then stimulated with *T. gondii* lysate or LPS. iNOS mRNA (A) and NO production (B) were determined after 3 hr, 6 hr, 18 hr, and 24 hr of incubation. NO production was measured with Griess reagent. PMA +A23187 was used as positive control. M, culture supernatant of HMC-1 cells alone; M*, culture supernatant of HMC-1 cells treated with IFN- γ ; TgL, *T. gondii* lysate. * $P < 0.05$ vs HMC-1 cells treated with IFN- γ .

an important role in protective immune responses to oral infection by *T. gondii* [3]. Exposure to *T. gondii* causes mast cell to activate, degranulate and release mediators, proteases and cytokines [12]. Increases in mast cells numbers may be involved in the balance between inflammation and wound healing [13].

In the current study we found that IL-8 and MCP-1 secretion was up-regulated after 18 hr exposure to *T. gondii* lysate. This result is in agreement with our previous report that infection with *T. gondii* tachyzoites leads to increased secretion of IL-8 and MCP-1 by HeLa cells and fibroblasts [14]. Up-regulation of TNF- α and IL-4 mRNAs was observed in HMC-1 cells treated with *T. gondii* lysate for 18 hr and 24 hr, respectively. Pro-

duction of both cytokines increased in *T. gondii* lysate- or LPS-treated HMC-1 cells indicating that *T. gondii* lysate induces inflammatory effect in the cells. *T. gondii* lysate alone did not increase cytokine production in HMC-1 cells. However, pretreatment of IFN- γ resulted in increased cytokine production in the cells. Therefore, IFN- γ was thought to be acted as a trigger for HMC-1 cells in cytokine production by *T. gondii* lysate.

The functions of chemokines induced by intracellular pathogens include leukocyte recruitment, cell mediated immunity and anti-protozoal activity. IL-8 induces a rapid and transient increase of Ca^{2+} in mast cells. HMC-1 cells express the CXCL8 chemokine receptors, CXCR1 and CXCR2 [15-18]. Increased levels of these receptors were observed on HMC-1

cells stimulated with *T. gondii* lysate for 1 hr (Fig. 1C).

The release of mediators by mast cell activation plays an important role in modulating inflammation during *T. gondii* infection [6,16]. In protozoal infections these mast cells are characterized by degranulation and the production of pro-inflammatory mediators, such as TNF- α [13]. The mast cell activator, compound 48/80, up-regulated Th1 cytokine mRNA (IFN- γ , IL-12p40, TNF- α), and the mast cell stabilizer, disodium cromoglycate, increased Th2 cytokine mRNA (IL-4, IL-10) [12].

Mast cells are effector cells involved in clearance of parasites including *Trichomonas vaginalis* (*T. vaginalis*) [6]. We observed that rat peritoneal mast cells (RPMCs) were involved in the inflammatory response to *T. vaginalis* infection. Increased release of histamine and TNF- α were observed in RPMC-stimulated *T. vaginalis* ESP or live trichomonads [19]. Activation of mast cells has been reported to trigger production of cytokines and inflammatory mediators by protozoan parasites such as *Plasmodium* sp., *Leishmania* sp., and *T. gondii* [12].

Several protozoan parasites have evolved strategies to evade NO-mediated anti-microbicidal activity. Inducible nitric oxide synthase (iNOS)-derived NO can inhibit leucocyte adhesion in the micro-vasculature during ongoing inflammation. *T. gondii* infection partially inhibits NO production by murine macrophages and thus promotes parasite survival in phagocytic cells [12,20,21]. *Leishmania* sp. and *Trypanosoma cruzi* are known to inhibit iNOS expression [22,23]. We investigated HMC-1 cells were treated with *T. gondii* lysate to determine whether *T. gondii* infection affects NO production. Increased iNOS mRNA and NO production were observed at 3 hr PI and it then declined by 6 hr.

Mast cells induce recruitment of other immune cells to initiate inflammation. In the present study, we observed that *T. gondii* lysate stimulated the production of chemokines (CXCL8, MCP-1), cytokines (TNF- α , IL-4) and NO production in HMC-1 cells. Therefore, activated mast cells by *T. gondii* lysate may be involved in innate and adaptive immune responses against *T. gondii* infection.

CONFLICT OF INTEREST

The authors declare no conflict of interest related to this study.

REFERENCES

- Henderson WR Jr, Chi EY. The importance of leukotrienes in mast cells-mediated *Toxoplasma gondii* cytotoxicity. *J Infect Dis* 1998; 177: 1437-1443.
- Długońska H. *Toxoplasma gondii* and mast cells. *Ann Parasitol* 2014; 60: 235-238.
- Cruz A, Mendes ÉA, de Andrade MV, do Nascimento VC, Cartelle CT, Arantes RM, Melo JR, Gazzinelli RT, Ropert C. Mast cells are crucial in the resistance against *Toxoplasma gondii* oral infection. *Eur J Immunol* 2014; 44: 2949-2954.
- Rodríguez NE, Wilson ME. Eosinophils and mast cells in leishmaniasis. *Immunol Res* 2014; 59: 129-141.
- S Ferreira GL, Mineo JR, Oliveira JG, V Ferro EA, Souza MA, D Santos AA. *Toxoplasma gondii* and mast cell interactions in vivo and in vitro: experimental infection approaches in *Calomys callosus* (Rodentia, Cricetidae). *Microbes Infect* 2004; 6: 172-181.
- Huang B, Huang S, Chen Y, Zheng H, Shen J, Lun ZR, Wang Y, Kasper LH, Lu F. Mast cells modulate acute Toxoplasmosis in murine models. *PLoS One* 2013; 8: 10; e77327.
- Smith NL, Abi Abdallah DS, Butcher BA, Denkers EY, Baird B, Holowka D. *Toxoplasma gondii* inhibits mast cell degranulation by suppressing phospholipase C γ -mediated Ca²⁺ mobilization. *Front Microbiol* 2013; 4: 179.
- Park GC, Ryu JS, Min DY. The role of nitric oxide as an effector of macrophage-mediated cytotoxicity against *Trichomonas vaginalis*. *Korean J Parasitol* 1997; 35: 189-195.
- Bierly AL, Shufesky WJ, Sukhumavasi W, Morelli AE, Denkers EY. Dendritic cells expressing plasmacytoid marker PDCA-1 are Trojan horse during *Toxoplasma gondii* infection. *J Immunol* 2008; 181: 8485-8491.
- Lambert H, Barragen A. Modelling parasite dissemination: host cell subversion and immune evasion by *Toxoplasma gondii*. *Cell Microbiol* 2010; 12: 292-300.
- Leng J, Butcher BA, Denkers EY. Dysregulation of macrophage signal transduction by *Toxoplasma gondii*: past progress and recent advances. *Parasite Immunol* 2009; 31: 717-728.
- Lu F, Huang S. The roles of mast cells in parasitic protozoan infections. *Front Immunol* 2017; 8: 363.
- Vukman KV, Lalor R, Aldridge A, O'Neill SM. Mast cells: new therapeutic target in helminth immune modulation. *Parasite Immunol* 2016; 38: 45-52.
- Lee EJ, Heo YM, Choi JH, Song HO, Ryu JS, Ahn MH. Suppressed production of pro-inflammatory cytokines by LPS-activated macrophages after treatment with *Toxoplasma gondii* lysate. *Korean J Parasitol* 2008; 46: 145-151.
- Inamura H, Kurosawa M, Okano A, Kayaba H, Majima M. Expression of the interleukin-8 receptors CXCR1 and CXCR2 on cord-blood-derived cultured human mast cells. *Int Arch Allergy Immunol* 2002; 128: 142-150.
- Gaddi PJ, Yap GS. Cytokine regulation of immunopathology in toxoplasmosis. *Immunol Cell Biol* 2007; 85: 155-159.

17. Denney CF, Eckmann L, Reed S. Chemokine secretion of human cells in response to *Toxoplasma gondii* infection. *Infect Immun* 1999; 67: 1547-1552.
18. Nilsson G, Mikovits JA, Metcalfe DD, Taub DD. Mast cell migratory response to interleukin-8 is mediated through interaction with chemokine receptor CXCR2/interleukin-8RB. *Blood* 1999; 93: 2791-2797.
19. Im SJ, Ahn MH, Han IH, Song HO, Kim YS, Kim HM, Ryu JS. Histamine and TNF- α release by rat peritoneal mast cells stimulated with *Trichomonas vaginalis*. *Parasite* 2011; 18: 49-55.
20. Seabra SH, de Souza W, DaMatta RA. *Toxoplasma gondii* partially inhibits nitric oxide production of activated murine macrophages. *Exp Parasitol* 2002; 100: 62-70.
21. Lüder CG, Algner M, Lang C, Bleicher N, Gross U. Reduced expression of the inducible nitric oxide synthase after infection with *Toxoplasma gondii* facilitates parasite replication in activated murine macrophages. *Int J Parasitol* 2003; 33: 833-844.
22. Pakianathan DR, Kuhn RE. *Trypanosoma cruzi* affects nitric oxide production by murine peritoneal macrophages. *J Parasitol* 1994; 80: 432-437.
23. Proudfoot L, Nikolaev AV, Feng GJ, Wei WQ, Ferguson MA, Brimacombe JS, Liew FY. Regulation of the expression of nitric oxide synthase and leishmanicidal activity by glycoconjugates of *Leishmania* lipophosphoglycan in murine macrophages. *Proc Natl Acad Sci USA* 1996; 93: 10984-10989.