

Seroprevalence of *Toxoplasma gondii* among School Children in Pyin Oo Lwin and Naung Cho, Upper Myanmar

Thị Lam Thái^{1,2,†}, Hojong Jun^{3,†}, Seo-Hye Park³, Hương Giang Lê^{1,2}, Jinyoung Lee³, Seong Kyu Ahn³, Jung-Mi Kang^{1,2}, Moe Kyaw Myint⁴, Khin Lin⁴, Woon-Mok Sohn¹, Ho-Woo Nam⁵, Byoung-Kuk Na^{1,2,*}, Tong-Soo Kim^{3,*}

¹Department of Parasitology and Tropical Medicine, and Institute of Health Sciences, Gyeongsang National University School of Medicine, Jinju 52727, Korea; ²BK21Plus Team for Anti-aging Biotechnology and Industry, Department of Convergence Medical Science, Gyeongsang National University, Jinju 52727, Korea; ³Department of Tropical Medicine, and Inha Research Institute for Medical Sciences, Inha University School of Medicine, Incheon 22212, Korea; ⁴Department of Medical Research Pyin Oo Lwin Branch, Pyin Oo Lwin, Myanmar; ⁵Department of Parasitology, College of Medicine, Catholic University of Korea, Seoul 06591, Korea

Abstract: *Toxoplasma gondii* is an apicomplexan parasite that can cause toxoplasmosis in a wide range of warm-blooded animals including humans. In this study, we analyzed seroprevalence of *T. gondii* among 467 school children living in the rural areas of Pyin Oo Lwin and Naung Cho, Myanmar. The overall seroprevalence of *T. gondii* among school children was 23.5%; 22.5% of children were positive for *T. gondii* IgG, 0.4% of children were positive for *T. gondii* IgM, and 0.6% of children were positive for both *T. gondii* IgG and IgM. Geographical factors did not significantly affect the seroprevalence frequency between Pyin Oo Lwin and Naung Cho, Myanmar. No significant difference was found between males (22.2%) and females (25.0%). The overall seroprevalence among school children differed by ages (10 years old [13.6%], 11-12 years old [19.8%], 13-14 years old [24.6%], and 15-16 years old [28.0%]), however, the result was not significant. Polymerase chain reaction analysis for *T. gondii* B1 gene for IgG-positive and IgM-positive blood samples were negative, indicating no direct evidence of active infection. These results collectively suggest that *T. gondii* infection among school children in Myanmar was relatively high. Integrated and improved strategies including reinforced education on toxoplasmosis should be implemented to prevent and control *T. gondii* infection among school children in Myanmar.

Key words: *Toxoplasma gondii*, seroprevalence, IgG, IgM, school child, Myanmar

Toxoplasma gondii is an ubiquitous apicomplexan parasite that can cause toxoplasmosis in a wide range of warm-blooded animals including humans [1,2]. It is prevalent in most areas of the world, and approximately 1/3 of the human population is estimated to be chronically infected with this parasite. The global prevalence of *T. gondii* human infection varies widely depending on geographic and climatic factors, socioeconomic status, cultural habits and the nutritional customs [3]. Humans are usually infected with the parasite by ingesting tissue cysts in undercooked or raw meat or by consuming food or water contaminated with oocysts shed by cats [4,5]. Clinically serious *T. gondii* infection can be acquired by congenital transmission from mother to fetus through the placenta during pregnancy [4,5]. Blood transfusion and organ transplanta-

tion are also potential routes of transmission [1,2]. *T. gondii* infection is usually asymptomatic or induces only mild clinical symptoms in immunocompetent adults and children; however, it can result in diverse clinical manifestations such as lymphadenitis, meningoencephalitis, or ocular toxoplasmosis in immunocompromised patients. Congenital transmission can lead to serious clinical consequences such as intracranial calcification, mental retardation, chronic chorioretinitis, hydrocephalus, and even fetal death [2,6].

Serological tests for *T. gondii*-specific antibodies are one of the most commonly used laboratory diagnostic methods for toxoplasmosis as an initial screening approach. Immunoglobulin M (IgM) antibody has been proposed as a marker of acute infection as this antibody usually appears in 1 week after the initial exposure [7]. IgM antibody level increases in a short period, declines gradually, and maintains for several months. On the other hand, IgG antibody is usually produced at several weeks after the initial infection and provides long-term protection. Therefore, the detection of specific IgM antibodies followed by specific IgG antibodies has been established for the

•Received 17 March 2019, revised 6 April 2019, accepted 25 April 2019.

*Corresponding authors (bkna@gnu.ac.kr; tongsookim@inha.ac.kr)

†These authors contributed equally to this study.

© 2019, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

diagnosis of recently acquired toxoplasmosis [7]. Seroprevalence studies of *T. gondii* infection among school children have been conducted in several countries including Brazil, China, Iran, Mexico, Nigeria, the Republic of the Marshall Islands, the United Arab Emirates (UAE) and the United States of America (USA) [8-16]. It is important to conduct prevalence study in school children since the infected children can manifest symptomatic toxoplasmosis later in life, even if they do not present symptoms at present. Nevertheless, serologic studies of *T. gondii* infection in school children are limited in Myanmar. This study aimed to evaluate the seroprevalence of *T. gondii* among school children living in the rural areas of Pyin Oo Lwin and Naung Cho in Myanmar on 2017.

A total of 467 blood samples were collected from school children in 4 basic education high schools in Pyin Oo Lwin (22°02'21.04" N, 96°28'17.97" E), Mandalay Division, and 3 basic education high schools in Naung Cho (22°19'52.76" N, 96°48'00.97" E), Shan State, Myanmar, in August and December 2017 (Fig. 1). Written informed consent was obtained from all children or legal guidances before blood collection. Basic information, including gender, address, and age were also obtained. The age of the children enrolled in this study ranged from 10 to 16 years old. Approximately 1 ml of venous blood was drawn from each participant, labeled individually,

and cooled with ice packs to maintain the temperature at 4°C during transportation to the laboratory. The serum was separated from the whole blood by centrifugation at 3,000 rpm for 5 min and stored at -76°C until use. The study protocol was reviewed and approved by the Ethics Committee of the Ministry of Health, Myanmar (97/Ethics 2015), and the Biomedical Research Ethics Review Board of Inha University School of Medicine, Republic of Korea (INHA 15-013).

All serum samples were screened for anti-*T. gondii* IgG and IgM antibodies by enzyme-linked immunosorbent assay (ELISA). The ELISA kits used in this study were the Human Anti-*Toxoplasma gondii* IgG ELISA Kit (Abcam, Cambridge, UK) and the Human Anti-*Toxoplasma gondii* IgM ELISA Kit (Abcam). All ELISA processes were performed following the manufacturer's instructions. The optical density (OD) values were read at 450 nm using the Multiskan™ FC Microplate Photometer (Thermo Scientific, Waltham, Massachusetts, USA). An OD value similar to or greater than the threshold value was considered positive.

Blood samples were further analyzed by molecular method to detect active *T. gondii* infection. Genomic DNA was extracted from 200 µl of whole blood samples using the QIAamp Blood Kit (Qiagen, Valencia, California, USA) according to the manufacturer's instructions. The DNA samples were evaluated




Fig. 1. Map of the study areas. Community-based survey for anti-*Toxoplasma gondii* antibodies was performed.

by polymerase chain reaction (PCR) targeting the *T. gondii* B1 gene using the primers, Toxo-B5 (5'-TGAAGAGAGGAAACAG-GTGGTCC-3') and Toxo-B6 (5'-CCGCCTCCTTCGTCGGTCG-TA-3') [17]. The amplification step included 1 cycle of denaturation at 94°C for 5 min, 35 cycles of denaturation at 94°C for 1 min, annealing at 60°C for 1 min, and extension at 72°C for 1 min, and a final extension step at 72°C for 3 min. In each PCR amplification step, genomic DNA extracted from *T. gondii* RH strain was included as a positive control. The PCR products were subjected to electrophoresis on a 2% agarose gel, stained with RedSafe™ Nucleic Acid Staining Solution (Intron, Seongnam, Korea), and visualized under a UV transilluminator (Analytik Jena US LLC, Upland, California, USA).

Statistical analysis was performed with the SPSS v.20.0 software package (SPSS Inc., Armonk, New York, USA). The relative proportions were calculated using a confidence interval (CI) of 95%. The Pearson's chi-square and Fisher's exact tests were used to analyze the differences in the seroprevalences of anti-*T. gondii* IgG and IgM according to sex, age, and area. Odds ratio (OR) with their 95% CI for various risk factors associated with seropositivity were estimated and *P*-values of <0.05 were considered statistically significant.

A total of 467 blood samples collected from school children aged 10-16 years in Myanmar were analyzed for anti-*T. gondii* IgG and IgM by ELISA. The overall seroprevalence of *T. gondii*

was 23.5%; 22.5% (95% CI = 18.9-26.4) of children were positive for anti-*T. gondii* IgG antibodies, 0.4% (95% CI = 0.1-1.4) of children were positive for IgM, and 0.6% (95% CI = 0.2-1.7) of children were positive for both IgG and IgM (Table 1). There was no significant difference in the seroprevalence of *T. gondii* between males (22.2%) and females (25.0%) (*P* = 0.480) (Table 2). The overall seroprevalence of *T. gondii* was increased with increasing age (Table 2). The prevalence (28.0%, OR = 2.5, 95% CI = 0.7-9.0) among older school children (15-16-year-olds) showed highest positive serology compared with other groups, 10-year-olds (13.6%), 11-12-year-olds (19.8%), and 13-14-year-olds (24.6%). The seroprevalence of *T. gondii* among school children in Naung Cho (25.0%) was slightly higher than that among school children in Pyin Oo Lwin (22.6%); however, the result was not statistically significant (*P* = 0.497). There was no positive amplification of the *T. gondii* B1 in all of the seropositive samples.

Myanmar is a country in a subtropical area, and various kinds of parasitic diseases are highly prevalent like other Southeast Asian countries. High humidity and warm climate, which may affect the survival of oocysts of *T. gondii* in the environment [18], may provide the favorable conditions for the development of *T. gondii*. However, only a few serological studies have been conducted for toxoplasmosis in Myanmar. In this study, the seroprevalence of *T. gondii* among school

Table 1. Seroprevalence of anti-*Toxoplasma gondii* IgG and IgM antibodies in 467 school children in Naung Cho and Pyin Oo Lwin, Myanmar

Antibodies	No. of tested	Positive rate (%)	Confidence interval (95%)
IgG ⁺ , IgM ⁻	105	22.5	18.9-26.4
IgG ⁺ , IgM ⁺	3	0.6	0.2-1.7
IgG ⁻ , IgM ⁺	2	0.4	0.1-1.4
IgG ⁻ , IgM ⁻	357	76.5	72.4-80.1
Total	467	100.0	

+, positive; -, negative.

Table 2. Socio-demographic characteristics and seroprevalence of *Toxoplasma gondii* antibody among school children in Naung Cho and Pyin Oo Lwin, Myanmar

Variables	Group	No. of tested	Positive rate (%)	Odds ratio (95% CI)	<i>P</i> -value
Gender	Male	243	22.2	Referent	0.48
	Female	224	25.0	1.2 (0.8-1.8)	
Age	10	22	13.6	Referent	0.34
	11-12	121	19.8	1.6 (0.4-5.7)	
	13-14	224	24.6	2.1 (0.6-7.2)	
	15-16	100	28.0	2.5 (0.7-9.0)	
Geographical region	Naung Cho	149	25.0	Referent	0.497
	Pyin Oo Lwin	318	22.6	0.9 (0.5-1.3)	

children living in the rural areas of Pyin Oo Lwin and Naung Cho in Myanmar was investigated on 2017. The prevalence of *T. gondii* infection is different between countries or communities even in the same region [19]. In this study, the overall seroprevalence of *T. gondii* among the school children in this study was 23.5% (110/467), which is comparable to that in other studies involving primary school children in southern Nigeria (24.0%), school children in Iran (22.0%), and high school students in Mexico (22.4%) [9,10,20]. Meanwhile, the prevalence in Myanmar was higher than that in other countries including the USA (10.8%), the UAE (12.5%), China (16.0%), Cyrus (6.5%), and Brazil [14], but much lower than that in West Africa (63.1%) [11] and the Republic of the Marshall Islands (54.8%) [8,13-15,21]. The discrepancy in seroprevalence rates between different countries may be attributed to differences in ethnicity, tradition/culture, and food habits [2,3]. Different serodiagnostic methods and sample size may also affect the overall seroprevalence rate of each country. However, the results of this study suggested that *T. gondii* infection among school children in Myanmar should not be ignored. A previous study of the seroprevalence rates of *T. gondii* among school children in Myanmar in 1977 also revealed high levels of *T. gondii* antibodies in school children in Hlawga (43.8%) and Gyogon (28.4%) [22]. It is generally known that there is no gender-related difference in the seroprevalence of toxoplasmosis [2,23]. In the present study, the seroprevalence of *T. gondii* among female children (25.0%) was slightly higher than that among male children (22.2%), but not significant ($P=0.480$). This may be attributed to the similar routes of *T. gondii* infection shared by male and female school children through common risk factors, such as playing in the soil and water or consuming undercooked meat and vegetables. The prevalence of anti-*T. gondii* antibodies is also known to increase with age [2,10,13,15,23]. Consistent with the previous studies, the overall seroprevalence of *T. gondii* among Myanmar school children was increased with age.

The presence of cats, hygienic conditions, sociocultural habits, and environmental conditions, are closely related to the prevalence of *T. gondii* [2,23]. In this study, there was no systematic questionnaire for the school children to access risk factors for *T. gondii* infection. However, young children are usually vulnerable to *T. gondii* infection due to their behavioral habits, which involve a higher environmental exposure and a lack of awareness for avoiding the risk of infection [12-15]. Substantial levels of *T. gondii* infections were reported in domestic

goats (11.4%) and wild insectivorous bats (29.3%) [24,25] in Myanmar, suggesting widespread *T. gondii* infection among domestic and wild animals in Myanmar. The ingestion of undercooked meat and unpasteurized milk and cheese from *T. gondii*-infected goats can be a potential cause of infection in humans [26]. Therefore, direct or indirect contact with stray cats or animals, poor hygiene conditions, and consuming undercooked meat and raw vegetables may contribute to *T. gondii* infection among school children in Myanmar. Congenital transmission cannot be ruled out as a potential source of infection considering the high seroprevalence rate of *T. gondii* among Myanmar pregnant women [27,28]. The seroprevalence of *T. gondii* among refugee and migrant Myanmar pregnant women along the Thailand-Myanmar border reported as 31.7% [28]. The seroprevalence among Myanmar pregnant women in Yangon was reported as 30.70% (95% CI=27.9-37.2) [27]. A lack of awareness of toxoplasmosis is likely to be one of the most key factors contributing to the high prevalence among Myanmar pregnant women. Young girls who are infected with *T. gondii* in their childhood are not at risk for the transmission of *T. gondii* at the present stage; however, *T. gondii* infection can cause transplacental transmission when they are pregnant as adults. Therefore, more attention should be given to young girls to prevent any kind of potential transmission in the future. With the rapid spread of HIV/AIDS in Southeast Asian countries [29], toxoplasmosis should be controlled because it can lead to life-threatening or fatal conditions. Further studies are necessary to understand the risk factors and possible transmission routes of *T. gondii* infection in Myanmar school children. Reinforced education to enhance awareness of toxoplasmosis and prevent transmission of the disease also would be necessary for school children in Myanmar.

In conclusion, this is the first community-based survey to analyze the seroprevalence of *T. gondii* among school children in Myanmar. The present study demonstrated the prevalence of *T. gondii* among the school children with an overall seroprevalence rate of 23.5%. This study highlights the urgent necessity of integrated and improved strategies including public health education for the prevention and control of *T. gondii* infection in young school children, especially young girls. Additional larger community-based studies in a wide range of geographical locations are also needed to clarify the prevalence rates of toxoplasmosis among school children in Myanmar.

ACKNOWLEDGMENTS

The authors would like to thank the staffs in the Department of Medical Research Pyin Oo Lwin Branch and the health professionals in Naung Cho and Pyin Oo Lwin for their contribution and technical support in blood sampling. This study was supported by the National Research Foundation of Korea grant funded by the Korean government (NRF-2017M3A9B8069530).

CONFLICT OF INTEREST

The authors declare that they have no competing interests.

REFERENCES

1. Tenter AM, Heckeroth AR, Weiss LM. *Toxoplasma gondii*: from animals to humans. *Int J Parasitol* 2000; 30: 1217-1258.
2. Montoya JG, Liesenfeld O. Toxoplasmosis. *Lancet* 2004; 363: 1965-1976.
3. Flegr J, Prandota J, Sovicková M, Israili ZH. Toxoplasmosis-global threat. Correlation of latent toxoplasmosis with specific disease burden in a set of 88 countries. *PLoS One* 2014; 9: e90203.
4. Kapperud G, Jenum PA, Stray-Pedersen B, Melby KK, Eskild A, Eng J. Risk factors for *Toxoplasma gondii* infection in pregnancy. Results of a prospective case-control study in Norway. *Am J Epidemiol* 1996; 144: 405-412.
5. Jones JL, Muccioli C, Belfort R Jr, Holland GN, Roberts JM, Silveira C. Recently acquired *Toxoplasma gondii* infection, Brazil. *Emerg Infect Dis* 2006; 12: 582-587.
6. Ronman E, Zamir CS, Rilkis I, Ben-David H. Congenital toxoplasmosis-prenatal aspects of *Toxoplasma gondii* infection. *Reprod Toxicol* 2006; 4: 458-472.
7. Robert-Gangneux F, Dardé ML. Epidemiology of and diagnostic strategies for toxoplasmosis. *Clin Microbiol Rev* 2012; 25: 264-296.
8. Abu-Zeid YA. Serological evidence for remarkably variable prevalence rates of *Toxoplasma gondii* in children of major residential areas in United Arab Emirates. *Acta Trop* 2002; 83: 63-69.
9. Galván-Ramírez ML, Pérez LR, Ledesma Agraz SY, Sifuentes Ávila LM. Seroepidemiology of toxoplasmosis in high-school students in the metropolitan area of Guadalajara, Jalisco, Mexico. *Sci Medica* 2010; 20: 59-63.
10. Sharif M, Daryani A, Barzegar G, Nasrolahei M. Seroepidemiological survey for toxoplasmosis among schoolchildren of Sari, Northern Iran. *Trop Biomed* 2010; 27: 220-225.
11. Fan CK, Lee LW, Liao CW, Huang YC, Lee YL, Chang YT, da Costa Âdos S, Gil V, Chi LH, Nara T, Tsubouchi A, Akinwale OP. *Toxoplasma gondii* infection: relationship between seroprevalence and risk factors among primary schoolchildren in the capital areas of Democratic Republic of São Tomé and Príncipe, West Africa. *Parasit Vectors* 2012; 5: 141.
12. Muñoz-Zanzi C, Williams-Nguyen J, Belongia EA, Belongia EA. A sero-survey of toxoplasmosis in farm and non-farm children from Wisconsin, United States, 1997-1999. *BMC Public Health* 2013; 13: 837.
13. Fu CJ, Chuang TW, Lin HS, Wu CH, Liu YC, Langinlur MK, Lu MY, Hsiao WW, Fan CK. *Toxoplasma gondii* infection: seroprevalence and associated risk factors among primary school children in the capital area of the Republic of the Marshall Islands. *Jpn J Infect Dis* 2014; 67: 405-410.
14. Marchioro AA, Colli CM, Ferreira E'C, Viol BM, Araujo SM, Favavigna-Guilherme AL. Risk factors associated with toxoplasmosis and toxocaríasis in populations of children from nine cities in southern Brazil. *J Helminthol* 2015; 89: 428-432.
15. Xin KS, Liu H, Wang HB, Yao ZL. Seroprevalence of *Toxoplasma gondii* among primary school children in Shandong Province, China. *Korean J Parasitol* 2015; 53: 489-492.
16. Yang N, Wang D, Xing M, Li C, Li J, Wu A, Sang X, Feng Y, Jiang N, Chen Q. Seroepidemiology and risk factors of *Toxoplasma gondii* infection among the newly enrolled undergraduates and post-graduate students in China. *Front Microbiol* 2017; 8: 2092.
17. Gontijo da Silva M, Clare Vinaud M, de Castro AM. Prevalence of toxoplasmosis in pregnant women and vertical transmission of *Toxoplasma gondii* in patients from basic units of health from Gurupi, Tocantins, Brazil, from 2012 to 2014. *PLoS One* 2015; 10: e0141700.
18. Meerburg BG, Kijlstra A. Changing climate-changing pathogens: *Toxoplasma gondii* in North-Western Europe. *Parasitol Res* 2009; 105: 17-24.
19. Pappas G, Roussos N, Falagas ME. Toxoplasmosis snapshots: global status of *Toxoplasma gondii* seroprevalence and implications for pregnancy and congenital toxoplasmosis. *Int J Parasitol* 2009; 39: 1385-1394.
20. Gyang VP, Akinwale OP, Lee YL, Chuang TW, Orok A, Ajibaye O, Liao CW, Cheng PC, Chou CM, Huang YC, Fan KH, Fan CK. *Toxoplasma gondii* infection: seroprevalence and associated risk factors among primary schoolchildren in Lagos City, Southern Nigeria. *Rev Soc Bras Med Trop* 2015; 48: 56-63.
21. Liassides M, Christodoulou V, Moschandreas J, Karagiannis C, Mitis G, Koliou M, Antoniou M. Toxoplasmosis in female high school students, pregnant women and ruminants in Cyprus. *Trans R Soc Trop Med Hyg* 2016; 110: 359-366.
22. Tin F. Serological responses to *Plasmodium* and *Toxoplasma* in school children from two areas in Burma. *Southeast Asian J Trop Med Public Health* 1977; 8: 552-557.
23. Wilking H, Thamm M, Stark K, Aebischer T, Seeber F. Prevalence, incidence estimations, and risk factors of *Toxoplasma gondii* infection in Germany: a representative, cross-sectional, serological study. *Sci Rep* 2016; 6: 22551.
24. Bawm S, Maung WY, Win MY, Thu MJ, Chel HM, Khaing TA, Wai SS, Htun LL, Myaing TT, Tiwananthagorn S, Igarashi M, Karakura K. Serological survey and factors associated with *Toxoplasma gondii* infection in domestic goats in Myanmar. *Scientifica*

- (Cairo) 2016; 2016: 4794318.
25. Sun H, Wang Y, Zhang Y, Ge W, Zhang F, He B, Li Z, Fan Q, Wang W, Tu C, Li J, Liu Q. Prevalence and genetic characterization of *Toxoplasma gondii* in bats in Myanmar. *Appl Environ Microbiol* 2013; 79: 3526-3528.
 26. Dubey JP, Verma SK, Ferreira LR, Oliveira S, Cassinelli AB, Ying Y, Kwok OC, Tuo W, Chiesa OA, Jones JL. Detection and survival of *Toxoplasma gondii* in milk and cheese from experimentally infected goats. *J Food Prot* 2014; 77: 1747-1753.
 27. Andiappan H, Nissapatom V, Sawangjaroen N, Nyunt MH, Lau YL, Khaing SL, Aye KM, Mon NC, Tan TC, Kumar T, Onichandran S, bin Mat Adenana NA. Comparative study on *Toxoplasma* infection between Malaysian and Myanmar pregnant women. *Parasit Vectors* 2014; 7: 564.
 28. van Enter BJD, Lau YL, Ling CL, Watthanaworawit W, Sukthana Y, Lee WC, Nosten F, McGready R. Seroprevalence of *Toxoplasma gondii* infection in refugee and migrant pregnant women along the Thailand-Myanmar border. *Am J Trop Med Hyg* 2017; 97: 232-235.
 29. Pendse R, Gupta S, Yu D, Sarkar S. HIV/AIDS in the South-East Asia region: progress and challenges. *J Virus Erad* 2016; 2 Suppl 4: 1-6.