

Development of a Molecular Marker Linked to the *A4* Locus and the Structure of HD Genes in *Pleurotus eryngii*

Song Hee Lee^a, Asjad Ali^b, Byeongsuk Ha^b, Min-Keun Kim^b, Won-Sik Kong^c and Jae-San Ryu^a

^aDepartment of Mushroom Science, Korea National College of Agriculture and Fisheries, Jeonju, Republic of Korea; ^bEnvironment-Friendly Research Division, Gyeongsangnam-do Agricultural Research and Extension Services, Jinju, Republic of Korea; ^cMushroom Research Division, National Institute of Horticultural & Herbal Science, Rural Development Administration, Eumseong, Republic of Korea

ABSTRACT

Allelic differences in *A* and *B* mating-type loci are a prerequisite for the progression of mating in the genus *Pleurotus eryngii*; thus, the crossing is hampered by this biological barrier in inbreeding. Molecular markers linked to mating types of *P. eryngii* KNR2312 were investigated with randomly amplified polymorphic DNA to enhance crossing efficiency. An *A4*-linked sequence was identified and used to find the adjacent genomic region with the entire motif of the *A* locus from a contig sequenced by PacBio. The sequence-characterized amplified region marker 7-2₉₉ distinguished *A4* mating-type monokaryons from KNR2312 and other strains. A BLAST search of flanked sequences revealed that the *A4* locus had a general feature consisting of the putative *HD1* and *HD2* genes. Both putative HD transcription factors contain a homeodomain sequence and a nuclear localization sequence; however, valid dimerization motifs were found only in the HD1 protein. The ACAAT motif, which was reported to have relevance to sex determination, was found in the intergenic region. The SCAR marker could be applicable in the classification of mating types in the *P. eryngii* breeding program, and the *A4* locus could be the basis for a multi-allele detection marker.

ARTICLE HISTORY

Received 4 April 2019
Accepted 12 May 2019

KEYWORDS

Mating-type gene; *A4* locus; *Pleurotus eryngii*; SCAR marker; homeodomain

1. Introduction


The sexual development of mushrooms starts with the mating step and progresses through haploid and diploid phases. A dikaryotic mycelium results from the fusion of two genetically distant monokaryons, and the two nuclei coexist during vegetative growth and fruiting body formation until the onset of the maturation stage, when these nuclei fuse to form basidia and then produce haploid spores [1,2]. The mating-type loci that control developmental pathways and play a role in maintaining a fertile dikaryon are termed *A* and *B* [3]. Similarly, mating compatibility in *Pleurotus eryngii* is determined by two multiallelic mating-type loci, *A* and *B* [4]. The developmental sequence leading to the formation of the dikaryon and further to the binucleate cells that characterize it are present in both sets of mating-type genes [5]. The initiation of *A*-regulated steps in dikaryon development begins with compatible mating to generate the active transcription factor complex [6], while *B* genes encode lipopeptide pheromones and pheromone receptors [7]. Previously, Ryu et al. [8] classified mating-type factors for *A* and *B* alleles by

crossing monokaryons with testers of each allele, thereby a SCAR marker linked to the *B3* locus was developed to determine the mating type of *P. eryngii*.

Since ~25% of the world's production of cultivated fungi is the production of edible oyster mushrooms, strain improvement through breeding might ease the production procedure with enhanced knowledge about their mating-type loci [9]. *P. eryngii*, king oyster mushroom, is among the highly demanded mushrooms in Asia, Europe and the USA, and this species has been grown on a commercial scale in Italy, China, Japan, and Korea [10]. Immunostimulatory, antiproliferative, and antifungal actions against cells make this mushroom medically valuable [11]. *P. eryngii* species are also famous for their flavor and relatively low-cost cultivation methods. Researchers have focused on obtaining high yielding strains using strain selection and improvements in cultivation technology [12], but limited research has been reported on mating-type specificity [4,8].

The DNA marker approach has benefited modern plant breeding in different ways, such as genotype identification, gene isolation, and the analysis of

CONTACT Jae-San Ryu  coolmush88@gmail.com

 Supplemental data for this article can be accessed [here](#).

different agronomic traits. Marker-assisted selection (MAS) has been used in mushroom breeding to distinguish cultivars to protect breeders' rights [13], and its use in relation to agronomic traits for breeding has made MAS a fast, easy, and inexpensive method [14]. SCAR markers are identified by PCR using specifically designed oligo-nucleotide primers based on sequence data. These markers have been reported as a suitable tool for routine diagnostics [15].

In the present study, we demonstrate the molecular structure of the *A4* locus and the development of a SCAR marker specifically linked to this locus.

2. Materials and methods

2.1. Strains and growth conditions

All of the strains used in this study were derived from the previous work by Ryu et al. [8,16] (Supplemental Table S1). Two compatible neohaplonts, P5 and P6, were isolated from dikaryotic *P. eryngii* KNR2312 by dedikaryotization through homogenization of the protoplast with a lysing enzyme. Mycelial mats were grown in MCM (mushroom complete medium) broth as described by a previous report [8] and were then harvested, freeze-dried, and ground with mortar and pestle for DNA extraction.

2.2. Genomic DNA extraction, RAPD analysis, and genome sequencing

Genomic DNA was extracted from the lyophilized mycelia using GenEx plant plus! (GeneAll, Seoul, Korea). PCR was performed with RAPD to screen the mating-type-specific bands according to the method in a previous report [17] with minor modifications. Four gDNAs of monokaryons with the same mating types (*A3B3*, *A3B4*, *A4B3*, and *A4B4*) derived from *P. eryngii* KNR2312 were bulked and adjusted to 30 ng/ul for bulked segregant analysis (BSA). Monokaryons in bulked gDNA showing mating-type-specific bands with random primers were subjected to PCR for individual monokaryons with the same primers. PCRs were conducted in a 10 μ L mixture containing 30 ng of template genomic DNA, 0.2 mM dNTPs, 0.25 U of *e-Taq* DNA polymerase (SolGent, Daejeon, Korea), 1 \times buffer containing 2.5 mM Tris-HCl (pH 8.2) and 1.5 mM MgCl₂, and 0.25 pmol of each 10-mer random oligonucleotides including L series (Operon Technologies, Alameda, CA). Amplification conditions were set as follows: initial denaturation for 4 min at 95 °C, followed by 35 cycles of a 1 min denaturation step at 94 °C, 1 min annealing at 37 °C, and 1 min and 30-sec extension at 72 °C. PCR

products were run on 3% (w/v) agarose gels in TAE buffer (400 mM Tris, 200 mM sodium acetate, and 20 mM EDTA, pH 8.3), stained Safeview classic (iNtRON Biotechnology, eongnam, Korea) and visualized under ultraviolet light. The 100 bp plus DNA ladder (Bioneer, Daejeon, Korea) was used as a standard size marker.

We prepared gDNA samples from KNR2312P6 (neohaplont) into SMRTbell DNA template libraries of 20-kb average insert size according to the manufacturer's specification, and Covaris G-tubes were used for fragmentation. SMRT sequencing was carried out on the PacBio RS II platform according to standard protocols, with the XL binding kit used in conjunction with the C4 sequencing kit. All runs were performed with diffusion-based loading and analyzed using standard primary data analysis. Contigs were assembled using CANU v1.7 with long reads.

2.3. SCAR marker development and data analysis

For isolation and identification of DNA in the gel, the mating-type-specific bands were cut out and purified by Gel Elution Kit (Solgent) and directly sequenced on an ABI 377 sequencer (Macrogen, Seoul, Korea) using the Big-Dye cycle sequencing kit (Applied Biosystems, Carlsbad, CA). The *A4* mating-type-specific sequence, designated 7-2, was used to find the 7-2 adjacent genomic region and the entire motif of the *A4* locus. The SCAR primer was manually designed using the 7-2 sequence and a flanked region (~1 kb) obtained from a contig of KNR2312P6 sequenced by PacBio. The developed primer sequences were F: 5'-AATCACGGGAAGATCTGGTG-3' and loci-R: 5'-GTGGTAGGGTCCC GCCT-3'. PCR conditions for the SCAR marker were optimized as follows: initial denaturation at 98 °C for 30 sec, followed by 35 cycles of a 10 sec denaturation at 98 °C, 15 sec annealing at 70 °C, and 10 sec extension at 72 °C, and a final extension of 10 min at 72 °C. The primer set was applied to determine the mating type of monokaryons from KNR2312 and other strains whose mating types were different (Supplemental Table S1) in addition to the *Pleurotus* genus.

2.4. Gene prediction and sequence alignment

The sequence including the *A4* allele was obtained from a contig of KNR2312P6 using a sequence search with DNAMAN (Lynnon Biosoft, Quebec, Canada) with an *A4*-specific genomic sequence. *A4* sequence has been deposited in the National Center for Biotechnology Information (NCBI) database and

assigned the Accession Number MK522809. Sequences were analyzed using FGENESH (<http://www.softberry.com/>) and NCBI Conserved Domain Search [18] to predict the genes present and their functions. COILS (window width 14 and MTIDK matrix) [19] and NLStradamus (prediction cutoff, 0.5) [20] were adapted for coiled-coil domains (putative dimerization motif) and nuclear localization signals, respectively. Genomic data of *A* mating-type genes for *P. eryngii* KNR2312P5, *P. eryngii* ATCC90797, *P. tuoliensis* CCMSSC00489 and CCMSSC00485, and *P. ostreatus* PC15 were obtained from an edible mushroom DB (<http://112.220.192.2/per/>) [21], NCBI's GenBank and Joint Genomic Institute (<https://genome.jgi.doe.gov/portal/>). Alignment analysis was performed with DNAMAN (Lynnon Biosoft) with default parameters to determine the nucleotide frequency, including the transcription initiation site.

3. Results

3.1. *A4* mating-type-linked SCAR marker

Among 200 primers, several primers were segregated with mating types (data not shown), including the OPL-07 primer specifying *A4Bx* by an ~500 bp PCR product (designated 7-2) (Figure 1). All RAPD markers showing mating-type specificity were converted into SCAR markers based on their sequences. Most SCAR markers showed low or no mating-type specificity (data not shown). The marker from 7-2 was unstable and weak; thus, a more accurate marker was designed based on the flanked sequences of 7-2 obtained from a contig that was one of the 498 contigs (~50 Mb) of KNR2312P6 sequenced by PacBio. Cosegregation of the specific 299 bp bands with the *A4Bx* locus was observed in KNR2312-derived monokaryons with the SCAR marker (designated 7-2₂₉₉). A critical annealing temperature of 70 °C was determined to best identify the mating type. Thus, to observe the strong linkage of 7-2₂₉₉ with the *A4Bx* locus, 17 monokaryons from KNR2312 were tested. Eight monokaryotic isolates showed a specific band for mating-type *A4B3*,

A4B4, and P6 (*A4B3*), while no amplification was detected for *A3B3*, *A3B4*, and P5 (*A3B4*) (Figure 2). All 98 monokaryons from KNR2312 [8] showed consistent results of the 299 bp genotype coinciding with the *A4Bx* phenotype at all times (data not shown).

3.2. Application of the SCAR marker (7-2₂₉₉) to other strains of *P. eryngii* and other species

Different monokaryons derived from KNR2501, KNR2522, and KNR2523 were used to evaluate the suitability of the 7-2₂₉₉ marker. Because the *A4* allele was shared in KNR2523 and KNR2522 but not in KNR2501 [8], 8 monokaryons, two from each mating type, were selected for the new marker. The 7-2₂₉₉ marker amplified unique bands from KNR2523(2)-15, 34 (*A4B3*) and KNR2523(2)-24, 26 (*A4B12*) but not from the remaining strains (KNR2523(2)-6 and -28 for *A12B3* and KNR2523(2)-23 and -32 for *A12B12*). Similarly, a unique band of 299 bp was observed from KNR2522(2)-1, 6 (*A4B12*) and KNR2522(2)-7, 15 (*4B3*). No band was shown in the strains carrying the *A12B12* and *A12B3* mating types in addition to the KNR2501(2)-derived monokaryons with *A3B5*, *A5B5*, *A3B4*, and *A5B4* (Figure 3). A 299 bp band was amplified from all monokaryons with *A4Bx* and P6 (*A4B3*), which indicated its suitability for the detection of the *A4Bx* locus in other strains of *P. eryngii*. Some other mushroom fungi of the same genus, such as *P. florida*, *P. sajor-caju*, *P. salmoneostamineus*, *P. cornucopiae*, and *P. ostreatus*, were also used to examine the presence of the *A4Bx* locus with the 7-2₂₉₉ marker, but none of them were positive for the *A4Bx* locus (data not shown).

3.3. Molecular structure of the *A4* locus

The BLASTp search of the flanked sequence of the *A4*-specific marker identified two homeodomains (HD1, designated *Pea1*, and HD2, designated *Pea2*), located 200 kb apart from the marker. The two genes were located side-by-side, and the direction of




Figure 1. RAPD patterns for 16 monokaryons of *Pleurotus eryngii* KNR2312 with OPL-07 (10-mer). Box shows the *A4* specific band. M: 100 bp ladder; 30, 33, 37, 40: *A3B3*; 12, 25, 29, 32: *A3B4*; 7, 26, 35, 39, 49: *A4B3*; 11, 15, 17, 21: *A4B4*.




Figure 2. Amplification patterns with 7-2₉₉ (*A4* specific marker) in different sets of monokaryons derived from *Pleurotus eryngii* KNR2312. M: 100 bp ladder; Upper: KNR2312(3)-4, 13, 33, 37: *A3B3*; 6, 7, 8, 26: *A3B4*. Lower: 12, 29, 34, 42: *A4B3*; 8, 10, 20, 22: *A4B4*; P5: *A3B4*; P6: *A4B3*; D: dikaryon (P5 × P6).




Figure 3. SCAR marker amplification patterns for the *A4* locus in diverse phenotypes. Amplifications were performed with SCAR (7-2₉₉) primers using monokaryons derived from KNR2523, KNR2522, and KNR2501. Two monokaryons were selected from each mating type. (A) monokaryons from KNR2523(2) (6, 28: *A12B3*; 15, 34: *A4B3*; 23, 32: *A12B12*; and 24, 26: *A4B12*), P5: *A3B4*; P6: *A4B3*; D: dikaryon (P5 × P6); (B) monokaryons from KNR2522(2) (1, 6: *A4B12*; 5, 21: *A12B12*; 24, 26: *A12B3*; and 7, 15: *A4B3*); (C) monokaryons from KNR2501(2) (15, 35: *A3B4*; 1, 31: *A3B5*; 21, 39: *A5B4*; and 5, 27: *A5B5*).

transcription was outward from an intergenic region for both genes, similar to other basidiomycetes [22,23]. The two HD genes were located across 4.6 kb, *HD1* was assumed to be 2197 bp gDNA and 694 a.a. with two introns, while *HD2* was likely 2062 bp gDNA and 627 a.a. with 3 introns (Supplemental Figure S1). The two HD genes were

considered a general feature of HD in *Pleurotus* [23,24]; however, *P. eryngii* ATCC90797 and *P. ostreatus* PC15 contained three HD genes (Supplemental Figure S2). *Pea1* (HD1) showed high similarity, ranging from 51.9–70.4% with those of other *Pleurotus*, while *Pea2* (HD2) had high similarity with HD2 of *Pleurotus*, ranging from 69.0–49.6%




Figure 4. Molecular structure of the *A4* locus in *Pleurotus eryngii* KNR2312P6. The *A4* locus spans ~4.6 kb in length and consists of two homeodomain transcription genes with a 164 bp intergenic region. *Pea1* and *Pea2* consist of 2197 bp and 2062 bp gDNA and 694 aa and 627 aa lengths, respectively. Conserved sequences in the intergenic region are ACAAT, GCCG, and TCCCAC. The gray block in the intergenic region indicates the relative position of the conserved sequence area shown below. Bold indicates conserved sequences revealed by alignment analysis with that of another genus *Pleurotus* (see M & M). Overlapped conserved domains are adjusted block height for identification of locations. HD: homeodomain; Di: dimerization domain (coiled coil); NLS: nuclear localization sequence.

(Supplemental Table S2). The intergenic region between *Pea1* (HD1) and *Pea2* (HD2) was 163 bp long, which was similar to those of other *Pleurotus* species ranging from 166 to 178 bp (Supplemental Figure S3). Alignment of the intergenic regions among *Pleurotus* showed several conserved sequences, including two ACAAT motifs (one was reversed) (Figure 4), GCCG and TCCCAC. *Pea1* contained two NLS and three dimerization motifs (coiled-coil domain). Both genes had homeodomains located at the 129–168th a.a. (*Pea1*, 39 a.a. in length) and 150–206th a.a. (*Pea2*, 56 a.a. in length). Some NLS domains overlapped with coiled-coil motifs and homeodomains (Figure 4). Interestingly, coiled coils were not detected at high levels in *Pea2* (three with 0.021 as the highest value were detected). Multiple alignment with HD mating-type proteins from the genus *Pleurotus* showed two sectors: one had highly diverse sequences, and the other had highly similar sequences. Diverged sequences were terminated at the beginning of the homeodomain (Supplemental Figure S2).

4. Discussion

4.1. Molecular marker linked to the *A4* locus

The development of new desired strains and the methods to distinguish them are some of the main issues in mushroom breeding. Previously, different techniques were employed to verify strains such as small subunit ribosomal DNA (SSU rDNA) [25], multiple nuclear genes [26], or PCR-based DNA fingerprinting methodologies including RAPD, AFLP, and SCAR [27]. These marker techniques save time by identifying the mating type more reliably than traditional methods such as crossing. SCAR markers are comparatively stable and overcome the limitations of RAPD markers in the form of reproducibility. SCARs have been used for identification purposes in *Lentinula edodes* [14] and *Laccaria bicolor* [28]. In the present study, polymorphisms were frequently detected in the RAPD products of

the *A4Bx*-specific compatible monokaryons (*A4B3* and *A4B4*) (Figure 1). As SCAR markers can identify single bands instead of complex patterns, they are more accurate than RAPDs. Therefore, a SCAR marker (7-2₂₉₉) was developed to achieve well-verified and consistent results (Figures 2 and 3). Although the 7-2₂₉₉ marker is located 200 kb away from the *A4* locus, it cosegregated with the *A4* locus trait among all monokaryons from KNR2312 and a wide range of other strains. This result suggested that the inter-region between the 7-2₂₉₉ and *A4* loci exerts strong selection pressure. The genomic region around the *A* locus in the genus *Pleurotus* was reported to be conserved during long evolutions [24]. Thus, due to the high specificity, consistency, and wide range of applicability of the PCR-based method, the 7-2₂₉₉ marker might be an efficient tool for distinguishing *A4Bx*-locus strains in breeding programs for *P. eryngii*. The newly developed method might be preferred over the traditional method for the identification of mating-type, based on the shorter duration, i.e., 1–3 days instead of 15 days. Rapid colony PCR for mycelium is well established for massive selections [29]; therefore, breeders could save time and materials by mating only compatible strains with SCAR markers in the case of inbreeding. Moreover, *P. eryngii* KNR2312 has been used as a parent for many cultivars [30–32] due to a narrow genetic pool; thus, this marker will be more effective for breeding using KNR2312-derived cultivars.

4.2. Molecular structure of the *A4* locus

The homeodomain proteins HD1 and HD2 play a role in the regulation of sexual development in several fungi [33]. The *A4* locus showed a general feature of the *A* mating locus of the genus *Pleurotus* reported [23,24]. The two HD genes were located side-by-side and separated by a 163 bp intergenic region. Exceptions were the three HDs in *P. eryngii* ATCC90797 and *P. ostreatus* PC15, suggesting that

these strains are likely closer to a prototype. However, whether all HDs are linked to the *A* mating type is unknown. Alignment of the intergenic region revealed conserved sequences. ACAAT motifs were reported to be enhancer elements for eukaryotic transcription factors [34]. Interestingly, some of them were reported to be involved in sex determination in animals, suggesting that the motif has been conserved across kingdoms for sex or sex-related functions. The function of the remaining GCCG and TCCCAC motifs has not been reported. ACAAT motifs were also found in *L. edodes*, whereas GCGGAGT, found in *L. edodes*, was not detected, indicating that cognate DNA subsites of transcription factors were likely genus specific. The conserved domain in the HD proteins was homeodomain, coiled-coil, and NLS. A mating-type protein could interact with a dimerized form and in the nucleus. Similarly, an interaction between HD1 and HD2 homeodomains has been reported for sexual development regulation in basidiomycetes [2,35]. Interestingly, *Pea2* contained only three coiled-coil domains with a value less than 0.021 for coiled-coil possibility. In this case, an HD protein with a low value was found in the *P. tuoliensis* CC00489 HD2 protein. Whether these low values are sufficient for dimerization is unknown; however, the dimerization motif in HD2 is reported to be weak due to the existence of additional forces, such as protein-protein interactions [2].

Many *A* alleles were found in *P. eryngii* [8], and thus, the development of more accurate molecular markers is required. The molecular structure of the *A4* locus will provide a basis for the development of vast and diverse *A*-allele detection markers. The newly developed method might be preferred over the traditional method for the identification of mating type due to its shorter duration, i.e., 1–3 days instead of 15 days. Rapid colony PCR for mycelium was well established for massive selections [29]; therefore, breeders could save time and materials by mating only compatible strains with SCAR markers in the case of inbreeding. Moreover, *P. eryngii* KNR2312 has been used as a parent for many cultivars [30–32] due to a narrow genetic pool; thus, this marker will be more effective for breeding using KNR2312-derived cultivars.

Disclosure statement

No potential conflict of interest was reported by the authors.

Funding

This study was supported by a grant from the Golden seed project [IPET No. 213007-05-3-SBI30], MIFAFF.

References

- [1] Kies U. Life history and developmental processes in the basidiomycete *Coprinus cinereus*. *Microbiol Mol Biol Rev.* 2000;64:316–353.
- [2] Kies U. From two to many: multiple mating types in Basidiomycetes. *Fungal Biol Rev.* 2015;29:126–166.
- [3] Raudaskoski M, Kothe E. Basidiomycete mating type genes and pheromone signaling. *Eukaryot Cell.* 2010;9:847–859.
- [4] Kim K-H, Kang YM, Im CH, et al. Identification and functional analysis of pheromone and receptor genes in the B3 mating locus of *Pleurotus eryngii*. *PLoS One.* 2014;9:e104693.
- [5] Casselton LA, Olesnicky NS. Molecular genetics of mating recognition in basidiomycete fungi. *Microbiol Mol Biol Rev.* 1998;62:55–70.
- [6] Banham AH, Asante-Owusu RN, Gottgens B, et al. An N-terminal dimerization domain permits homeodomain proteins to choose compatible partners and initiate sexual development in the mushroom *Coprinus cinereus*. *Plant Cell.* 1995;7:773–783.
- [7] O’Shea SF, Chaure PT, Halsall JR, et al. A large pheromone and receptor gene complex determines multiple B mating type specificities in *Coprinus cinereus*. *Genetics.* 1998;148:1081–1090.
- [8] Ryu J-S, Kim MK, Ro H-S, et al. Identification of mating type loci and development of SCAR marker genetically linked to the B3 locus in *Pleurotus eryngii*. *J Microbiol Biotechnol.* 2012;22:1177–1184.
- [9] Kothe E. Mating-type genes for basidiomycete strain improvement in mushroom farming. *Appl Microbiol Biotechnol.* 2001;56:602–612.
- [10] Estrada AR, Royse D. Yield, size and bacterial blotch resistance of *Pleurotus eryngii* grown on cottonseed hulls/oak sawdust supplemented with manganese, copper and whole ground soybean. *Bioresour Technol.* 2007;98:1898–1906.
- [11] Ngai PH, Ng T. A hemolysin from the mushroom *Pleurotus eryngii*. *Appl Microbiol Biotechnol.* 2006;72:1185–1191.
- [12] Zervakis G, Venturella G. Mushroom breeding and cultivation enhances ex situ conservation of Mediterranean *Pleurotus taxa*. In: Engels JMM, Ramanantha Rao V, Brown AHD, Jackson MT, editors. *Managing plant genetic diversity*. New York, USA: CABI Publishing; 2002. p. 351–358.
- [13] Zhang R, Hu D, Zhang J, et al. Development and characterization of simple sequence repeat (SSR) markers for the mushroom *Flammulina velutipes*. *J Biosci Bioeng.* 2010;110:273–275.
- [14] Tanaka A, Miyazaki K, Murakami H, et al. Sequence characterized amplified region markers tightly linked to the mating factors of *Lentinula edodes*. *Genome.* 2004;47:156–162.
- [15] Scovel G, Ben-Meir H, Ovadis M, et al. RAPD and RFLP markers tightly linked to the locus controlling carnation (*Dianthus caryophyllus*) flower type. *Theor Appl Genet.* 1998;96:117–122.
- [16] Ryu J, Kim K, Im C, et al., editors. Complete genome sequence of *Pleurotus eryngii* KNR2312 using the next generation sequencing (NGS). 80th Meeting of the Mycological Society of America.

- The Mycological Society of America, New Haven, CT, USA; 2012.
- [17] Michelmore RW, Paran I, Kesseli R. Identification of markers linked to disease-resistance genes by bulked segregant analysis: a rapid method to detect markers in specific genomic regions by using segregating populations. *Proc Natl Acad Sci USA*. 1991;88:9828–9832.
- [18] Marchler-Bauer A, Bryant SH. CD-Search: protein domain annotations on the fly. *Nucleic Acids Res*. 2004;32:W327–W331.
- [19] Lupas A, Van Dyke M, Stock J. Predicting coiled coils from protein sequences. *Science*. 1991;252:1162–1164.
- [20] Ba ANN, Pogoutse A, Provart N, et al. NLStradamus: a simple Hidden Markov Model for nuclear localization signal prediction. *BMC Bioinformatics*. 2009;10:202.
- [21] Im CH, Park Y-H, Hammel KE, et al. Construction of a genetic linkage map and analysis of quantitative trait loci associated with the agronomically important traits of *Pleurotus eryngii*. *Fungal Genet Biol*. 2016;92:50–64.
- [22] Au CH, Wong MC, Bao D, et al. The genetic structure of the A mating-type locus of *Lentinula edodes*. *Gene*. 2014;535:184–190.
- [23] Zhao M, Zhang J, Chen Q, et al. The famous cultivated mushroom Bailinggu is a separate species of the *Pleurotus eryngii* species complex. *Sci Rep*. 2016;6:33066.
- [24] James TY, Liou S-R, Vilgalys R. The genetic structure and diversity of the A and B mating-type genes from the tropical oyster mushroom, *Pleurotus djamor*. *Fungal Genet Biol*. 2004;41:813–825.
- [25] Berbee ML, Taylor JW. Detecting morphological convergence in true fungi, using 18S rRNA gene sequence data. *Biosystems*. 1992;28:117–125.
- [26] Hansen K, LoBuglio KF, Pfister DH. Evolutionary relationships of the cup-fungus genus *Peziza* and *Pezizaceae* inferred from multiple nuclear genes: RPB2, beta-tubulin, and LSU rDNA. *Mol Phylogenet Evol*. 2005;36:1–23.
- [27] Lopandic K, Molnár O, Prillinger H. Application of ITS sequence analysis, RAPD and AFLP fingerprinting in characterising the yeast genus *Fellomyces*. *Microbiol Res*. 2005;160:13–26.
- [28] Weber J, Díez J, Selosse M-A, et al. SCAR markers to detect mycorrhizas of an American *Laccaria bicolor* strain inoculated in European Douglas-fir plantations. *Mycorrhiza*. 2002;12:19–27.
- [29] Izumitsu K, Hatoh K, Sumita T, et al. Rapid and simple preparation of mushroom DNA directly from colonies and fruiting bodies for PCR. *Mycoscience*. 2012;53:396–401.
- [30] Im CH, Kim M-K, Kim K-H, et al. Breeding of king oyster mushroom, *Pleurotus eryngii* with a high yield and earliness of harvest trait and its sensory test. *Kor J Mycol*. 2013;41:91–96.
- [31] Shin P, Park Y, Yoo Y, et al. Characteristics and breeding of a new cultivar *Pleurotus eryngii*, Song-A. *J Mushroom Sci Prod*. 2011;9:59–62.
- [32] Lee Y, Kim Y, Seuk S, et al. Mband characterization of a cultivar “DanBi 5Ho” with a long shelf life. *J Mushrooms*. 2016;14:64–69.
- [33] Burglin T. A comprehensive classification of homeobox genes. In: Duboule D, editor. *Guidebook to the homeobox genes*. Oxford: Oxford University Press; 1994. p. 25–27.
- [34] Narayana N, Weiss MA. Crystallographic analysis of a sex-specific enhancer element: sequence-dependent DNA structure, hydration, and dynamics. *J Mol Biol*. 2009;385:469–490.
- [35] Kües U, Göttgens B, Stratmann R, et al. A chimeric homeodomain protein causes self-compatibility and constitutive sexual development in the mushroom *Coprinus cinereus*. *EMBO J*. 1994;13:4054–4059.