

**COMMON FIXED POINT RESULTS FOR
NON-COMPATIBLE R-WEAKLY COMMUTING
MAPPINGS IN PROBABILISTIC SEMIMETRIC SPACES
USING CONTROL FUNCTIONS**

KRISHNAPADA DAS

ABSTRACT. In common fixed point problems in metric spaces several versions of weak commutativity have been considered. Mappings which are not compatible have also been discussed in common fixed point problems. Here we consider common fixed point problems of non-compatible and R-weakly commuting mappings in probabilistic semimetric spaces with the help of a control function. This work is in line with research in probabilistic fixed point theory using control functions. Further we support our results by examples.

1. Introduction

The problem of finding common fixed points of more than one mappings have been considered in several contemporary works. In this context the commuting condition on a pair of mappings has been relaxed mainly in two directions. In one direction the notion of compatibility of various types have been introduced and common fixed point results of several types of compatible mappings have been established. This line of research was initiated by Jungck [11]. It has also been noted that fixed point problems of non-compatible mappings are also important and have been considered in a number of recent works [20, 23].

Received October 3, 2018. Revised June 28, 2019. Accepted July 27, 2019.

2010 Mathematics Subject Classification: 47H10, 54H25, 54E70.

Key words and phrases: Menger space, compatible maps, (ϕ, R) -weakly commuting maps, fixed point.

© The Kangwon-Kyungki Mathematical Society, 2019.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

In another direction weaker versions of commutativity have been considered in a large number of works. One such concept is R-weakly commutativity, which was introduced by Pant [19]. This is an extension of weakly commuting mappings [18, 28]. Some of other references dealing with R-weakly commuting mappings are [20–22] and [23]. Recently commutativity conditions have also been used to find coupled coincidence point in [5–7].

In metric fixed point theory a new direction was opened by Khan et al in [13]. They introduced a new contraction mapping principle and proved a fixed point result with the help of a control function which they called altering distance function. Altering distance function has been used in a number of papers in metric fixed point theory. Some of these results are noted in [17, 24] and [25].

K. Menger [15] introduced the notion of probabilistic metric space in 1942 and since then the theory of probabilistic metric spaces has emerged as an active area of research. Menger utilized distribution functions instead of non-negative real numbers as the values of the metric. The notion of probabilistic metric space corresponds to the situation when we do not know exactly the distance between two points, we only know the probabilities of possible values of the distances. Probabilistic generalizations of the metric spaces appears to be interesting in the investigation of physical quantities and useful in modelling some physical phenomena.

First fixed point result in probabilistic metric spaces appeared in literature in the work of Sehgal and Bharucha-Reid [27]. After that result fixed point and common fixed point properties for mappings defined on probabilistic spaces has been studied by many authors. Hadzic and Pap in [12] has given a comprehensive survey of this line of research.

Recently in [2] Choudhury and Das have extended the idea of altering distance function to probabilistic metric spaces and have established a generalization of Sehgal's contraction, where a probabilistic contraction mapping principle has been established through the application of a control function. The introduction of control function in probabilistic spaces opens new possibilities of establishing new fixed point results. Some recent works in probabilistic spaces where this control function have been utilized are noted in [1, 3, 4, 8–10, 14, 16, 29] and [30]. Here we make another use of this control function to fixed point problems in probabilistic spaces. Precisely in this work we use the control function to establish common fixed point results in probabilistic semimetric spaces

for mappings which are non-compatible and satisfy R-weakly commuting condition. Our results are supported by examples.

2. Definitions and Mathematical Preliminaries

In this section we give some definitions and results which are needed for our results.

DEFINITION 2.1. T-NORM [12, 26]

A t-norm is a function $\Delta : [0, 1] \times [0, 1] \rightarrow [0, 1]$ which satisfies the following conditions

- (i) $\Delta(1, a) = a,$
- (ii) $\Delta(a, b) = \Delta(b, a),$
- (iii) $\Delta(c, d) \geq \Delta(a, b)$ whenever $c \geq a$ and $d \geq b,$
- (iv) $\Delta(\Delta(a, b), c) = \Delta(a, \Delta(b, c)).$

DEFINITION 2.2. [12, 26] A mapping $F : R \rightarrow R^+$ is called a distribution function if it is non-decreasing and left continuous with $\inf_{t \in R} F(t) = 0$ and $\sup_{t \in R} F(t) = 1,$ where R is the set of real numbers and R^+ denotes the set of non-negative real numbers.

DEFINITION 2.3. PROBABILISTIC SEMIMETRIC SPACE [12, 26]

Let M be a non empty set, F is a function defined on $M \times M$ to the set of distribution functions. (M, F) is said to be a probabilistic semimetric spaces if the following are satisfied:

- (i) $F_{x,y}(0) = 0$ for all $x, y \in M,$
- (ii) $F_{x,y}(s) = 1$ for all $s > 0$ and $x, y \in M$ if and only if $x = y,$
- (iii) $F_{x,y}(s) = F_{y,x}(s)$ for all $x, y \in M, s > 0.$

DEFINITION 2.4. Menger Space [12, 26]

A Menger space is a triplet (M, F, Δ) where M is a non empty set, F is a function defined on $M \times M$ to the set of distribution functions and Δ is a t-norm, such that the following are satisfied:

- (i) $F_{x,y}(0) = 0$ for all $x, y \in M,$
- (ii) $F_{x,y}(s) = 1$ for all $s > 0$ and $x, y \in M$ if and only if $x = y,$
- (iii) $F_{x,y}(s) = F_{y,x}(s)$ for all $x, y \in M, s > 0$ and
- (iv) $F_{x,y}(u + v) \geq \Delta(F_{x,z}(u), F_{z,y}(v))$ for all $u, v \geq 0$ and $x, y, z \in M.$

A sequence $\{x_n\} \subset M$ converges to some point $x \in M$ if for given $\epsilon > 0, \lambda > 0$ we can find a positive integer $N_{\epsilon, \lambda}$ such that for all $n > N_{\epsilon, \lambda}$,

$$F_{x_n, x}(\epsilon) > 1 - \lambda.$$

DEFINITION 2.5. ALTERING DISTANCE FUNCTION [13]

A function $h : [0, \infty) \rightarrow [0, \infty)$ is an altering distance function if

- (i) h is monotone increasing and continuous and
- (ii) $h(t) = 0$ if and only if $t = 0$.

Khan et al. proved the following generalization of Banach contraction mapping principle.

THEOREM 2.6. [13] Let (X, d) be a complete metric space, h be an altering distance function and let $f : X \rightarrow X$ be a self mapping which satisfies the following inequality

$$h(d(fx, fy)) \leq ch(d(x, y))$$

for all $x, y \in X$ and for some $0 < c < 1$. Then f has a unique fixed point.

In fact Khan et al. proved a more general theorem (Theorem-2 in [13]) of which the above result is a corollary.

DEFINITION 2.7. Φ -FUNCTION [2]

A function $\phi : [0, \infty) \rightarrow [0, \infty)$ is said to be a Φ -function if it satisfies the following conditions:

- (i) $\phi(t) = 0$ if and only if $t = 0$,
- (ii) $\phi(t)$ is strictly increasing and $\phi(t) \rightarrow \infty$ as $t \rightarrow \infty$,
- (iii) ϕ is left continuous in $(0, \infty)$ and
- (iv) ϕ is continuous at 0.

An altering distance function with the additional property that $h(t) \rightarrow \infty$ as $t \rightarrow \infty$ generates a Φ -function in the following way.

$$\phi(t) = \begin{cases} \sup\{s : h(s) < t\}, & \text{if } t > 0, \\ 0, & \text{if } t = 0. \end{cases}$$

It can be easily seen that ϕ is a Φ -function.

The following result has been established in [2].

THEOREM 2.8. [2] Let (M, F, Δ) be a complete Menger space with $\Delta(a, b) = \min\{a, b\}$ and $f : M \rightarrow M$ be a self mapping such that the following inequality is satisfied.

$$F_{fx, fy}(\phi(t)) \geq F_{x,y}(\phi(t/c))$$

where ϕ is a Φ -function, $0 < c < 1$, $t > 0$ and $x, y \in M$. Then f has a unique fixed point.

DEFINITION 2.9. Two self-mappings f and g of a probabilistic semimetric space (M, F) are called compatible if $\lim_{n \rightarrow \infty} F_{fgx_n, gfx_n}(t) = 1$ for all $t > 0$ whenever $\{x_n\}$ is a sequence such that $\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = z$ for some z in M .

DEFINITION 2.10. Two self-mappings f and g of a probabilistic semimetric space (M, F) are called (ϕ, R) -weakly commuting if there exists some real number $R > 0$ such that

$$F_{fgx, gfx}(\phi(t)) \geq F_{fx, gx}(\phi(\frac{t}{R})) \text{ for all } t > 0 \text{ and for all } z \in M.$$

Now f and g will be called pointwise (ϕ, R) -weakly commuting if given x in M , there exists $R > 0$ such that

$$F_{fgx, gfx}(\phi(t)) \geq F_{fx, gx}(\phi(\frac{t}{R})) \text{ for all } t > 0.$$

DEFINITION 2.11. Ψ -FUNCTION

A function $\psi : [0, 1] \times [0, 1] \rightarrow [0, 1]$ is said to be a Ψ -function if

- (i) ψ is monotone increasing and continuous,
- (ii) $\psi(x, x) > x$ for all $0 < x < 1$,
- (iii) $\psi(0, 0) = 0$,
- (iv) $\psi(x, x) = 1$ if and only if $x = 1$.

An example of Ψ -function.

$$1. \quad \psi(x, y) = \frac{p\sqrt{x}+q\sqrt{y}}{p+q}, \text{ } p \text{ and } q \text{ are positive real numbers.}$$

3. Main Results

THEOREM 3.1. Let (M, F) be a probabilistic semimetric space and $f, g : M \rightarrow M$ be non-compatible pointwise (ϕ, R) -weakly commuting mappings such that

$$[i] \quad fM \subset gM \tag{2.1}$$

$$[ii] F_{fx, fy}(\phi(t)) > \min\{F_{gx, gy}(\phi(t)), \psi(F_{fx, gx}(\phi(\frac{t_1}{a_1})), F_{fy, gy}(\phi(\frac{t_2}{b_1}))), \\ \psi(F_{fy, gx}(\phi(\frac{t_3}{a_2})), F_{fx, gy}(\phi(\frac{t_4}{b_2})))\} \quad (2.2)$$

where $x, y \in M$ with $x \neq y$, $a_1, a_2, b_1, b_2 > 0$, with $0 < a_1 + b_1 < 1, 0 < a_2 + b_2 < 1$, $t, t_1, t_2, t_3, t_4 > 0$ with $t_1 + t_2 = t = t_3 + t_4$, ϕ is a Φ -function and ψ is a Ψ -function. If the range of f or g is a complete subspace of M then f and g have a unique common fixed point.

Proof. The mappings f and g are non-compatible maps, hence there exists a sequence $\{x_n\}$ in M such that $\lim_{n \rightarrow \infty} fx_n = \lim_{n \rightarrow \infty} gx_n = z$ for some $z \in M$ but $\lim_{n \rightarrow \infty} F_{fgx_n, gfx_n}(\phi(t)) \neq 1$ for some $t > 0$ or the limit does not exist.

Let us suppose that the range of g is a complete subspace of M . Then there exists some point $u \in M$ such that $z = gu$ where $z = \lim_{n \rightarrow \infty} gx_n$. If possible let $fu \neq gu$.

$$\text{Also let us take } t_1 = \frac{a_1 t}{a_1 + b_1}, t_2 = \frac{b_1 t}{a_1 + b_1}, t_3 = \frac{a_2 t}{a_2 + b_2}, t_4 = \frac{a_2 t}{a_2 + b_2}, \quad (2.3)$$

$$\text{and } c_1 = a_1 + b_1, c_2 = a_2 + b_2 \text{ and } \max\{c_1, c_2\} = c. \quad (2.4)$$

Now we have

$$F_{fx_n, fu}(\phi(t)) > \min\{F_{gx_n, gu}(\phi(t)), \psi(F_{fx_n, gx_n}(\phi(\frac{t_1}{a_1})), F_{fu, gu}(\phi(\frac{t_2}{b_1}))), \\ \psi(F_{fu, gx_n}(\phi(\frac{t_3}{a_2})), F_{fx_n, gu}(\phi(\frac{t_4}{b_2})))\} \\ \geq \min\{F_{gx_n, gu}(\phi(t)), \psi(F_{fx_n, gx_n}(\phi(\frac{t}{c})), F_{fu, gu}(\phi(\frac{t}{c}))), \\ \psi(F_{fu, gx_n}(\phi(\frac{t}{c})), F_{fx_n, gu}(\phi(\frac{t}{c})))\} \\ \geq \min\{F_{gx_n, gu}(\phi(t)), \psi(F_{fx_n, gx_n}(\phi(t)), F_{fu, gu}(\phi(t))), \\ \psi(F_{fu, gx_n}(\phi(t)), F_{fx_n, gu}(\phi(t)))\} \quad (2.5)$$

Letting $n \rightarrow \infty$ we have from (2.5),

$$F_{gu, fu}(\phi(t)) \geq \min\{F_{gu, gu}(\phi(t)), \psi(F_{gu, gu}(\phi(t)), F_{fu, gu}(\phi(t))), \\ \psi(F_{fu, gu}(\phi(t)), F_{gu, gu}(\phi(t)))\} \\ \geq \min\{1, \psi(1, F_{fu, gu}(\phi(t))), \psi(F_{fu, gu}(t), 1)\}. \quad (2.6)$$

We claim $F_{fu, gu}(\phi(t)) = 1$ for all $t > 0$. If not then for some $t > 0$, $0 < F_{fu, gu}(\phi(t)) < 1$ and by the property of ψ we have from (2.6)

$$F_{gu, fu}(\phi(t)) \geq \min\{\psi(1, F_{fu, gu}(t)), \psi(F_{fu, gu}(\phi(t)), 1)\} \\ \geq \min\{\psi(F_{fu, gu}(\phi(t)), F_{fu, gu}(\phi(t))), \\ \psi(F_{fu, gu}(\phi(t)), F_{fu, gu}(\phi(t)))\}$$

$$\begin{aligned}
 &= \psi(F_{fu,gu}(\phi(t)), F_{fu,gu}(\phi(t))) \\
 &> F_{fu,gu}(\phi(t)) , \quad \text{which is a contradiction.}
 \end{aligned}$$

Therefore for all $t > 0$ we have , $F_{gu,fu}(\phi(t)) = 1$. (2.7)
 From the property of ϕ it follows that given $s > 0$ we can find $t > 0$ such that $s > \phi(t) > 0$.

Therefore for all $s > 0$ we have , $F_{gu,fu}(s) = 1$ that is, $gu = fu$.

Now f and g are (ϕ, R) -weakly commuting maps, therefore there exists $R_1 > 0$ such that for all $t > 0$

$$F_{fgu,gfu}(\phi(t)) \geq F_{fu,gu}(\phi(\frac{t}{R_1})) = F_{fu,fu}(\phi(\frac{t}{R_1})) = 1.$$

Therefore we can say as in above $fgu = gfu$ and hence

$$ffu = fgu = gfu = ggu.$$

Now we claim that $fu = ffu$. If $fu \neq ffu$ then we have,

$$\begin{aligned}
 F_{fu,ffu}(\phi(t)) &> \min\{F_{gu,gfu}(\phi(t)), \psi(F_{fu,gu}(\phi(\frac{t_1}{a_1})), F_{ffu,gfu}(\phi(\frac{t_2}{b_1}))), \\
 &\quad \psi(F_{ffu,gu}(\phi(\frac{t_3}{a_2})), F_{fu,gfu}(\phi(\frac{t_4}{b_2})))\} \\
 &\geq \min\{F_{fu,ffu}(\phi(t)), \psi(1, 1), \\
 &\quad \psi(F_{ffu,fu}(\phi(\frac{t_3}{a_2})), F_{fu,ffu}(\phi(\frac{t_4}{b_2})))\}
 \end{aligned}$$

By taking $t_3 = \frac{a_2t}{a_2+b_2}$, $t_4 = \frac{a_2t}{a_2+b_2}$ and $c_2 = a_2 + b_2$ we have from above,

$$\begin{aligned}
 F_{fu,ffu}(\phi(t)) &> \min\{F_{fu,ffu}(\phi(t)), 1, \psi(F_{ffu,fu}(\phi(\frac{t}{c_2})), F_{fu,ffu}(\phi(\frac{t}{c_2})))\} \\
 &\geq \min\{F_{fu,ffu}(\phi(t)), 1, F_{ffu,fu}(\phi(\frac{t}{c_2}))\} \\
 &\geq \min\{F_{fu,ffu}(\phi(t)), 1, F_{ffu,fu}(\phi(t))\} \\
 &= \min\{F_{fu,ffu}(\phi(t)), 1\}
 \end{aligned}$$

If $\min\{F_{fu,ffu}(\phi(t)), 1\} = F_{fu,ffu}(\phi(t))$, then we arrived at contradiction. On the other hand if $\min\{F_{fu,ffu}(\phi(t)), 1\} = 1$, then we have $F_{fu,ffu}(\phi(t)) > 1$, which is impossible.

Therefore $fu = ffu = gfu$. Hence fu is a common fixed point of f and g .

For uniqueness let fu and fv be the two common fixed points of f and g . If $fu \neq fv$ then we have,

$$F_{fu,fv}(\phi(t)) > \min\{F_{gu,gv}(\phi(t)), \psi(F_{fu,gu}(\phi(\frac{t_1}{a_1})), F_{fv,gv}(\phi(\frac{t_2}{b_1}))),$$

$$\begin{aligned}
& \psi(F_{fv,gu}(\phi(\frac{t_3}{a_2})), F_{fu,gv}(\phi(\frac{t_4}{b_2}))) \\
= & \min\{F_{gu,gv}(\phi(t)), \psi(1, 1), \\
& \psi(F_{fv,fu}(\phi(\frac{t_3}{a_2})), F_{fu,fv}(\phi(\frac{t_4}{b_2})))\}
\end{aligned}$$

Now taking t_1, t_2, t_3, t_4 as in (2.3) and c_1, c_2, c as in (2.4) we have ,
 $F_{fu,fv}(\phi(t)) > \min\{F_{fu,fv}(\phi(t)), \psi(1, 1), \psi(F_{fv,fu}(\phi(\frac{t}{c})), F_{fu,fv}(\phi(\frac{t}{c})))\}$
 $\geq \min\{F_{fu,fv}(\phi(t)), 1, F_{fv,fu}(\phi(\frac{t}{c}))\}$
 $\geq \min\{F_{fu,fv}(\phi(t)), 1, F_{fv,fu}(\phi(t))\}$
 $= \min\{F_{fu,fv}(\phi(t)), 1\}$

If $\min\{F_{fu,fv}(\phi(t)), 1\} = F_{fu,fv}(\phi(t))$ then we have $F_{fu,fv}(\phi(t)) > F_{fu,fv}(\phi(t))$, which is a contradiction. Again if $\min\{F_{fu,fv}(\phi(t)), 1\} = 1$ then we have $F_{fu,fv}(\phi(t)) > 1$ which is impossible.

Therefore $fu = fv$. This completes the proof of the theorem. \square

Taking $a_1 = b_1 = a_2 = b_2 = a$ in above theorem we get the following corollary.

COROLLARY 3.2. *Let (M, F) be a probabilistic semimetric space and $f, g : M \rightarrow M$ be non-compatible pointwise (ϕ, R) -weakly commuting mappings such that*

$$[i] \quad fM \subset gM \tag{2.8}$$

$$\begin{aligned}
[ii] \quad F_{fx,fy}(\phi(t)) > \min\{F_{gx,gy}(\phi(t)), \psi(F_{fx,gx}(\phi(\frac{t_1}{a})), F_{fy,gy}(\phi(\frac{t_2}{a}))), \\
\psi(F_{fy,gx}(\phi(\frac{t_3}{a})), F_{fx,gy}(\phi(\frac{t_4}{a})))\}
\end{aligned} \tag{2.9}$$

where $x, y \in M$ with $x \neq y$, $0 < a < \frac{1}{2}$, $t, t_1, t_2, t_3, t_4 > 0$ with $t_1 + t_2 = t$ and $t_3 + t_4 = t$, ϕ is a Φ -function and ψ is a Ψ -function. If the range of f or g is a complete subspace of M then f and g have a unique common fixed point.

THEOREM 3.3. *Let (M, F) be a probabilistic semimetric space, (A, S) and (B, T) be pairwise (ϕ, R) -weakly commuting self-mappings on (M, F, Δ) such that*

$$[i] \quad AM \subset TM \text{ and } BM \subset SM \tag{2.10}$$

$$[ii] F_{Ax,By}(\phi(t)) > \min\{F_{Sx,Ty}(\phi(t)), \psi(F_{Ax,Sx}(\phi(\frac{t_1}{a_1})), F_{By,Ty}(\phi(\frac{t_2}{b_1}))), \\ \psi(F_{Ax,Ty}(\phi(\frac{t_3}{a_2})), F_{By,Sx}(\phi(\frac{t_4}{b_2})))\} \quad (2.11)$$

where $x, y \in M$ with $x \neq y$, $a_1, a_2, b_1, b_2 > 0$, with $0 < c_1 = a_1 + b_1 < 1$, $0 < c_2 = a_2 + b_2 < 1$, $t, t_1, t_2, t_3, t_4 > 0$ with $t_1 + t_2 = t = t_3 + t_4$, ϕ is a Φ -function and ψ is a Ψ -function. Let (A, S) or (B, T) be a non-compatible pair of mappings. If the range of one of the mappings is a complete subspace of M then A, B, S and T have a unique common fixed point.

Proof. Let B and T be non-compatible pair of mappings. Then there exists a sequence $\{x_n\}$ in M such that $\lim_{n \rightarrow \infty} Bx_n = \lim_{n \rightarrow \infty} Tx_n = z$ for some $z \in M$ but $\lim_{n \rightarrow \infty} F_{BTx_n, TBx_n} \phi(t) \neq 1$ for some $t > 0$ or the limit does not exists. Since $BM \subset SM$, for each x_n there exists y_n in M such that $Bx_n = Sy_n$. Thus $Bx_n \rightarrow z$, $Tx_n \rightarrow z$ and $Sy_n \rightarrow z$ as $n \rightarrow \infty$.

We claim that $Ay_n \rightarrow z$ as $n \rightarrow \infty$,
that is, $\lim_{n \rightarrow \infty} F_{Ay_n, z} \phi(t) \rightarrow 1$ for all $t > 0$,

that is, there exists a positive integer N_1 such that $\lim_{n \rightarrow \infty} F_{Ay_n, z} \phi(t) > 1 - \lambda$ for all $t > 0$ and for all $n \geq N_1$.

Otherwise, we have for some $t' > 0$ a subsequence $\{Ay_m\}$ of $\{Ay_n\}$ and a positive number r , $0 < r \leq 1 - \lambda$ such that $\lim_{m \rightarrow \infty} F_{Ay_m, z} \phi(t') = r$.

$$F_{Ay_m, Bx_m}(\phi(t')) \\ > \min\{F_{Sy_m, Tx_m}(\phi(t')), \psi(F_{Ay_m, Sy_m}(\phi(\frac{t'_1}{a_1})), F_{Bx_m, Tx_m}(\phi(\frac{t'_2}{b_1}))), \\ \psi(F_{Ay_m, Tx_m}(\phi(\frac{t'_3}{a_2})), F_{Bx_m, Sy_m}(\phi(\frac{t'_4}{b_2})))\}. \quad (2.12)$$

Taking $t'_1 = \frac{a_1 t'}{a_1 + b_1}$, $t'_2 = \frac{b_1 t'}{a_1 + b_1}$, $t'_3 = \frac{a_2 t'}{a_2 + b_2}$, $t'_4 = \frac{a_2 t'}{a_2 + b_2}$ and c_1, c_2, c as in (2.4) we have from (2.12),

$$F_{Ay_m, Bx_m}(\phi(t')) \\ > \min\{F_{Sy_m, Tx_m}(\phi(t')), \psi(F_{Ay_m, Sy_m}(\phi(\frac{t'}{c})), F_{Bx_m, Tx_m}(\phi(\frac{t'}{c}))), \\ \psi(F_{Ay_m, Tx_m}(\phi(\frac{t'}{c})), F_{Bx_m, Sy_m}(\phi(\frac{t'}{c})))\}. \\ \geq \min\{F_{Sy_m, Tx_m}(\phi(t')), \psi(F_{Ay_m, Sy_m}(\phi(t')), F_{Bx_m, Tx_m}(\phi(t'))), \\ \psi(F_{Ay_m, Tx_m}(\phi(t')), F_{Bx_m, Sy_m}(\phi(t')))\}.$$

Letting $n \rightarrow \infty$ we get,

$$r \geq \min\{F_{z,z}(\phi(t')), \psi(r, F_{z,z}(\phi(t'))), \psi(r, F_{z,z}(\phi(t')))\}. \\ \geq \min\{1, \psi(r, 1), \psi(r, 1)\}$$

$$\begin{aligned}
&= \psi(r, 1) \\
&\geq \psi(r, r) \\
&> r, \text{ which is a contradiction.}
\end{aligned}$$

Therefore, $Ay_n \rightarrow z$ as $n \rightarrow \infty$. (2.13)

Suppose that SM is a complete subspace of M . Then, since $Sy_n \rightarrow z$, there exists a point u in M such that $z = Su$. If $Au \neq Su$ then we have,

$$\begin{aligned}
&F_{Au, Bx_n}(\phi(t)) \\
&> \min\{F_{Su, Tx_n}(\phi(t)), \psi(F_{Au, Su}(\phi(\frac{t_1}{a_1})), F_{Bx_n, Tx_n}(\phi(\frac{t_2}{b_1}))), \\
&\quad \psi(F_{Au, Tx_n}(\phi(\frac{t_3}{a_2})), F_{Bx_n, Su}(\phi(\frac{t_4}{b_2})))\}. \tag{2.14}
\end{aligned}$$

Now as (2.3) and (2.4) the same choice of t_1, t_2, t_3, t_4 and c_1, c_2, c gives us,

$$\begin{aligned}
&F_{Au, Bx_n}(\phi(t)) \\
&> \min\{F_{Su, Tx_n}(\phi(t)), \psi(F_{Au, Su}(\phi(\frac{t}{c})), F_{Bx_n, Tx_n}(\phi(\frac{t}{c}))), \\
&\quad \psi(F_{Au, Tx_n}(\phi(\frac{t}{c})), F_{Bx_n, Su}(\phi(\frac{t}{c})))\}. \\
&\geq \min\{F_{Su, Tx_n}(\phi(t)), \psi(F_{Au, Su}(\phi(t)), F_{Bx_n, Tx_n}(\phi(t))), \\
&\quad \psi(F_{Au, Tx_n}(\phi(t)), F_{Bx_n, Su}(\phi(t)))\}.
\end{aligned}$$

Therefore taking limit as $n \rightarrow \infty$ we get,

$$\begin{aligned}
F_{Au, Su}(\phi(t)) &\geq \min\{F_{Su, Su}(\phi(t)), \psi(F_{Au, Su}(\phi(t)), F_{Su, Su}(\phi(t))), \\
&\quad \psi(F_{Au, Su}(\phi(t)), F_{z, z}(\phi(t)))\}. \tag{2.15}
\end{aligned}$$

If for all $t > 0$, $F_{Au, Su}(\phi(t)) = 1$ then as above we have $Au = Su$. If not then there exists some $t > 0$ such that $0 < F_{Au, Su}(\phi(t)) < 1$.

Therefore,

$$\begin{aligned}
F_{Au, Su}(\phi(t)) &\geq \min\{1, \psi(F_{Au, Su}(\phi(t)), 1), \psi(F_{Au, Su}(\phi(t)), 1)\} \\
&= \psi(F_{Au, Su}(\phi(t)), 1) \\
&\geq \psi(F_{Au, Su}(\phi(t)), F_{Au, Su}(\phi(t))) \\
&> F_{Au, Su}(\phi(t))
\end{aligned}$$

which is a contradiction.

Therefore $Au = Su$.

Now pointwise R -weak commutativity of A and S implies that there exist a $R_1 > 0$ such that for all $t > 0$,

$$F_{ASu,SAu}(\phi(t)) \geq F_{Au,Su}(\phi(\frac{t}{R_1})) = 1,$$

that is, $ASu = SAu$ and that of $AAu = ASu = SAu = SSu$. Since $AM \subset TM$, there exists a w in M such that $Au = Tw$. We next show that $Tw = Bw$. If not then we have,

$$\begin{aligned} & F_{Au,Bw}(\phi(t)) \\ & > \min\{F_{Su,Tw}(\phi(t)), \psi(F_{Au,Su}(\phi(\frac{t_1}{a_1})), F_{Bw,Tw}(\phi(\frac{t_2}{b_1}))), \\ & \quad \psi(F_{Au,Tw}(\phi(\frac{t_3}{a_2})), F_{Bw,Su}(\phi(\frac{t_4}{b_2})))\}. \end{aligned} \tag{2.16}$$

By the similar choice of t_1, t_2, t_3, t_4 and c_1, c_2, c as in (2.3) and (2.4) respectively gives us,

$$\begin{aligned} & F_{Au,Bw}(\phi(t)) \\ & > \min\{F_{Su,Tw}(\phi(t)), \psi(F_{Au,Su}(\phi(\frac{t}{c})), F_{Bw,Tw}(\phi(\frac{t}{c}))), \\ & \quad \psi(F_{Au,Tw}(\phi(\frac{t}{c})), F_{Bw,Su}(\phi(\frac{t}{c})))\} \\ & = \min\{F_{Au,Au}(\phi(t)), \psi(F_{Au,Au}(\phi(\frac{t}{c})), F_{Bw,Au}(\phi(\frac{t}{c}))), \\ & \quad \psi(F_{Au,Au}(\phi(\frac{t}{c})), F_{Bw,Au}(\phi(\frac{t}{c})))\} \\ & = \min\{1, \psi(1, F_{Bw,Au}(\phi(\frac{t}{c}))), \psi(1, F_{Bw,Au}(\phi(\frac{t}{c})))\} \\ & \geq \min\{1, \psi(F_{Bw,Au}(\phi(\frac{t}{c}))), F_{Bw,Au}(\phi(\frac{t}{c}))), \\ & \quad \psi(F_{Bw,Au}(\phi(\frac{t}{c})), F_{Bw,Au}(\phi(\frac{t}{c})))\} \\ & \geq \min\{1, F_{Bw,Au}(\phi(\frac{t}{c})), F_{Bw,Au}(\phi(\frac{t}{c}))\} \\ & \geq F_{Bw,Au}(\phi(\frac{t}{c})) \\ & \geq F_{Bw,Au}(\phi(t)), \text{ which is a contradiction.} \end{aligned}$$

Hence $Au = Bw = Tw = Su$. Pointwise R -weakly commutativity of B and T implies that $BTw = TBw$ and $BBw = BTw = TBw = TTW$. Now if $Au \neq AAu$ then applying the same procedure we have,

$$\begin{aligned} & F_{Au,AAu}(\phi(t)) \\ & = F_{AAu,Au}(\phi(t)) = F_{AAu,Bw}(\phi(t)) \\ & > \min\{F_{SAu,Tw}(\phi(t)), \psi(F_{AAu,SAu}(\phi(\frac{t}{c})), F_{Bw,Tw}(\phi(\frac{t}{c}))), \\ & \quad \psi(F_{AAu,Tw}(\phi(\frac{t}{c})), F_{Bw,SAu}(\phi(\frac{t}{c})))\} \\ & = \min\{F_{AAu,Au}(\phi(t)), \psi(F_{AAu,AAu}(\phi(\frac{t}{c})), F_{Au,Au}(\phi(\frac{t}{c}))), \\ & \quad \psi(F_{AAu,Au}(\phi(\frac{t}{c})), F_{Au,AAu}(\phi(\frac{t}{c})))\} \\ & \geq \min\{F_{AAu,Au}(\phi(t)), \psi(1, 1), F_{AAu,Au}(\phi(\frac{t}{c}))\} \\ & \geq \min\{F_{AAu,Au}(\phi(t)), 1, F_{AAu,Au}(\phi(\frac{t}{c}))\} \end{aligned}$$

$$\begin{aligned} &\geq \min\{F_{AAu,Au}(\phi(t)), F_{AAu,Au}(\phi(t))\} \\ &= F_{AAu,Au}(\phi(t)), \text{ which is a contradiction.} \end{aligned}$$

Therefore, $Au = AAu = SAu$ and Au is a common fixed point of A and S . Similarly $Au = Bu$ is fixed point of B and T . The proof is similar when TM is assumed to be a complete subspace of M . When AM or BM is a complete subspace of M the proof will be similar to the case when TM or SM is complete respectively as $AM \subset TM$ and $BM \subset SM$.

For uniqueness if possible let u and v be two distinct common fixed points of A, B, S and T that is, $Au = Bu = Su = Tu = u$ and $Av = Bv = Sv = Tv = v$. Since u and v be two distinct points we can find a $t > 0$ such that, $0 < F_{u,v}(\phi(t)) < 1$ and we have

$$\begin{aligned} &F_{u,v}(\phi(t)) \\ &= F_{Au,Bv}(\phi(t)) \\ &> \min\{F_{Su,Tv}(\phi(t)), \psi(F_{Au,Su}(\phi(\frac{t_1}{a_1})), F_{Bv,Tv}(\phi(\frac{t_2}{b_1}))), \psi(F_{Au,Tv}(\phi(\frac{t_3}{a_2})), F_{Bv,Su}(\phi(\frac{t_4}{b_2})))\} \\ &= \min\{F_{u,v}(\phi(t)), \psi(F_{u,u}(\phi(\frac{t_1}{a_1})), F_{v,v}(\phi(\frac{t_2}{b_1}))), \psi(F_{u,v}(\phi(\frac{t_3}{a_2})), F_{v,u}(\phi(\frac{t_4}{b_2})))\} \\ &= \min\{F_{u,v}(\phi(t)), \psi(1, 1), \psi(F_{u,v}(\phi(\frac{t_3}{a_2})), F_{v,u}(\phi(\frac{t_4}{b_2})))\} \end{aligned} \quad (2.17)$$

By the same choice of t_3, t_4 and c as in (2.3) and (2.4) respectively we have from (2.17),

$$\begin{aligned} F_{u,v}(\phi(t)) &> \min\{F_{u,v}(\phi(t)), \psi(1, 1), \psi(F_{u,v}(\phi(\frac{t}{c})), F_{v,u}(\phi(\frac{t}{c})))\} \\ &\geq \min\{F_{u,v}(\phi(t)), 1, F_{u,v}(\phi(\frac{t}{c}))\} \\ &\geq \min\{F_{u,v}(\phi(t)), F_{u,v}(\phi(t))\} \\ &= F_{u,v}(\phi(t)), \text{ which is a contradiction.} \end{aligned}$$

Hence the uniqueness of the common fixed point is proved.

This completes the proof of the theorem. \square

EXAMPLE 3.4. Let (M, F) be a probabilistic semimetric space, where $M = [2, 20]$ and $F_{x,y}(t) = e^{-\frac{(x-y)^2}{t}}$. Define $f, g : M \rightarrow M$ as follows:

$$fx = \begin{cases} 2, & \text{if } x = 2, \\ 6, & \text{if } 2 < x \leq 5, \\ 2, & \text{if } x > 5, \end{cases}$$

$$gx = \begin{cases} 2, & \text{if } x = 2, \\ 12, & \text{if } 2 < x \leq 5, \\ \frac{x+1}{3}, & \text{if } x > 5. \end{cases}$$

If we take $\phi(t) = \begin{cases} t^2, & \text{if } 0 \leq x \leq 1, \\ \frac{t^2+1}{2}, & \text{if } x > 1. \end{cases}$ and $\psi(x, y) = \frac{\sqrt{x}+\sqrt{y}}{2}$ then f and g satisfies all the conditions of Theorem 3.1 have a unique common fixed point at $x = 2$.

EXAMPLE 3.5. Let (M, F) be a probabilistic semimetric space, where $M = [2, 20]$ and $F_{x,y}(t) = e^{-\frac{(x-y)^2}{t}}$. Define $A, B, S, T : M \rightarrow M$ as follows:

$$Ax = 2 \quad \text{if } 2 \leq x \leq 20,$$

$$Bx = \begin{cases} 2, & \text{if } x = 2, \\ 8, & \text{if } 2 < x \leq 5, \\ 2, & \text{if } x > 5, \end{cases}$$

$$Sx = \begin{cases} x, & \text{if } 2 \leq x \leq 8, \\ 8, & \text{if } x > 8, \end{cases}$$

$$Tx = \begin{cases} 2, & \text{if } x = 2, \\ 12 + x, & \text{if } 2 < x \leq 5, \\ x - 3, & \text{if } x \geq 5. \end{cases}$$

If we take $\phi(t) = \begin{cases} t^2, & \text{if } 0 \leq x \leq 1, \\ \frac{t^2+1}{2}, & \text{if } x > 1. \end{cases}$ and $\psi(x, y) = \frac{\sqrt{x}+\sqrt{y}}{2}$ then A, B, S and T satisfies all the conditions of Theorem 3.3 have a unique common fixed point at $x = 2$.

4. Conclusion

The structure of the probabilistic metric space allows us to develop fixed point theory in several ways not always available in ordinary metric space. In this paper we have proved some fixed point results for non-compatible R-weakly commuting mappings. Our paper is also an

instance of the use of control functions in fixed point theory in probabilistic metric spaces. This control function appears to be helpful in exploring the geometric aspects of Menger spaces which is also relevant to the study of geometry at the quantum level.

References

- [1] C.T. Aage, B.S. Choudhury and K. Das, *Some fixed point results in fuzzy metric spaces using a control function*, Surveys in Mathematics and its Applications **12** (2017), 23–34.
- [2] B.S. Choudhury and K. Das, *A new contraction principle in Menger spaces*, Acta Mathematica Sinica **24** (8) (2008), 1379–1386.
- [3] B.S. Choudhury, P.N. Dutta and K. Das, *A fixed points result in Menger space using a real function*, Acta. Math. Hungar. **122** (2008), 203–216.
- [4] B.S. Choudhury and K. Das, *A coincidence point result in Menger spaces using a control function*, Chaos, Solitons and Fractals **42** (2009), 3058–3063.
- [5] B.S. Choudhury, K.P. Das and P. Das, *Coupled coincidence point results for compatible mappings in partially ordered fuzzy metric spaces*, Fuzzy Sets and Systems **222** (2013), 84–97.
- [6] B.S. Choudhury, K.P. Das and P. Bhattacharya, *A Common Fixed Point Theorem for Weakly Compatible Mappings in Complete Fuzzy Metric Space*, Review Bulletin of the Calcutta Mathematical Society **21** (2013), 181–192.
- [7] B.S. Choudhury, K.P. Das and P. Das, *Coupled coincidence point results in partially ordered fuzzy metric spaces*, Annals of Fuzzy Mathematics and Informatics **7** (2014), 619–628.
- [8] P.N. Dutta, B.S. Choudhury and K. Das, *Some fixed point results in Menger spaces using a control function*, Surveys in Mathematics and its Applications **4** (2009), 41–52.
- [9] S.N. Jesic, N.A. Cirovic and, D. O'Regan, *Altering distances and a common fixed point theorem in Menger probabilistic metric spaces*, FILOMAT **31** (2017), 175–181.
- [10] C. Ji, C. Xhu, and Z. Wu, *Some new fixed point theorems in generalized probabilistic metric space*, J. Nonlinear Sci. Appl. **9** (2016), 3735–3743.
- [11] G. Jungk, *Compatible mappings and common fixed points*, Int. J. Math. Math. Sci. **9** (1986), 771–779.
- [12] O. Hadzic and E. Pap, *Fixed Point Theory in Probabilistic Metric Spaces*, Kluwer Academic Publishers, 2001.
- [13] M.S. Khan, M. Swaleh and S. Sessa, *Fixed point theorems by altering distances between the points*, Bull. Austral. Math. Soc. **30** (1984), 1–9.
- [14] P. Ma, J. Guan, Y. Tang, Xu, and Y. Su, *ψ -contraction and $(\phi - \psi)$ -contraction in Menger probabilistic metric space*, Springer Plus **5** (2016), 1–12.
- [15] K. Menger, *Statistical metrics*, Proc. Nat. Acad. Sci. USA **28** (1942), 535–537.

- [16] D. Mihet, *Altering distances in probabilistic Menger spaces*, Nonlinear Analysis **71** (2009), 2734–2738.
- [17] S.V.R. Naidu, *Some fixed point theorems in metric spaces by altering distances*, Czechoslovak Mathematical Journal **53** (2003), 205–212.
- [18] R.P. Pant, *Common fixed points of weakly commuting mappings*, Math. Student **62** (1993), 97–102.
- [19] R.P. Pant, *Common fixed points of noncommuting mappings*, J. Math. Anal. Appl. **188** (1994), 436–440.
- [20] R.P. Pant, *Common fixed points of contractive maps*, J. Math. Anal. Appl. **226** (1998), 251–258.
- [21] R.P. Pant, *R-weakly commutativity and common fixed points*, Soochow J. Math. **25** (1999), 37–42.
- [22] R.P. Pant, *Common fixed points under strict contractive conditions*, J. Math. Anal. Appl. **248** (2000), 327–332.
- [23] V. Pant, *Contractive conditions and common fixed points*, Acta Math. Acad. Paed. Nyir. **24** (2008), 257–266.
- [24] K.P.R. Sastry and G.V.R. Babu, *Some fixed point theorems by altering distances between the points*, Ind. J. Pure. Appl. Math. **30** (6) (1999), 641–647.
- [25] K.P.R. Sastry, S.V.R. Naidu, G.V.R. Babu and G.A. Naidu, *Generalisation of common fixed point theorems for weakly commuting maps by altering distances*, Tamkang Journal of Mathematics **31** (3) (2000), 243–250.
- [26] B. Schweizer and A. Sklar, *Probabilistic Metric Spaces*, Elsevier, North-Holland, (1983).
- [27] V.M. Sehgal and A.T. Bharucha-Reid, *Fixed point of contraction mappings on PM space*, Math. Sys. Theory **6** (2) (1972), 97–100.
- [28] S. Sessa, *On a weak commutativity condition of mappings in fixed point considerations*, Publ. Inst. Math. **32** (1982), 149–153.
- [29] N. Wairojjana, T. Dosenovic, D. Rakik, and D. Gopal, P. Kumam, *An altering distance function in fuzzy metric fixed point theorems*, Fixed Point Theory and Applications **2015** (2015 : 69), 1–19.
- [30] C. Zaharia and N. Cirovic, *A probabilistic fixed point result using altering distance function*, Journal of Function Spaces, Volume 2015, Article ID 919202, 6 pages, <http://dx.doi.org/10.1155/2015/919202>

Krishnapada Das

Department of Mathematics

Sarojini Naidu College for Women

Kolkata, 700028, INDIA

E-mail: kestapm@yahoo.co.in