




Morphological and Molecular Identification of *Spirometra* Tapeworms (Cestoda: Diphyllbothriidae) from Carnivorous Mammals in the Serengeti and Selous Ecosystems of Tanzania

Barakaeli Abdieli Ndosi^{1,2}, Hansol Park¹, Dongmin Lee¹, Seongjun Choe¹ , Yeseul Kang¹, Tilak Chandra Nath^{1,3} , Mohammed Mebarek Bia¹, Chatanun Eamudomkarn⁴, Hyeong-Kyu Jeon^{1,*} , Keeseon S. Eom^{1,*}

¹Department of Parasitology, Parasitology Research Center and Parasite Resource Bank, Chungbuk National University, School of Medicine, Cheongju 28644, Korea; ²Tanzania Wildlife Management Authority, P.O. Box 2658 Morogoro, Tanzania; ³Department of Parasitology, Sylhet Agricultural University, Bangladesh; ⁴Department of Parasitology, Faculty of Medicine, Khon Kaen University, Khon Kaen, Thailand

Abstract: *Spirometra* tapeworms (Cestoda: Diphyllbothriidae) collected from carnivorous mammals in Tanzania were identified by the DNA sequence analysis of the mitochondrial cytochrome c oxidase subunit 1 (*cox1*) and internal transcribed spacer 1 (ITS1), and by morphological characteristics. A total of 15 adult worms were collected from stool samples and carcasses of *Panthera leo*, *Panthera pardus*, and *Crocuta crocuta* in the Serengeti and Selous ecosystems of Tanzania. Three *Spirometra* species: *S. theileri*, *S. ranarum* and *S. erinaceiropaei* were identified based on morphological features. Partial *cox1* sequences (400 bp) of 10 specimens were revealed. Eight specimens showed 99.5% similarity with *Spirometra theileri* (MK955901), 1 specimen showed 99.5% similarity with the Korean *S. erinaceiropaei* and 1 specimen had 99.5% similarity with Myanmar *S. ranarum*. Sequence homology estimates for the ITS1 region of *S. theileri* were 89.8% with *S. erinaceiropaei*, 82.5% with *S. decipiens*, and 78.3% with *S. ranarum*; and 94.4% homology was observed between *S. decipiens* and *S. ranarum*. Phylogenetic analyses were performed with 4 species of *Spirometra* and 2 species of *Dibothriocephalus* (= *Diphyllbothrium*). By both ML and BI methods, *cox1* and ITS1 gave well supported, congruent trees topology of *S. erinaceiropaei* and *S. theileri* with *S. decipiens* and *S. ranarum* forming a clade. The *Dibothriocephalus* species were sisters of each other and collectively forming successive outgroups. Our findings confirmed that 3 *Spirometra* species (*S. theileri*, *S. ranarum*, and *S. erinaceiropaei*) are distributed in the Serengeti and Selous ecosystems of Tanzania.

Key words: *Spirometra erinaceiropaei*, *Spirometra ranarum*, *Spirometra theileri*, *cox1*, ITS1

INTRODUCTION

Tapeworms of the genus *Spirometra* (Cestoda: Diphyllbothriidae) are intestinal parasites of feline and canine mammals. They have complex life cycles involving 2 intermediate hosts, such as freshwater copepods (*Cyclops* spp.), followed by amphibians (tadpoles and frogs) amphibians (tadpoles and frogs) or reptiles (snakes) that consume copepods whilst drinking water [1]. The plerocercoid larva, the sparganum, frequently infects humans by the consumption of raw or undercooked

meat from the second intermediate or transport host, or consumption of untreated water containing proceroid-infected copepods [2].

Few species of *Spirometra* have been reported in Africa. *S. pretoriensis* (= *Lühella pretoriense*) has been found in the bat-eared fox (*Otocyon megalotis*) in South Africa (Baer, 1924, 1926) [3,4] and in the African wild dog (*Lycan pictus*) in DRC Congo (Baer and Fain, 1955) [5]. *S. theileri* (= *Diphyllbothrium theileri*) has been found in the bush cat (*Leptailurus serval*) and the tiger cat (*Felis lybica*) near Pretoria, and in the lion (*Panthera leo*) in DRC Congo (Baer, 1959) [6]. The plerocercoids of *Spirometra*, known as spargana, have been detected in wild herbivorous mammals such as buffalo (*Syncerus caffer*), zebra (*Equus quagga*), wildebeest (*Connochaetes taurinus*), and warthogs (*Phacochoerus africanus*) in Tanzania and Kenya [7,8]. Adults of *S. theileri* were collected from dogs and cats experi-

•Received 9 December 2019, revised 19 October 2020, accepted 25 October 2020.

*Corresponding author (jeonhk@chungbuk.ac.kr; kseom@chungbuk.ac.kr)

© 2020, Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

mentally infected with spargana from warthogs (*Phacochoerus africanus*) in Tanzania [9]. *S. pretoriensis* was found in hyena in Ethiopia [10]. The eggs and proglottids of *Spirometra* had been found in fecal samples of spotted hyenas (*Crocuta crocuta*) and lions in previous studies [8,11,12]. A high prevalence of *Spirometra* infection was reported in lions from the Serengeti and Ngorongoro Crater of Tanzania [13]. *Spirometra* species was commonly found in the intestines of lion and wild dogs in the Luangwa Valley, Zambia [14]. In lions from Tarangire, northern Tanzania, the prevalence of *Spirometra* was higher than that of other intestinal helminths [15].

Previous studies reported *S. ranarum* from lions in the Seronera area [16] and *S. theileri* from leopard (*Panthera pardus*) and spotted hyena in Maswa Game Reserve [17], which were identified by molecular analysis of mitochondrial *cox1* and NADH dehydrogenase subunit 1 (*nad 1*), together with morphological observations. Both of these areas are within the Serengeti ecosystem, which suggested that further studies were needed to clarify the diversity of *Spirometra* in the various geographical areas of Tanzania.

In this study, we investigated the occurrence of *Spirometra* in carnivorous mammals (the main definitive hosts for *Spirometra*) in 2 widely separated and ecologically important areas, the Serengeti ecosystem in northern Tanzania, and the Selous ecosystem in the far eastern part of Tanzania.

MATERIALS AND METHODS

Sample collection and morphological analysis

Tapeworm sample collection was carried out in the Selous and Serengeti ecosystems in Tanzania between 2011 and 2018. These areas were selected owing to its diverse wildlife species. A total of 15 *Spirometra* samples was collected from the stool and the intestines of the leopard, *Panthera pardus*, (n=8), spotted hyena, *Crocuta crocuta*, (n=1), and lion, *Panthera leo*, (n=6). Of 15 samples, 12 were obtained from Loliondo, Maswa, Tomm camp, and Tototo in the Serengeti ecosystem, and 3 were obtained from Utunge Kingupira in the Selous ecosystem (Fig. 1; Table 1).

The collected worms were preserved separately in 70% etha-




Fig. 1. Map of the Serengeti ecosystem in Tanzania indicating the locations of 5 areas where 15 adult worms were collected in 2011-2018.

Table 1. Morphological and molecular identifications of *Spirometra* species tapeworms by the host animals and the collecting areas in Tanzania between 2011 and 2018

Host	Collecting area	cox1/ ITS1 (GenBank No.)	Morphology
Leopard	Loliondo	<i>S. erinacei</i> (MT775875/MN733011) ^b	<i>S. erinacei</i>
Leopard	Loliondo	<i>S. theileri</i> (MT775871/MN737751) ^b	<i>S. theileri</i>
Spotted hyena	Loliondo	<i>S. theileri</i> (MT775870)	<i>S. theileri</i>
Lion	Maswa	<i>S. theileri</i> (MT775876)	<i>S. theileri</i>
Lion	Maswa	<i>S. theileri</i> (MT775877)	<i>S. theileri</i>
Lion	Maswa	np ^a	<i>S. theileri</i>
Lion	Maswa	<i>S. theileri</i> (MT775879)	<i>S. theileri</i>
Lion	Maswa	np	<i>S. theileri</i>
Lion	Maswa	<i>S. theileri</i> (MT775878)	<i>S. theileri</i>
Leopard	Tomm camp	<i>S. theileri</i> (MT775873)	<i>S. theileri</i>
Leopard	Tomm camp	np	<i>S. theileri</i>
Leopard	Tototo	<i>S. ranarum</i> (MT775874/MN733012) ^b	<i>S. ranarum</i>
Leopard	Utunge, Kingupira	<i>S. theileri</i> (MT775872)	<i>S. theileri</i>
Leopard	Utunge, Kingupira	np	<i>S. theileri</i>
Leopard	Utunge, Kingupira	np	<i>S. theileri</i>

^anp, not performed.

^bcox1 and ITS1 were sequenced.

nol and 10% formalin for molecular and morphological observations. For morphological observations, each gravid proglottid of 15 samples preserved in 10% formalin was gently compressed between 2 glass slides before carmine staining. The morphological characteristics of the samples were compared with those described by Baer 1926 [4] and Faust et al. 1929 [18] for carmine-stained specimens, including the width and length of mature and gravid proglottids, number of uterine loops, position of the uterine pore, genital pore, vaginal opening, size and number of testes, and size of eggs.

PCR and DNA sequencing

Genomic DNA was extracted individually from 15 adult tapeworms using a QIAamp DNA Mini Kit following the manufacturer's instructions (Qiagen, Valencia, California, USA). A partial sequence of the mitochondrial *cox1* gene was employed, as described by Jeon et al. [20]. PCR primers for the amplification of internal transcribed spacer 1 (ITS1) were designed from conserved sequences of the end regions of 18S and 5.8S ribosomal DNA to examine the relationships between congeneric and other closely-related species. Primers used to amplify the ITS1 region were Di-ITS1F (5'-AAC AAG GTT TCC GTA GGT GA-3') and Di-ITS1R (5'-AGC AGT CTG CGA TTC ACA TT-3'). The 25 µl reaction mixture contained 50 ng of nuclear DNA, 10 pmol of each ITS1 primer, 2.5 µl of 10×buffer, 12.25 µl of 2×buffer (MgCl₂, dNTP), and 1.25 units of *Taq* polymerase (TAKARA Bio, Inc., Kusatsu, Japan).

Reaction conditions were 3 min at 95°C, over 35 cycles of 1 min at 95°C, 1 min at 58°C, 1 min at 72°C, and a final extension step of 10 min at 72°C. PCR products were purified and ligated into the pGEM-T vector (Promega, Madison, Wisconsin, USA), transformed into *Escherichia coli*, and plated onto LB agar containing X-Gal (20 mg/ml) and ampicillin (100 µg/ml). Plasmid DNA from selected colonies was digested by *Eco*RI to confirm the size of the insert. Ten positive clones from each PCR product were used for nucleotide sequence confirmation.

DNA sequence analysis

DNA sequences of the mitochondrial *cox1* gene and ITS1 were assembled using Geneious R9.1 (Biometer, Auckland, New Zealand). The sequences were identified by comparisons with *cox1* sequences of *Spirometra erinacei*, *S. decipiens*, *S. ranarum*, and *S. theileri*, and with ITS1 sequences of *Diphyllobothrium latum* and *Diphyllobothrium nihonkaiense* in the GenBank database.

A phylogenetic analysis was performed by Bayesian inference (BI) and maximum likelihood (ML) using partial mitochondrial *cox1* (390 bp) sequences of *S. erinacei* (KJ599680), *S. decipiens* (KJ599679), *S. ranarum* (MH298843), *D. nihonkaiense* (EF420138), and *D. latum* (DQ985706) and ITS1 (526 bp) sequences of *D. nihonkaiense* (AB288368 and AB288369). The ML analyses of *cox1* and ITS1 were implemented in MEGA7 [19] with the HKY + G substitution model, determined using a Mod-

eltest. BI analyses were performed using Bayesian Evolutionary Analysis Sampling Trees (BEAST) (version 1.10.4) [20,21] with the HKY substitution, which was chosen using MEGA. Nodes were assessed by bootstrapping with 1,000 pseudoreplicates.

RESULTS

Morphological features of *Spirometra* species in Tanzania

A strobila of *S. erinaceiropaei* collected from a leopard measured 45 cm long. The scolex was spoon-shaped, measuring 0.4 mm in diameter and 1.0 mm in length. The proglottid segments measured 5 to 9 mm in width and 5 to 8 mm in length and became broad on the terminal gravid segments. The cirrus pouch appeared just behind the male genital opening close to the vaginal pore, projecting towards the vaginal duct. The uterus covered a narrow field comprising 7 to 8 complete coils, with 4 internal loose coils located slightly behind the outer pile. The eggs were numerous, ellipsoidal, operculated, and up to 57 μm with a diameter of 34 μm , and testes and vitellaria extended to the lateral margin of the uterus and united at the anterior portion (Fig. 2A).

A strobila of *S. ranarum* collected from the stool of a lion was 75 cm long and 3 mm wide. The mature and gravid proglottids were 12.0 mm in width and 3 mm in length, on average. The uterus comprised 3-3.5 loops with a butterfly or bow-tie appearance, and adopted a diagonal direction in the second turn. The genital and vaginal pores were situated ventrally on the midline in the anterior third of the proglottid. The male

genital aperture extended to the anterior border of the segment and the median was close to the female aperture on its lateral side. The testes measured 74 μm in diameter. The ovaries were highly dendritic and connected to the uterus. The eggs were numerous, 57 to 60 μm in length and 34.2 to 35.3 μm in transverse diameter (Fig. 2B; Table 2).

The strobilae of *S. theileri* collected from the intestine of a leopard and spotted hyena were 45 cm and 134 cm in length, respectively, with varying shapes and sizes. The first segment measured 0.4 mm wide and 0.02 mm long, up to 0.6 to 1.0 mm in width, and 0.15 to 0.25 mm in length. The uterus consisted of 3-4.5 loops on each side of the proglottids. The genital pore was located ventrally on the midline in the anterior 1/5 of the proglottids. The vaginal pore was behind the male genital pore. The uterine pore was situated posterior to the vagina, and numerous thick-shelled and operculated eggs were present, 54 to 59 μm in length and 20 to 24 μm in width. The dumbbell-shaped ovary was located near the posterior margin of the proglottid and connected to the uterus (Fig. 2C; Table 2).

Sequence similarity and phylogenetic relationships

Among the 15 samples, 10 partial *cox1* sequences were successfully obtained and deposited in GenBank (under accession numbers MT775875 for *S. erinaceiropaei*, MT775874 for *S. ranarum*, and MT775870-775873, MT775876-775879 for *S. theileri*). Partial *cox1* sequences (-400 bp) from 8 isolates from Tanzania showed 99.5% (400/430) similarity with *cox1* of previously sequenced Tanzanian *S. theileri* (MK955901). Addi-




Fig. 2. Mature proglottids of *Spirometra* species collected from carnivorous mammals in the Serengeti and Selous ecosystems of Tanzania (stained with acetocarmine). (A) *S. erinaceiropaei*. (B) *S. ranarum*. (C) *S. theileri*. Characteristic genital organs, cirrus sac (CS), genital pore (GP), ovary (OV), vaginal pore (VP), and uterus (U) are designated in (A-C).

Table 2. Morphometric features of 3 *Spirometra* species from carnivorous mammals in Serengeti National Park of Tanzania (unit = mm)

Organs	Organ feature		<i>S. erinaceiropaei</i> ^a	<i>S. ranarum</i> ^b	<i>S. theileri</i> ^c
Scolex	Spatulate	Width	0.42	-	-
		Length	1.03	-	-
Proglottids (n = 10)	Trapezoid	Width	5.00-9.00	12.00	0.60-0.65
		Length	5.00-8.00	3.01	0.15-0.25
Uterus	Coiling	Loops	7.00-8.00	3.00	3-4.50
Cirrus pouch	Oval	Length		0.17-0.25	0.16
		Width		0.17-0.25	0.13
Cirrus	Cylindrical	Length	0.30	0.30	0.30
		Width	0.13	0.19	0.13
Seminal vesicle	Elliptical	Length	0.13	0.19-0.22	0.13
		Width	0.14	0.19-0.22	0.14
Uterine terminal ball	Oval	Diameter	0.06	0.04-0.04	0.07-0.08
Vaginal opening	Crescentic	Width	0.26	0.25	0.25
Testes	Polygonal	Diameter	0.06	0.06	0.11-0.12
Eggs	Operculate	Length	0.06	0.05-0.07	0.05-0.06
		Width	0.03	0.03-0.04	0.03-0.04

^aIsolated from a leopard (*Panthera pardus*), ^ba lion (*Panthera leo*), and ^ca hyena (*Crocuta crocuta*).

tionally, *cox1* of one specimen (No. B473) showed 99.5% similarity with that of Korean *S. erinaceiropaei* (KJ599680), and *cox1* of another specimen (No. 457-3) showed 99.5% similarity with that of Myanmar *S. ranarum* (MH298843). Sequence differences in the partial *cox1* sequence of Tanzanian *Spirometra* ranged from 0% to 10.1%. Samples were identified as *S. theileri* (n=8), *S. erinaceiropaei* (n=1), and *S. ranarum* (n=1) by *cox1* sequence analysis. *Spirometra theileri*, *S. erinaceiropaei* and *S. ranarum* were all found in Serengeti ecosystem while only *S. theileri* was revealed in Selous ecosystem (Table 1).

The ITS1 sequences of 4 *Spirometra* species, *S. theileri* (MN737751, 716 bp), *S. erinaceiropaei* (MN733011, 732 bp), *S. decipiens* (MN733010, 801 bp), and *S. ranarum* (MN733012, 842 bp), showed length differences. The ITS1 sequence of *S. theileri* showed 89.8% similarity with that of *S. erinaceiropaei* (MN733011), 82.5% similarity with *S. decipiens* (MN733010), and 78.3% similarity with *S. ranarum* (MN733012), and 94.4% similarity was observed between *S. decipiens* and *S. ranarum*.

Phylogenetic analyses were performed with 4 species of *Spirometra* and 2 species of *Dibothriocephalus* (= *Diphyllobothrium*).

The *cox1* and ITS1, by both ML and BI methods, gave well supported, congruent trees topology of *S. erinaceiropaei* and *S. theileri* with *S. decipiens* and *S. ranarum* forming a clade (Fig. 3). The *Dibothriocephalus* species were sisters of each other, and, collectively, forming successive outgroups.

DISCUSSION

Spirometra can be identified based on morphological and molecular characters. In this study, *S. erinaceiropaei*, *S. theileri*, and *S. ranarum* were obtained from lions, spotted hyenas, and leopards in the Selous and Serengeti ecosystems, the 2 most important ecosystems in Tanzania, with a high diversity of wildlife species. Adult *Spirometra* were analyzed based on genital morphology and mitochondrial *cox1* and ITS1 sequences. Partial *cox1* sequences obtained from 10 of the 15 samples revealed 3 species of *Spirometra* were present.

Based on morphological observations, uterine shapes of collected *Spirometra* species vary with respect to the number of coiled shapes, with 7-8 complete coils in *S. erinaceiropaei*, 3-3.5 in *S. ranarum*, and 3 on one side and 4 complete coils on another side in *S. theileri*, similar to previous findings [5,22]. The ovaries were clearly dendritic in shape, but the eggs were operculate with slight differences in size among taxa.

Major differences between *S. theileri* and *S. pretoriensis* included the number of uterine loops and shape of the cirrus pouch. The uterus of *S. theileri* consisted of 3-4.5 coils and a cirrus pouch with communication through a short canal with a second smaller vesicular seminis, while the uterus of *S. pretoriensis* formed a single large loop on either side and the cirrus pouch contained a large vesicular seminis of variable shape. The distinct morphological features of *S. ranarum* relative to other *Spirometra* species included posterior uterine coils larger




Fig. 3. Phylogenetic trees of *Spirometra* species based on mitochondrial (A) *cox1* and (B) ITS1 sequences. Numbers above the branches represent bootstrap values for trees obtained by maximum likelihood and Bayesian inference.

than the terminal uterine ball and coiling of the uteri diagonally rather than spirally. The widths of gravid proglottids and the uterine ball of *S. ranarum* were larger than those of other *Spirometra* species, including *S. erinaceieuropaei*, *S. decipiens*, *S. theileri*, and *S. pretoriensis*.

Mitochondrial *cox1* and *nad1* sequences of *S. ranarum* and *S. theileri* have been reported [16,17]. Our results provide a basis for the accurate molecular identification of *Spirometra* species from Tanzania by providing sequences for ITS1 and partial *cox1* of 10 specimens. The 3 *Spirometra* species identified have now been found in Tanzania and various Asian countries, including China, Japan, Korea, Laos, and Thailand. The genetic divergence of the partial *cox1* sequences among the several species of *Spirometra* (*S. erinaceieuropaei*, *S. decipiens*, *S. ranarum*, and *S. theileri*) ranged from 0% to 10.1%. Phylogenetic

analyses based on mitochondrial DNA sequences with ML and BI methods showed that *S. erinaceieuropaei* and *S. theileri* were separated from other *Spirometra* species, and *S. decipiens* and *S. ranarum* belonged to the same clade.

In addition to Seronera and Maswa, which have been evaluated in previous studies [16,17], our study is reporting 4 new endemic areas of *Spirometra* species such as Loliondo Game Controlled Area, Tomm camp, and Tototo in the Serengeti ecosystem in northern Tanzania and Utunge-Kingupira in Selous ecosystem in far eastern Tanzania. Among the 15 specimens collected, 13 specimens were identified as *S. theileri* collected from the lion ($n=6$) in Maswa area, 3 from the leopard in Utunge-Kingupira area, 2 from the leopard in Tomm camp area and 2 specimens collected from leopard ($n=1$) and spotted hyena ($n=1$) in Loliondo area. *S. ranarum* was obtained from a

leopard in the Tototo area and *S. erinaceieuropaei* was obtained from a leopard in the Loliondo area.

In conclusion, we provide the first evidence that the leopard is a definitive host of *S. erinaceieuropaei*, *S. theileri* and *S. ranarum* and clarified the diversity of *Spirometra* species in 4 new endemic areas. These findings suggest that wild carnivorous mammals could be as reservoir hosts for *Spirometra*, which cause human sparganosis in Africa. The infection of wildlife species with *Spirometra* is likely related to the distribution of *Spirometra* species in various geographical locations. However, further studies are needed to evaluate the *Spirometra* species responsible for human sparganosis in the region.

ACKNOWLEDGMENTS

This work was supported by the National Research Foundation of Korea (NRF- 2017K1A3A1A09085607). Materials were provided by the Parasite Resource Bank of Korea (PRB000720).

CONFLICT OF INTEREST

The authors declare no conflict of interest related to this work.

REFERENCES

- Nobrega-Lee M, Hubbard G, Loverde P, Carvalho-Queiroz C, Conn DB, Rohde K, Dick EJ Jr, Nathanielsz P, Martin D, Siler-Khodr T, Schlabritz-Loutsevitch N. Sparganosis in wild-caught baboons (*Papio cynocephalus anubis*). *J Med Primatol* 2007; 36: 47-54. <https://doi.org/10.1111/j.1600-0684.2006.00177.x>
- Anantaphruti MT, Nawa Y, Vanvanitchai Y. Human sparganosis in Thailand: an overview. *Acta Trop* 2011; 118: 171-176. <https://doi.org/10.1016/j.actatropica.2011.03.011>
- Baer JG. Contribution à la faune helminthologique Sud-Africaine-Note préliminaire. *Annales de Parasitologie* 1924; 2: 237-247 (in French).
- Contribution to the helminth fauna of South Africa. Mammalian cestodes. In Baer JG ed, 11th and 12th Report of the Director of Veterinary Education and Research. Pretoria, South Africa. Department of Agriculture. 1926, pp 61-136.
- Baer JG, Fain A. Cestodes. Report d'Exploration Parcs Nationaux de l'Upemba. Fasc. 1955, pp 36.
- Baer JG. Helminthes parasites. Exploration des Parcs Nationaux du Congo Belge. Brussels, Belgium. 1959, pp 163.
- Nelson GS, Pester FRN, Rickman R. The significance of wild animals in the transmission of cestodes of medical importance in Kenya. *Trans R Soc Trop Med Hyg* 1965; 59: 507-524. [https://doi.org/10.1016/0035-9203\(65\)90153-7](https://doi.org/10.1016/0035-9203(65)90153-7)
- Müller-Graf CD. A coprological survey of intestinal parasites of wild lions (*Panthera leo*) in the Serengeti and the Ngorongoro Crater, Tanzania, east Africa. *J Parasitol* 1995; 8: 812-814.
- Opuni EK, Muller RL. Studies on *Spirometra theileri* (Baer, 1925) n. comb. 1. Identification and biology in the laboratory. *J Helminthol* 1974; 48: 15-23. <https://doi.org/10.1017/s0022149x00022550>
- Graber M. Diphyllobothriosis and sparganosis in tropical Africa. *Rev Elev Méd Vét Pays Trop* 1981; 34: 303-311 (in French).
- Bjork KE, Averbeck AG, Stromberg BE. Parasites and parasite stages of free-ranging wild lions (*Panthera leo*) of northern Tanzania. *J Zoo Wildl Med* 2000; 31: 56-61. [https://doi.org/10.1638/1042-7260\(2000\)031\[0056:PAPSOF\]2.0.CO;2](https://doi.org/10.1638/1042-7260(2000)031[0056:PAPSOF]2.0.CO;2)
- Müller-Graf CD, Woolhouse ME, Packer C. Epidemiology of an intestinal parasite (*Spirometra* spp.) in two populations of African lions (*Panthera leo*). *Parasitol* 1999; 118: 407-415. <https://doi.org/10.1017/s0031182098003813>
- Engh AL, Nelson KG, Peebles R, Hernandez AD, Hubbard KK, Holekamp KE. Coprologic survey of parasites of spotted hyenas (*Crocuta crocuta*) in the Masai Mara National Reserve, Kenya. *J Wildl Dis* 2003; 39: 224-227. <https://doi.org/10.7589/0090-3558-39.1.224>
- Berensten AR, Becker MS, Stockdale-Walden H, Matandiko W, McRobb R, Dunbar MR. Survey of gastrointestinal parasite infection in African lion (*Panthera leo*), African wild dog (*Lycaon pictus*) and spotted hyaena (*Crocuta crocuta*) in the Luangwa Valley, Zambia. *Afr Zool* 2012; 47: 363-368. <https://doi.org/10.1080/15627020.2012.11407561>
- Kavana NJ, Kassuku AA, Kasanga CJ. Prevalence of *Spirometra* species and other gastrointestinal helminths in wild lions (*Panthera leo*) in Tarangire National Park, northern Tanzania. *Int J Microbiol Immunol Res* 2015; 4: 10-14.
- Eom KS, Park HS, Lee DM, Choe SJ, YS Kang, Bia MM, Lee SH, Keyyu J, Fymagwa R, Jeon HK. Molecular and morphologic identification of *Spirometra ranarum* found in the stool of African lion, *Panthera leo* in the Serengeti plain of Tanzania. *Korean J Parasitol* 2018; 56: 379-383. <https://doi.org/10.3347/kjp.2018.56.4.379>
- Eom KS, Park H, Lee D, Choe S, Kang Y, Bia MM, Ndosi BA, Nath TC, Eamudomkarn C, Keyyu J, Fymagwa R, Mduma S, Jeon HK. Identity of *Spirometra theileri* from a leopard (*Panthera pardus*) and spotted hyena (*Crocuta crocuta*) in Tanzania. *Korean J Parasitol* 2019; 57: 639-645. <https://doi.org/10.3347/kjp.2019.57.6.639>
- Faust EC, Campbell HE, Kellogg CR. Morphological and biological studies on the species of *Diphyllobothrium* in China. *Am J Epidemiol* 1929; 9: 560-583. <https://doi.org/10.1093/oxfordjournals.aje.a121670>
- Jeon HK, Park HS, Lee DM, Choe SJ, Kim KH, Huh S, Sohn WM, Chai JY, Eom KS. Human infections with *Spirometra* decipiens plerocercoids identified by morphologic and genetic analyses in Korea. *Korean J Parasitol* 2015; 53: 299-305. <https://doi.org/10.3347/kjp.2015.53.3.299>
- Tamura K, Peterson D, Peterson N, Stecher G, Nei M, Kumar S. MEGA7 molecular evolutionary genetics analysis using maxi-

- mum likelihood evolutionary distance, and maximum parsimony methods. *Mol Biol Evol* 2011; 25: 2731-2739. <https://doi.org/10.1093/molbev/msr121>
21. Suchard MA, Kitchen CMR, Sinsheimer JS, Weiss RE. Hierarchical phylogenetic models for analyzing multipartite sequence data. *Systema Biol* 2003; 52: 649-664. <https://doi.org/10.1080/10635150390238879>
22. Jeon HK, Park HS, Lee DM, Choe SJ, YS Kang, Bia MM, Lee SH, Sohn WM, Hong SJ, Chai JY, Eom KS. Genetic and morphologic identification of *Spirometra ranarum* in Myanmar. *Korean J Parasitol* 2018; 56: 275-280. <https://doi.org/10.3347/kjp.2018.56.3.275>