


Effects of *Bacillus*-based probiotics on growth performance, nutrient digestibility, and intestinal health of weaned pigs

Daye Mun^{1#}, Hyunjin Kyoung^{2#}, Myunghwan Kong^{2#}, Sangdon Ryu¹, Ki Beom Jang³, Jangryeol Baek², Kyeong Il Park², Minho Song^{2*} and Younghoon Kim^{1*}

¹Department of Agricultural Biotechnology and Research Institute of Agriculture and Life Science, Seoul National University, Seoul 08826, Korea

²Division of Animal and Dairy Science, Chungnam National University, Daejeon 34134, Korea

³Department of Animal Science, North Carolina State University, Raleigh, NC 2769, USA



Received: Aug 5, 2021
Revised: Aug 19, 2021
Accepted: Sep 27, 2021

#These authors contributed equally to this work.

*Corresponding author

Minho Song
Division of Animal and Dairy Science,
Chungnam National University,
Daejeon 34134, Korea.
Tel: +82-42-821-5776
E-mail: mhsong@cnu.ac.kr

Younghoon Kim
Department of Agricultural
Biotechnology and Research Institute
of Agriculture and Life Science, Seoul
National University, Seoul 08826,
Korea.
Tel: +82-2-880-4808
E-mail: ykeys2584@snu.ac.kr

Copyright © 2021 Korean Society of
Animal Sciences and Technology.
This is an Open Access article
distributed under the terms of the
Creative Commons Attribution
Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted
non-commercial use, distribution, and
reproduction in any medium, provided
the original work is properly cited.

ORCID

Daye Mun
<https://orcid.org/0000-0002-3470-7632>

Abstract

Bacillus is characterized by the formation of spores in harsh environments, which makes it suitable for use as a probiotic for feed because of thermostability and high survival rate, even under long-term storage. This study was conducted to investigate the effects of *Bacillus*-based probiotics on growth performance, nutrient digestibility, intestinal morphology, immune response, and intestinal microbiota of weaned pigs. A total of 40 weaned pigs (7.01 ± 0.86 kg body weight [BW]; 28 d old) were randomly assigned to two treatments (4 pigs/pen; 5 replicates/treatment) in a randomized complete block design (block = BW and sex). The dietary treatment was either a typical nursery diet based on corn and soybean meal (CON) or CON supplemented with 0.01% probiotics containing a mixture of *Bacillus subtilis* and *Bacillus licheniformis* (PRO). Fecal samples were collected daily by rectal palpation for the last 3 days after a 4-day adaptation. Blood, ileal digesta, and intestinal tissue samples were collected from one pig in each pen at the respective time points. The PRO group did not affect the feed efficiency, but the average daily gain was significantly improved ($p < 0.05$). The PRO group showed a trend of improved crude protein digestibility ($p < 0.10$). The serum transforming growth factor- $\beta 1$ level tended to be higher ($p < 0.10$) in the PRO group on days 7 and 14. There was no difference in phylum level of the intestinal microbiota, but there were differences in genus composition and proportions. However, β -diversity analysis showed no statistical differences between the CON and the PRO groups. Taken together, *Bacillus*-based probiotics had beneficial effects on the growth performance, immune system, and intestinal microbiota of weaned pigs, suggesting that *Bacillus* can be utilized as a functional probiotic for weaned pigs.

Keywords: *Bacillus*-based probiotics, Weaned pigs, Growth performance, Immune response, Intestinal health

Hyunjin Kyoung
<https://orcid.org/0000-0001-5742-5374>
 Myunghwan Kong
<https://orcid.org/0000-0002-7207-9382>
 Sangdon Ryu
<https://orcid.org/0000-0001-5338-8385>
 Ki Beom Jang
<https://orcid.org/0000-0001-6794-9569>
 Jangryeol Baek
<https://orcid.org/0000-0002-1722-2641>
 Kyeong Il Park
<https://orcid.org/0000-0002-3590-3993>
 Minho Song
<https://orcid.org/0000-0002-4515-5212>
 Younghoon Kim
<https://orcid.org/0000-0001-6769-0657>

Competing interests

No potential conflict of interest relevant to this article was reported.

Funding sources

This work was supported by grants from National Agricultural Cooperative Federation (NACF) and a grant from the Agricultural and livestock products safe distribution and consumption technology development program (Project No. 320111-1), Korea Institute of Planning and Evaluation for Technology in Food, Agriculture and Forestry (IPET), Korea and the National Research Foundation of Korea (NRF) Grant funded by the Korean government (NRF-2021R1A2C3011051).

Acknowledgements

Not applicable.

Availability of data and material

Upon reasonable request, the datasets of this study can be available from the corresponding author.

Authors' contributions

Conceptualization: Mun D, Kyoung H, Kong M, Song M, Kim Y.
 Data curation: Mun D, Kyoung H, Kong M, Ryu S, Jang KB, Baek J, Park KI, Song M, Kim Y.
 Formal analysis: Mun D, Song M, Kim Y.
 Methodology: Mun D, Kyoung H, Kong M, Ryu S.
 Software: Mun D, Kyoung H, Kong M, Ryu S.
 Validation: Jang KB, Baek J, Park KI, Song M, Kim Y.
 Investigation: Mun D, Kyoung H, Kong M.
 Writing - original draft: Mun D, Kyoung H, Kong M, Song M, Kim Y.
 Writing - review & editing: Mun D, Kyoung H, Kong M, Ryu S, Jang KB, Baek J, Park KI, Song M, Kim Y.

Ethics approval and consent to participate

The experimental protocol for this study was reviewed and approved by the Institutional Animal Care and Use Committee of Chungnam National University, Daejeon, Korea (approval #CNU-00910).

INTRODUCTION

Weaning is a stressful period for young pigs because several stressors including separation from mother sow, diet form change, and environmental changes [1]. In addition, weaned pigs must acclimatize to less digestible plant-based solid feed rather than easily digestible sow milk [2]. These multiple stress factors cause loss of appetite, which leads to insufficient energy intake for the maintenance of gastrointestinal health [3]. Additionally, the damage of intestinal epithelial layers because of weaning stress reduces trans-mucosal resistance [2]. Therefore, piglets in the weaning period are sensitive to infection by enteric pathogens, such as *Escherichia coli*, which leads to diarrhea [4,5].

To prevent post-weaning diarrhea and to improve growth performance and gut health of piglets, various antibiotics have been used as growth promoters [6,7]. However, because of potential safety issues, such as antimicrobial resistance or antibiotic residue, that may occur when antibiotics are overused, most countries have banned the use of antibiotics as growth promoters [8,9]. Therefore, a number of studies are being conducted to develop powerful alternatives to antibiotics for maintaining the health of weaned pigs [10].

Probiotics are defined by the FAO/WHO group as “live microorganisms which, when administered in adequate amounts, confer a good health benefit on the host” [11,12]. It has been well studied that probiotics could be alternatives to antibiotics because of their ability to reduce diarrhea and improve the growth performance of weaned pigs [13]. General strains of probiotic species are mostly isolated from consumers of probiotics, including humans, chickens, or pigs. Commonly, *Bacillus*, *Lactobacillus*, *Bifidobacterium*, *Enterococcus*, and *Streptococcus* are used as probiotics [11]. Among them, *Bacillus* is an attractive bacterium for administration as a probiotic in swine feed because of its ability to form spores, which allows the bacterium to withstand high-temperature and the low-pH of gastric juices [14]. Additionally, spore-forming bacteria, such as *Bacillus subtilis* and *Bacillus amyloliquefaciens*, allow for long-term storage without losing viability [15]. According to previous studies, supplementation of *Bacillus* has been shown to improve growth performance by enhancing gut barrier integrity and relieving post-weaning diarrhea [14,16].

Although several previous studies have shown that *Bacillus* has beneficial properties for the host, the results of its effects on the immune system or intestinal microbiota of weaned pigs are still inconsistent. Therefore, this study was conducted to investigate the effect of *Bacillus*-based probiotics on growth performance, nutrient digestibility, intestinal morphology, immune response and intestinal microbiota of weaned pigs.

MATERIALS AND METHODS

Experimental design, animals, and diets

The procedure of experiment was reviewed and approved by the Institutional Animal Care and Use Committee of Chungnam National University, Daejeon, Korea (approval #CNU-00910). This study was carried out at the facility of Animal Research Center of Chungnam National University. A total of healthy 40 weaned pigs ([Landrace × Yorkshire] × Duroc; 7.01 ± 0.86 kg; 28 d old) were assigned to 2 treatments in a randomized complete block design (block = body weight [BW] and sex) with 4 pigs per pen and 5 replicated pens per treatment. The dietary treatment was either a typical nursery diet based on corn and soybean meal (CON) or CON supplemented with 0.01% probiotics (PRO). Both dietary treatments were formulated to meet or exceed the requirements of the National Research Council [17] of weaned pigs and to have comparable metabolizable energy, crude protein (CP), and lysine levels (Table 1). Animal plasma, antibiotics, and zinc

Table 1. Composition of experimental diet for weaning pigs (as-fed basis)

Items	Weaner
Ingredients (%)	
Corn	49.86
Whey powder	12.50
Soybean meal (44%)	25.00
Soy protein concentrate	6.25
Soybean oil	3.00
Limestone	1.14
Mono-calcium phosphate	1.05
Vitamin premix ¹	0.20
Mineral premix ²	0.20
L-Lysine-HCl	0.45
DL-Methionine	0.16
L-Threonine	0.13
L-Valine	0.06
Calculated energy and nutrient contents	
Metabolizable energy (kcal/kg)	3,465
Crude protein (%)	21.26
Calcium (%)	0.81
Phosphorus (%)	0.65

¹Provided per kilogram of complete diet: vitamin A, 12,000 IU; vitamin D₃, 2,500 IU; vitamin E, 30 IU; vitamin K₃, 3 mg; D-pantothenic acid, 15 mg; nicotinic acid, 40 mg; choline, 400 mg; vitamin B₁₂, 12 µg.

²Provided per kilogram of complete diet: Fe, 90 mg from iron sulfate; Cu, 8.8 mg from copper sulfate; Zn, 100 mg from zinc oxide; Mn, 54 mg from manganese oxide; I, 0.35 mg from potassium iodide; Se, 0.30 mg from sodium selenite.

oxide were excluded from the piglet diets to avoid antimicrobial or physiological effects, and all feeds were provided in meal form. The dietary probiotics used in this study were a commercial product containing a mixture of *Bacillus subtilis* and *Bacillus licheniformis*. Pigs were housed in an environmentally controlled room and each pen was equipped with a feeder and water. Also, piglets were allowed *ad libitum* access to feed and water throughout the experiment.

Data and sample collection

The BW of pigs was recorded at d 0 of the experiment and at the last day of the experiment. The amount of feed added per pen was recorded, and feed residue were weighed from each pen on the last day of the experiment. The difference between the last BW and the initial BW was divided by the number of experimental days to determine the average daily gain (ADG). The feed intake per piglet was divided by the number of experimental days and was determined as the average daily feed intake (ADFI). To obtain the feed efficiency, the gain to feed ratio (G:F) was calculated by dividing ADG by ADFI for each pen. Diarrhea scores of all piglets during the experiment were recorded daily for 2 weeks from the first day after weaning. The diarrhea score for each pig was visually assessed daily on a scale of 1 to 5 (1 = dry feces, 2 = normal feces, 3 = slightly moist feces, 4 = semiliquid feces, mild diarrhea, and 5 = watery feces, severe diarrhea). The frequency of diarrhea (%) was calculated by counting the number of days the pigs had a diarrhea score of 4 or greater [18]. Whole blood samples were collected from one randomly selected pig in each pen using the jugular vein and stored in ethylenediaminetetraacetic acid tubes with anticoagulant or serum tubes (Becton Dickinson Vacutainer Systems, Franklin Lakes, NJ, USA) to measure total white blood cells (WBC) counts or inflammatory cytokines on d 7 and 14. Pigs were each fed a dietary

treatment containing 0.2% indigestible chromium oxide during the last week of the experimental period to determine the apparent ileal and total tract digestibility of nutrients. For the collection of fresh fecal samples, rectal palpation was performed from one randomly selected pig in each pen for the last 3 days after the 4-day adjustment period. The collected stool samples and diet samples were stored at -20°C until analysis. Fecal samples for metagenome analysis were freshly collected from three randomly selected pigs in each treatment group on the last day of the experiment by rectal palpation with a sterile cotton swab, and the samples were placed in a sterile tube. The samples were taken to the laboratory on ice and stored at -80°C in a deep freezer until further analysis. On the last day of the experiment, one randomly selected piglet in each pen was anesthetized using 2 mL of suxamethonium chloride (Succicholine[®], Ilsung Pharm, Seoul, Korea) by an intramuscular injection. Immediately after anesthesia, pigs were euthanized by CO_2 gas [19]. Digesta samples of the pig gastrointestinal tract were collected from the terminal ileum before the ileocecal valve and were stored at -20°C until the time of analysis. For histological analysis, 3-cm sections were taken from the middle of the duodenum, jejunum, and ileum. The tissue samples were gently flushed with 0.05 M phosphate buffered saline (pH 7.2) and were then fixed in 10% neutral buffered formalin.

Sample measurements and analysis

Before chemical analysis, frozen ileal digesta, fecal samples, and feed samples were thawed, dried at 60°C for 72 h and finely ground. Dry matter (DM, method 930.15) and nitrogen (method 999.03) of all samples were analyzed according to the procedures outlined in the Association of Official Analytical Chemists [20]. The gross energy (GE) was analyzed for each sample using a bomb calorimeter (Parr 1281 Bomb Calorimeter, Parr Instrument, Moline, IL, USA), and nitrogen was measured using a Kjeltec 2300 Analyzer (Foss Tecator AB, Hoeganaes, Sweden). Chromium levels in the samples were measured by UV spectrophotometer (Hitachi Z-5000, Hitachi High-Technologies, Tokyo, Japan). The apparent total tract digestibility (ATTD) and apparent ileal digestibility (AID) of the DM, CP, and GE were calculated for each sample as described by Stein et al. [21]. The fixed tissue samples taken from the small intestine were embedded in paraffin wax, sectioned to $5\ \mu\text{m}$ using a microtome, and stained with hematoxylin and eosin. Stained sections were examined using a charge-coupled device camera-equipped microscope (TE2000, Nikon, Tokyo, Japan) with digital imaging software for measurements and counts (NIS Elements BR; Nikon). The 15 longest and straightest villi and their associated crypts were selected to evaluate morphological changes [22]. The morphometric measurements of intestinal tissue were carried out as follows: villus height; crypt depth; villus height to crypt depth ratio (VH:CD); and the number of goblet cells. Whole blood samples were analyzed by an automated hematology analyzer (scil Vet abc hematology analyzer; scil animal care company, Viernheim, Germany) calibrated for porcine blood for total WBC counts. The blood samples for serum were allowed to clot for 2 h at room temperature and then overnight at 4°C followed by centrifugation at $3,000\times g$ for 15 min at 4°C . Serum concentrations of cytokines and cortisol were measured by using a porcine-specific tumor necrosis factor- α (TNF- α) or transforming growth factor- β 1 (TGF- β 1) ELISA kit (R&D Systems, Minneapolis, MN, USA), a porcine-specific cortisol ELISA kit (Cusabio, Wuhan, China), a porcine C-reactive protein or an immunoglobulin ELISA kit (Abnova, Taipei City, Taiwan), and measurements were conducted according to the manufacturer's protocols. The optical density was measured at 450 nm, and all concentrations were calculated using a standard curve.

DNA extraction, amplification, and sequencing

DNA was extracted from fecal samples using a PowerSoil DNA Isolation kit (MO BIO Laboratories, Carlsbad, CA, USA) according to the manufacturer's protocol. DNA concentration

and quality were determined by measuring absorbance at 230, 260, and 280 nm using a spectrophotometer (SpectraMax ABS Plus, Molecular Devices, San Jose, CA, USA). To amplify the V4 region from the 16S rRNA gene, 341F/805R primers (forward, 5'-CCT ACG GGN GGC WGC AG-3'; reverse, 5'-GAC TAC HVG GGT ATC TAA TCC-3') were used. 16S rRNA gene amplicon sequencing was performed on the Illumina MiSeq platform at Macrogen (Seoul, Korea).

Metagenomic analysis

The raw paired-end reads were processed using Quantitative Insights Into Microbial Ecology 2 (QIIME2; version 2020.6.0). Quality control (e.g., filtering, trimming the sequence and removing chimeric sequences) was performed on demultiplexed sequences with the DADA2 pipeline. Taxonomic assignment was performed with the Greengenes reference database (13_8 version) into operational taxonomic units (OTUs) with 97% similarity. To assess within-sample diversity, microbial α -diversity was calculated based on several metrics (Shannon, Chao1, and Simpson index) with the script “diversity α -group-significance” after making phylogenetic trees using the mafft and FastTree programs of QIIME2. To evaluate between-group diversity, β -diversity using phylogenetic-based weight and unweighted UniFrac distances were measured through the script “diversity beta-group-significance”. Additionally, based on the weight and unweighted UniFrac distance metrics, principal coordinate analysis (PCoA) plots were used to visualize β -diversity.

Statistical analyses

The PROC GLM procedures of SAS (SAS Inst., Cary, NC, USA) were used to analyze all data except frequency of diarrhea and fecal microbiota in a randomized complete block design, and the experimental unit was the pen. The statistical model for growth performance, nutrient digestibility, intestinal morphology, and immune responses of piglets included effects of dietary treatment as a fixed effect. The results are presented as the mean \pm standard error of the mean. A chi-squared test was used for the diarrhea frequency. Statistical analysis of the fecal microbial composition was performed using QIIME2 and GraphPad Prism software version 7 (La Jolla, CA, USA). Permutational multivariate analysis of variance (PERMANOVA) was performed using QIIME2 to establish whether the profile of fecal microbiota between the two groups was significantly different and was based on UniFrac distance matrices. Statistical significance and tendency were considered at $p < 0.05$ and $0.05 \leq p < 0.10$, respectively.

RESULTS AND DISCUSSION

Growth performance and nutrient digestibility

Weaned pigs fed PRO had significantly improved ADG (428.57 vs. 386.19 g/d; $p < 0.05$) (Table 2). There were no significant differences in ADFI, G:F, and frequency of diarrhea between CON and PRO. Additionally, as shown in Table 3, there were no differences in the AID of DM, CP, and GE and the ATTD of DM and GE between CON and PRO at the end of the experiment. Pigs fed PRO tended to have improved ATTD of CP (87.25% vs. 84.13%; $p < 0.10$). Several studies have shown that the addition of *Bacillus* to pig feed could improve growth performance or reduce the incidence of diarrhea [16,23]. However, some previous studies failed to observe such positive effects on growth performance [24–26]. These discrepancies in the effect of *Bacillus* on pig growth performance may be caused by various factors, such as diet formula, different *Bacillus* species, dose, pig age, or environmental features [15]. In this study, the AID of DM, CP, and GE and the ATTD of either DM or GE were not affected by *Bacillus*-based probiotics. However, pigs supplemented with *Bacillus*-based probiotics tended to show improved ATTD of CP. Based on these results, it can

Table 2. Effects of *Bacillus*-based probiotics on overall growth performance of weaned pigs¹

Item	Dietary treatments		SEM	p-value
	CON	PRO		
Initial BW (kg)	7.01	7.02	0.41	0.917
Final BW (kg)	23.23	25.02	0.81	0.155
Feed intake (kg)	27.59	29.66	1.83	0.491
ADG (g/d)	386.19	428.57	11.85	0.040
ADFI (g/d)	656.90	706.19	43.58	0.491
G:F (g/g)	0.588	0.607	0.03	0.739
Frequency of diarrhea (%) ²	14.29	13.27		0.845

¹Each value presented as the least square mean of 5 replicates (4 pigs/pen).

²Frequency of diarrhea (%) = (Number of pigs with diarrhea / number of pen days) × 100. The data was analyzed by chi-squared test.

CON, control diet based on soy-bean meal diet; PRO, control + 0.01% *Bacillus*-based probiotics; BW, body weight; ADG, average daily gain; ADFI, average daily feed intake; G:F, gain-to-feed ratio.

Table 3. Effects of *Bacillus*-based probiotics on nutrient digestibility of weaned pigs¹

Item	Dietary treatments		SEM	p-value
	CON	PRO		
Apparent ileal digestibility (%)				
DM	84.11	85.18	3.201	0.877
CP	80.78	80.99	3.949	0.872
GE	83.36	83.86	3.233	0.865
Apparent total tract digestibility (%)				
DM	87.48	86.36	1.127	0.500
CP	84.13	87.25	1.155	0.067
GE	87.15	86.22	1.141	0.577

¹Each value presented as the least square mean of 5 replicates.

CON, control diet based on soy-bean meal diet; PRO, control + 0.01% *Bacillus*-based probiotics; DM, dry matter; CP, crude protein; GE, gross energy.

be speculated that improved growth rate of pigs may be a result of a tendency to increase ATTD in CP. Similarly, some previous studies demonstrated that pigs supplemented with *Bacillus*-based probiotics showed improved nutrient digestibility [27,28]. One of the major well-known benefits of *Bacillus* spp. is to eliminate pathogens through nutrient competition in the intestinal mucosa, thereby increasing the balance of host gut microbiota and maximizing nutrient utilization [29,30]. Additionally, *Bacillus* spp. can synthesize various types of extracellular enzymes and proteases [31,32]. Therefore, it is predicted that nutrient digestibility may be increased due to *Bacillus*-based probiotic supplements in feed.

Intestinal morphology

Weaning leads to reduction of VH and VH:CD, impaired intestinal barrier function, and decreased mucin levels [2,3]. Intestinal epithelium cells play an important role in digestion, nutrient absorption, and protection from pathogens and toxins in the gastrointestinal tract [33,34]. Therefore, analyzing the morphology of the intestine is a profitable indicator for assessing the health and function of the intestine [35]. The goblet cells at the intestinal mucus layer produce secretory mucin glycoproteins or membrane-bound mucins, the proteins that form the mucus layer that acts as a line of defense against enteric pathogens and microbial adhesion and invasion [36,37].

In this study, the effects of *Bacillus*-based probiotics on the intestinal morphology of the small intestine (duodenum, jejunum, and ileum) and the number of goblet cells were evaluated (Table 4). *Bacillus*-based probiotics did not affect the intestinal morphology in the duodenum, jejunum, and ileum, but tended to increase the number of goblet cells in the ileum (16.91 vs. 10.83; $p < 0.10$). Similarly, several studies reported that small intestinal morphology was not affected by probiotic products [38–40]. However, other studies have demonstrated a positive effect on intestinal histomorphology in pigs that consume *Bacillus*-based probiotics [37,41,42]. The inconsistent results may be because of different strains or levels of probiotics and the overall health of the pigs.

Immune response

Probiotics are well known to regulate the production of lymphocyte cytokines and have a major effect on the immune system [43]. To investigate the effects of *Bacillus*-based probiotics on the immune systems of piglets, blood parameters were analyzed, including the number of WBC and serum cytokines. As shown in Table 5, the blood parameters were not significantly different between CON and PRO. The TGF- β 1 in the serum tended to be higher in the probiotic group at d 7 (987.28 vs. 590.85 pg/mL; $p < 0.10$) and d 14 (1,095.56 vs. 636.03 pg/mL; $p < 0.10$). The TGF- β 1 is an anti-inflammatory cytokine that is known to maintain intestinal epithelial integrity and support the differentiation of immature dendritic cells [44]. According to a previous study, administration of a probiotic mixture (*Bacillus licheniformis* and *Bacillus subtilis*) increased the expression of mRNA for the gene encoding TGF- β 1 in ileum tissues of weaned pigs [45]. Multi-microbial probiotics also increased TGF- β concentration in the serum of sows [46]. The TNF- α is a pro-inflammatory cytokine that activates the secretion of other cytokines, macrophages, and neutrophils [47]. In addition, it plays an important role in the initial immune response by stimulating immature immune cells to become mature cells and enhancing the antigen-killing ability of these cells [48,49]. Previous studies have reported that feeding a probiotic mixture

Table 4. Effects of *Bacillus*-based probiotics on intestinal morphology of weaned pigs¹⁾

Item	Dietary treatments		SEM	p-value
	CON	PRO		
Duodenum				
VH (μ m)	269.78	286.78	10.30	0.270
CD (μ m)	120.98	122.11	5.04	0.877
VH:CD	2.23	2.35	0.08	0.309
Number of goblet cells (n)	6.10	5.85	0.98	0.858
Jejunum				
VH (μ m)	309.71	279.90	15.76	0.211
CD (μ m)	143.37	131.24	8.54	0.339
VH:CD	2.16	2.13	0.11	0.911
Number of goblet cells (n)	5.07	5.77	0.80	0.548
Ileum				
VH (μ m)	271.84	279.14	18.82	0.789
CD (μ m)	123.62	121.60	6.96	0.842
VH:CD	2.20	2.30	0.16	0.675
Number of goblet cells (n)	10.83	16.91	2.38	0.091

¹⁾Each value presented as the least square mean of 5 replicates.

CON, control diet based on soy-bean meal diet; PRO, control + 0.01% *Bacillus*-based probiotics; VH, villus height; CD, crypt depth.

Table 5. Effects of *Bacillus*-based probiotics on immune responses of weaned pigs¹⁾

Item	Dietary treatments		SEM	p-value
	CON	PRO		
Day 7				
Number of white blood cells ($\times 10^3/\mu\text{L}$)	17.47	17.92	2.277	0.892
CRP (ng/mL)	152.98	282.30	56.45	0.167
Cortisol (ng/mL)	1.21	2.47	0.56	0.172
TGF- β 1 (pg/mL)	590.85	987.28	139.03	0.092
TNF- α (pg/mL)	317.94	366.82	17.98	0.079
IgA (ng/mL)	72.17	66.71	5.89	0.527
IgG (ng/mL)	945.69	943.74	13.29	0.919
IgM (ng/mL)	234.02	199.70	43.52	0.589
Day 14				
Number of white blood cells ($\times 10^3/\mu\text{L}$)	20.32	20.13	2.113	0.952
CRP (ng/mL)	149.88	191.47	43.33	0.529
Cortisol (ng/mL)	1.44	2.02	0.40	0.353
TGF- β 1 (pg/mL)	636.03	1,095.56	153.67	0.089
TNF- α (pg/mL)	341.43	386.42	17.47	0.903
IgA (ng/mL)	88.17	80.53	9.59	0.586
IgG (ng/mL)	972.57	956.67	17.79	0.251
IgM (ng/mL)	161.92	182.53	26.49	0.594

¹⁾Each value presented as the least square mean of 5 replicates.

CON, control diet based on soy-bean meal diet; PRO, control + 0.01% *Bacillus*-based probiotics; CRP, C-reactive protein; TGF- β 1, transforming growth factor- β 1; TNF- α , tumor necrosis factor- α ; IgA, immunoglobulin A; IgG, immunoglobulin G; IgM, immunoglobulin M.

(*Bacillus licheniformis* and *Bacillus subtilis*) increased the mRNA expression of TNF- α , but did not show a significant change in the concentration of TNF- α in the serum of weaned pigs [45]. Similarly, in the current study, the concentration of TNF- α in the serum of the probiotic group tended to be higher on d 7 (366.82 vs. 317.94 pg/mL; $p < 0.10$), but the difference between the two groups disappeared on d 14 (386.42 vs. 341.43 pg/mL; $p = 0.903$). However, there was no significant difference between the two groups in the overall immunity index, which may be due to the minimal immune response of weaned pigs under good hygienic experimental conditions [28]. Consistent with this result, it was previously shown that the treatment with multi-strain probiotics had no effect on immune markers [50].

16S rRNA sequencing data and fecal microbial composition

To investigate the effect of *Bacillus*-based probiotics on microbial composition, taxonomic analysis was performed. A total of 6 fecal samples yielded a total of 233,408 reads in this study, and the number of sequence reads ranged from 35,128 to 49,995 (mean = 38,901; median = 37,070). After quality control and removal of chimeras in OIIME2, the sequences were clustered into 1,799 OTUs based on 97% identity in our fecal samples. The relative abundances of the fecal microbiota of the control and probiotic groups at the phylum and genus levels are shown in Fig. 1. At the phylum level, the bacterial composition of the CON group on the last day of the experiment was predominated by the phyla Firmicutes (74.38%), Bacteroidetes (18.38%), and Spirochaetes (1.18%). Other phyla with relative abundances lower than 1% were Proteobacteria, Spirochaetes, Actinobacteria, and Tenericutes. By comparison, the core fecal microbiota at the phylum level of the probiotic group was Firmicutes (74.76%), Bacteroidetes (15.28%), Proteobacteria (1.63%),




Fig. 1. Effects of *Bacillus*-based probiotics on fecal microbial composition of weaned pigs. Relative abundance of fecal microbial taxa at the phylum level (A) and genus level (B) of both groups. CON, control diet based on soy-bean meal diet; PRO, control + 0.01% *Bacillus*-based probiotics.

and Spirochaetes (1.5%). The low-abundance (< 1%) phyla were Actinobacteria, Chlamydiae, and Cyanobacteria. The abundances of unclassified bacteria from the fecal microbiota of the CON and PRO groups were 4.40% and 5.34%, respectively. A total of 107 genera were classified from all six samples at the genus level. The major genera of the CON group were *Lactobacillus* (21.50%), *Prevotella* (5.98%), *Clostridium* (5.91%), *Ruminococcus* (2.57%), *Barnesiella* (1.59%), and *Roseburia* (1.04%). Furthermore, the predominant taxa at the genus level of the PRO group were *Lactobacillus* (27.77%), *Prevotella* (6.71%), *Clostridium* (4.51%), *Ruminococcus* (1.71%), *Barnesiella* (1.13%), *Treponema* (1.17%), and *Selenomonas* (1.04%). The abundance of unclassified genera in each group was 49.26% and 44.75%, respectively. Intestinal microbes have various functions, such as the production of volatile fatty acids and vitamin K, digestion of cellulose, and regulation of the immune system. Because these functions are closely related to the gut health of pigs, they have been a major issue in recent years. [1,8]. Our findings show that, as in previous studies, Firmicutes and Bacteroidetes in fecal samples accounted for more than 90% of the phyla in both the CON and PRO groups [51,52]. Most unclassified genera belong to the Ruminococcaceae family, which is known to play a beneficial role in gut health by breaking down various polysaccharides and fibers and producing short-chain fatty acids [53,54]. The most dominant genera at the genus level in both the CON and PRO groups were *Prevotella* and *Lactobacillus*. Since weaned pigs generally consume grain-based carbohydrate-rich feed, it is presumed that *Prevotella* and *Lactobacillus*, which are involved in the fermentation of non-starch polysaccharides and the production of short-chain fatty acids, are dominated [55,56].

Fecal microbial diversity analysis

To determine microbial richness and evenness, Chao1, Shannon, and Simpson indices were calculated. The alpha diversity did not differ between the control and probiotic groups (Figs. 2A, B, and C, respectively). To compare the microbial diversity of fecal samples between the two groups, beta diversity was analyzed. PERMANOVA of weighted UniFrac distances showed that there was no remarkable difference between the two groups in microbial diversity (Fig. 2D). The PCoA based on unweighted UniFrac distances also indicated that there were no significant differences between the CON and PRO groups (Fig. 2E). A previous study showed that when *Bacillus subtilis* was fed to weaned pigs, there was no change in the alpha diversity of the small intestinal microbiota, but the microbial composition of the ileum and jejunum contents differed between the control and probiotic groups [57]. In addition, supplementing the feed with *B. subtilis* and/or yeast cultures did not affect the richness or diversity of feces from weaned pigs compared to the control group [58]. It was also reported that feeding weaned pigs higher doses of *B. subtilis* did not change the




Fig. 2. Effects of probiotics on fecal microbial diversity of weaned pigs. Alpha diversity determined by the Chao 1 (A), Shannon (B), and Simpson (C) index. PCoA plot based on weighted UniFrac (D) and unweighted UniFrac (E) distances. CON, control diet based on soy-bean meal diet; PRO, control + 0.01% *Bacillus*-based probiotics; PC, principal component.

alpha diversity compared to the control group but caused a distinct separation between groups [41]. Additional studies are needed for various doses and strains of *Bacillus*-based probiotics that cause clear changes in the intestinal microbiota between groups.

CONCLUSION

The addition of *Bacillus*-based probiotics has beneficial effects on the growth performance including ADG and modulates immune system and gut microbiota of weaned pigs, suggesting that *Bacillus* could be utilized as a functional probiotic in animal feed. However, further study is needed to assess the effects of different *Bacillus* species and doses on the growth performance and intestinal health of weaned pigs and to evaluate the mechanism of action.

REFERENCES

- Lallès JP, Bosi P, Smidt H, Stokes CR. Nutritional management of gut health in pigs around weaning. *Proc Nutr Soc.* 2007;66:260-8. <https://doi.org/10.1017/S0029665107005484>
- Pluske JR, Hampson DJ, Williams IH. Factors influencing the structure and function of the small intestine in the weaned pig: a review. *Livest Prod Sci.* 1997;51:215-36. [https://doi.org/10.1016/S0301-6226\(97\)00057-2](https://doi.org/10.1016/S0301-6226(97)00057-2)
- Kim JC, Hansen CF, Mullan BP, Pluske JR. Nutrition and pathology of weaner pigs: nutritional strategies to support barrier function in the gastrointestinal tract. *Anim Feed Sci Technol.* 2012;173:3-16. <https://doi.org/10.1016/j.anifeedsci.2011.12.022>
- Krause DO, Bhandari SK, House JD, Nyachoti CM. Response of nursery pigs to a synbiotic preparation of starch and an anti-*Escherichia coli* K88 probiotic. *Appl Environ Microbiol.* 2010;76:8192-200. <https://doi.org/10.1128/AEM.01427-10>

5. Dlamini Z, Langa RLS, Aiyegoro OA, Okoh AI. Effects of probiotics on growth performance, blood parameters, and antibody stimulation in piglets. *S Afr J Anim Sci.* 2017;47:766-75. <https://doi.org/10.4314/sajas.v47i6.4>
6. Pettigrew JE. Reduced use of antibiotic growth promoters in diets fed to weanling pigs: dietary tools, part 1. *Anim Biotechnol.* 2006;17:207-15. <https://doi.org/10.1080/10495390600956946>
7. Stein HH, Kil DY. Reduced use of antibiotic growth promoters in diets fed to weanling pigs: dietary tools, part 2. *Anim Biotechnol.* 2006;17:217-31. <https://doi.org/10.1080/10495390600957191>
8. Heo JM, Opapeju FO, Pluske JR, Kim JC, Hampson DJ, Nyachoti CM. Gastrointestinal health and function in weaned pigs: a review of feeding strategies to control post-weaning diarrhoea without using in-feed antimicrobial compounds. *J Anim Physiol Anim Nutr.* 2013;97:207-37. <https://doi.org/10.1111/j.1439-0396.2012.01284.x>
9. Lee JS, Kim TH, Song MH, Oh HJ, Yun W, Lee JH, et al. Effects of microencapsulated organic acids on growth performance, nutrient digestibility, fecal microbial counts, and blood profiles in weaning pigs. *J Anim Sci Technol.* 2021;63:104-13. <https://doi.org/10.5187/jast.2021.e16>
10. de Lange CFM, Pluske J, Gong J, Nyachoti CM. Strategic use of feed ingredients and feed additives to stimulate gut health and development in young pigs. *Livest Sci.* 2010;134:124-34. <https://doi.org/10.1016/j.livsci.2010.06.117>
11. Liao SF, Nyachoti M. Using probiotics to improve swine gut health and nutrient utilization. *Anim Nutr.* 2017;3:331-43. <https://doi.org/10.1016/j.aninu.2017.06.007>
12. Min Y, Choi Y, Choe J, Kim Y, Jeong Y, Kim D, et al. Effects of dietary mixture of protease and probiotics on growth performance, blood constituents, and carcass characteristics of growing-finishing pigs. *J Anim Sci Technol.* 2019;61:272-7. <https://doi.org/10.5187/jast.2019.61.5.272>
13. Kang J, Lee JJ, Cho JH, Choe J, Kyoung H, Kim SH, et al. Effects of dietary inactivated probiotics on growth performance and immune responses of weaned pigs. *J Anim Sci Technol.* 2021;63:520-30. <https://doi.org/10.5187/jast.2021.e44>
14. He Y, Kim K, Kovanda L, Jinno C, Song M, Chase J, et al. *Bacillus subtilis*: a potential growth promoter in weaned pigs in comparison to carbadox. *J Anim Sci.* 2020;98:skaa290. <https://doi.org/10.1093/jas/skaa290>
15. Mingmongkolchai S, Panbangred W. *Bacillus* probiotics: an alternative to antibiotics for livestock production. *J Appl Microbiol.* 2018;124:1334-46. <https://doi.org/10.1111/jam.13690>
16. Hu Y, Dun Y, Li S, Zhao S, Peng N, Liang Y. Effects of *Bacillus subtilis* KN-42 on growth performance, diarrhea and faecal bacterial flora of weaned piglets. *Asian-Australas J Anim Sci.* 2014;27:1131-40. <https://doi.org/10.5713/ajas.2013.13737>
17. NRC [National Research Council]. *Nutrient requirements of swine.* 11th ed. Washington, DC: National Academy Press; 2012.
18. Lee JJ, Kyoung H, Cho JH, Choe J, Kim Y, Liu Y, et al. Dietary yeast cell wall improves growth performance and prevents of diarrhea of weaned pigs by enhancing gut health and anti-inflammatory immune responses. *Animals.* 2021;11:2269. <https://doi.org/10.3390/ani11082269>
19. Park S, Lee JJ, Yang BM, Cho JH, Kim S, Kang J, et al. Dietary protease improves growth performance, nutrient digestibility, and intestinal morphology of weaned pigs. *J Anim Sci Technol.* 2020;62:21-30. <https://doi.org/10.5187/jast.2020.62.1.21>
20. AOAC [Association of Official Analytical Chemists] International. *Official methods of analysis of AOAC International.* 18th ed. Gaithersburg, MD: AOAC International; 2005.
21. Stein HH, Sève B, Fuller MF, Moughan PJ, de Lange CFM. Invited review: amino acid bio-

- availability and digestibility in pig feed ingredients: terminology and application. *J Anim Sci*. 2007;85:172-80. <https://doi.org/10.2527/jas.2005-742>
22. Kyoung H, Lee JJ, Cho JH, Choe J, Kang J, Lee H, et al. Dietary glutamic acid modulates immune responses and gut health of weaned pigs. *Animals*. 2021;11:504. <https://doi.org/10.3390/ani11020504>
 23. Zhang S, Yoo DH, Ao X, Kim IH. Effects of dietary probiotic, liquid feed and nutritional concentration on the growth performance, nutrient digestibility and fecal score of weaning piglets. *Asian-Australas J Anim Sci*. 2020;33:1617-23. <https://doi.org/10.5713/ajas.19.0473>
 24. Wang Y, Cho JH, Chen YJ, Yoo JS, Huang Y, Kim HJ, et al. The effect of probiotic BioPlus 2B® on growth performance, dry matter and nitrogen digestibility and slurry noxious gas emission in growing pigs. *Livest Sci*. 2009;120:35-42. <https://doi.org/10.1016/j.livsci.2008.04.018>
 25. Jørgensen JN, Laguna JS, Millán C, Casabuena O, Gracia MI. Effects of a Bacillus-based probiotic and dietary energy content on the performance and nutrient digestibility of wean to finish pigs. *Anim Feed Sci Technol*. 2016;221:54-61. <https://doi.org/10.1016/j.anifeeds-ci.2016.08.008>
 26. Luise D, Bertocchi M, Motta V, Salvarani C, Bosi P, Luppi A, et al. Bacillus sp. probiotic supplementation diminish the Escherichia coli F4ac infection in susceptible weaned pigs by influencing the intestinal immune response, intestinal microbiota and blood metabolomics. *J Anim Sci Biotechnol*. 2019;10:74. <https://doi.org/10.1186/s40104-019-0380-3>
 27. Lan R, Tran H, Kim I. Effects of probiotic supplementation in different nutrient density diets on growth performance, nutrient digestibility, blood profiles, fecal microflora and noxious gas emission in weaning pig. *J Sci Food Agric*. 2017;97:1335-41. <https://doi.org/10.1002/jsfa.7871>
 28. Wang H, Kim KP, Kim IH. Influence of Bacillus subtilis GCB-13-001 on growth performance, nutrient digestibility, blood characteristics, faecal microbiota and faecal score in weaning pigs. *J Anim Physiol Anim Nutr*. 2019;103:1919-25. <https://doi.org/10.1111/jpn.13199>
 29. Hwang KY, Cho JH, Lee JY, Kang KD, Seong SI, Kim IH. The benefits of using Bacillus as a probiotic. *J Anim Vet Adv*. 2012;11:3457-62. <https://doi.org/10.3923/javaa.2012.3457.3462>
 30. Piewngam P, Zheng Y, Nguyen TH, Dickey SW, Joo HS, Villaruz AE, et al. Pathogen elimination by probiotic Bacillus via signalling interference. *Nature*. 2018;562:532-7. <https://doi.org/10.1038/s41586-018-0616-y>
 31. Priest FG. Extracellular enzyme synthesis in the genus Bacillus. *Bacteriol Rev*. 1977;41:711-53. <https://doi.org/10.1128/br.41.3.711-753.1977>
 32. Das G, Prasad MP. Isolation, purification & mass production of protease enzyme from Bacillus subtilis. *Int Res J Microbiol*. 2010;1:26-31.
 33. Hampson DJ. Alterations in piglet small intestinal structure at weaning. *Res Vet Sci*. 1986;40:32-40. [https://doi.org/10.1016/S0034-5288\(18\)30482-X](https://doi.org/10.1016/S0034-5288(18)30482-X)
 34. Yang H, Xiong X, Wang X, Li T, Yin Y. Effects of weaning on intestinal crypt epithelial cells in piglets. *Sci Rep*. 2016;6:36939. <https://doi.org/10.1038/srep36939>
 35. Cai L, Indrakumar S, Kiarie E, Kim IH. Effects of a multi-strain Bacillus species-based direct-fed microbial on growth performance, nutrient digestibility, blood profile, and gut health in nursery pigs fed corn-soybean meal-based diets. *J Anim Sci*. 2015;93:4336-42. <https://doi.org/10.2527/jas.2015-9056>
 36. Liu Y, Song M, Che TM, Almeida JAS, Lee JJ, Bravo D, et al. Dietary plant extracts alleviate diarrhea and alter immune responses of weaned pigs experimentally infected with a pathogenic Escherichia coli. *J Anim Sci*. 2013;91:5294-306. <https://doi.org/10.2527/jas.2012-6194>
 37. Li Y, Zhang H, Su W, Ying Z, Chen Y, Zhang L, et al. Effects of dietary Bacillus amyloliquifaciens supplementation on growth performance, intestinal morphology, inflammatory re-

- response, and microbiota of intra-uterine growth retarded weanling piglets. *J Anim Sci Biotechnol.* 2018;9:22. <https://doi.org/10.1186/s40104-018-0236-2>
38. Walsh MC, Sholly DM, Hinson RB, Saddoris KL, Sutton AL, Radcliffe JS, et al. Effects of water and diet acidification with and without antibiotics on weanling pig growth and microbial shedding. *J Anim Sci.* 2007;85:1799-808. <https://doi.org/10.2527/jas.2006-049>
 39. Bhandari SK, Xu B, Nyachoti CM, Giesting DW, Krause DO. Evaluation of alternatives to antibiotics using an *Escherichia coli* K88+ model of piglet diarrhea: effects on gut microbial ecology. *J Anim Sci.* 2008;86:836-47. <https://doi.org/10.2527/jas.2006-822>
 40. Choi JY, Shinde PL, Ingale SL, Kim JS, Kim YW, Kim KH, et al. Evaluation of multi-microbe probiotics prepared by submerged liquid or solid substrate fermentation and antibiotics in weaning pigs. *Livest Sci.* 2011;138:144-51. <https://doi.org/10.1016/j.livsci.2010.12.015>
 41. Deng B, Wu J, Li X, Zhang C, Men X, Xu Z. Effects of *Bacillus subtilis* on growth performance, serum parameters, digestive enzyme, intestinal morphology, and colonic microbiota in piglets. *AMB Express.* 2020;10:212. <https://doi.org/10.1186/s13568-020-01150-z>
 42. Zhang W, Zhu YH, Zhou D, Wu Q, Song D, Dicksved J, et al. Oral administration of a select mixture of *Bacillus* probiotics affects the gut microbiota and goblet cell function following *Escherichia coli* challenge in newly weaned pigs of genotype MUC4 that are supposed to be enterotoxigenic *E. coli* F4ab/ac receptor negative. *Appl Environ Microbiol.* 2017;83:e02747-16. <https://doi.org/10.1128/AEM.02747-16>
 43. Klaenhammer TR, Kleerebezem M, Kopp MV, Rescigno M. The impact of probiotics and prebiotics on the immune system. *Nat Rev Immunol.* 2012;12:728-34. <https://doi.org/10.1038/nri3312>
 44. Zeuthen LH, Fink LN, Frokiaer H. Epithelial cells prime the immune response to an array of gut-derived commensals towards a tolerogenic phenotype through distinct actions of thymic stromal lymphopoietin and transforming growth factor- β . *Immunology.* 2008;123:197-208.
 45. Zhou D, Zhu YH, Zhang W, Wang ML, Fan WY, Song D, et al. Oral administration of a select mixture of *Bacillus* probiotics generates Tr1 cells in weaned F4ab/acR- pigs challenged with an F4+ ETEC/VTEC/EPEC strain. *Vet Res.* 2015;46:95. <https://doi.org/10.1186/s13567-015-0223-y>
 46. Laskowska E, Jarosz Ł, Grądziński Z. Effect of multi-microbial probiotic formulation bokashi on pro- and anti-inflammatory cytokines profile in the serum, colostrum and milk of sows, and in a culture of polymorphonuclear cells isolated from colostrum. *Probiotics Antimicrob Proteins.* 2019;11:220-32. <https://doi.org/10.1007/s12602-017-9380-9>
 47. McDermott MF. TNF and TNFR biology in health and disease. *Cell Mol Biol.* 2001;47:619-35.
 48. Zganiacz A, Santosuosso M, Wang J, Yang T, Chen L, Anzulovic M, et al. TNF- α is a critical negative regulator of type 1 immune activation during intracellular bacterial infection. *J Clin Invest.* 2004;113:401-13. <https://doi.org/10.1172/JCI18991>
 49. Izadpanah K, Freyer D, Weber JR, Braun JS. Brain parenchymal TNF- α and IL-1 β induction in experimental pneumococcal meningitis. *J Neuroimmunol.* 2014;276:104-11. <https://doi.org/10.1016/j.jneuroim.2014.08.625>
 50. Lan RX, Lee SI, Kim IH. Effects of multistrain probiotics on growth performance, nutrient digestibility, blood profiles, faecal microbial shedding, faecal score and noxious gas emission in weaning pigs. *J Anim Physiol Anim Nutr.* 2016;100:1130-8. <https://doi.org/10.1111/jpn.12501>
 51. Mach N, Berri M, Estellé J, Levenez F, Lemonnier G, Denis C, et al. Early-life establishment of the swine gut microbiome and impact on host phenotypes. *Environ Microbiol Rep.*

- 2015;7:554-69. <https://doi.org/10.1111/1758-2229.12285>
52. Gaukroger CH, Stewart CJ, Edwards SA, Walshaw J, Adams IP, Kyriazakis I. Changes in faecal microbiota profiles associated with performance and birthweight of piglets. *Front Microbiol.* 2020;11:917. <https://doi.org/10.3389/fmicb.2020.00917>
 53. Fohse JM, Zijlstra RT, Willing BP. The role of gut microbiota in the health and disease of pigs. *Anim Front.* 2016;6:30-6. <https://doi.org/10.2527/af.2016-0031>
 54. Shang Q, Shan X, Cai C, Hao J, Li G, Yu G. Dietary fucoidan modulates the gut microbiota in mice by increasing the abundance of *Lactobacillus* and *Ruminococcaceae*. *Food Funct.* 2016;7:3224-32. <https://doi.org/10.1039/C6FO00309E>
 55. Pieper R, Janczyk P, Zeyner A, Smidt H, Guiard V, Souffrant WB. Ecophysiology of the developing total bacterial and *Lactobacillus* communities in the terminal small intestine of weaning piglets. *Microb Ecol.* 2008;56:474-83. <https://doi.org/10.1007/s00248-008-9366-y>
 56. Kovatcheva-Datchary P, Nilsson A, Akrami R, Lee YS, De Vadder F, Arora T, et al. Dietary fiber-induced improvement in glucose metabolism is associated with increased abundance of *Prevotella*. *Cell Metab.* 2015;22:971-82. <https://doi.org/10.1016/j.cmet.2015.10.001>
 57. Ding H, Zhao X, Ma C, Gao Q, Yin Y, Kong X, et al. Dietary supplementation with *Bacillus subtilis* DSM 32315 alters the intestinal microbiota and metabolites in weaned piglets. *J Appl Microbiol.* 2020;130:217-32. <https://doi.org/10.1111/jam.14767>
 58. Cui K, Iv X, Diao Q, Zhang N. Effects of dietary supplementation with *Bacillus subtilis* and yeast culture on growth performance, nutrient digestibility, serum indices and faeces microbiota of weaned piglets. *J Anim Feed Sci.* 2019;28:328-36. <https://doi.org/10.22358/jafs/114238/2019>