

Invited Mini Review

Deciphering the molecular mechanisms of epitranscriptome regulation in cancer

Seung Hun Han^{1,2,3} & Junho Choe^{1,2,3,*}¹Department of Life Science, College of Natural Sciences, Hanyang University, Seoul 04763, ²Research Institute for Natural Sciences, Hanyang University, Seoul 04763, ³Research Institute for Convergence of Basic Sciences, Hanyang University, Seoul 04763, Korea

Post-transcriptional regulation is an indispensable cellular mechanism of gene expression control that dictates various cellular functions and cell fate decisions. Recently, various chemical RNA modifications, termed the “epitranscriptome,” have been proposed to play crucial roles in the regulation of post-transcriptional gene expression. To date, more than 170 RNA modifications have been identified in almost all types of RNA. As with DNA modification-mediated control of gene expression, regulation of gene expression via RNA modification is also accomplished by three groups of proteins: writers, readers, and erasers. Several emerging studies have revealed that dysregulation in RNA modification is closely associated with tumorigenesis. Notably, the molecular outcomes of specific RNA modifications often have opposite cellular consequences. In this review, we highlight the current progress in the elucidation of the mechanisms of cancer development due to chemical modifications of various RNA species. [BMB Reports 2021; 54(2): 89-97]

INTRODUCTION

Over the past few decades, it has been established that genetic dysregulation underlies various human diseases. For example, genetic errors arising from gene amplification, deletion, mutation, or chromosomal translocation have been associated with numerous cancers (1-3). Meanwhile, increasing evidence suggests that epigenetic modifications of chromatin structure also affect tumor formation and cancer development via abnormal regulation of gene expression (4). Furthermore, the impairment of epigenetic regulation of oncogenes and tumor suppressor genes has been linked to several signaling pathways that lead

*Corresponding author. Tel: +82-2-2220-0966; Fax: +82-2-2298-0319; E-mail: jcho2711@hanyang.ac.kr

<https://doi.org/10.5483/BMBRep.2021.54.2.212>

Received 11 September 2020, Revised 13 October 2020,
Accepted 25 November 2020

Keywords: Cancer, Epitranscriptomics, Gene expression, Post-transcriptional regulation, RNA modification

to cancer development (5-7). Similarly, several recent studies have suggested an important role for RNA modifications, termed the “epitranscriptome”, representing a new layer of post-transcriptional gene regulation (8, 9). Although several studies have investigated signaling pathways and transcriptional regulation in cancer, relatively little is known about the post-transcriptional regulation of cancer. Therefore, a better understanding of the gene regulatory mechanisms controlling tumorigenesis and cancer development will facilitate their therapeutic exploitation (10, 11).

To date, more than 170 chemical RNA modifications have been identified, including N6-methyladenosine (m⁶A), pseudouridine (ψ), N1-methyladenosine (m¹A), 2'-O-methylation (Nm), N5-methylcytosine (m⁵C), and internal N7-methylguanosine (m⁷G) (12, 13). Each of these modifications has been shown to have a preferential modification site within the mRNA.

Similar to DNA methylation in epigenetics, three classes of RNA-binding proteins, broadly classified as writers, readers, and erasers, mediate the regulation of RNA modification (Fig. 1) (14). Writer proteins install the modification, while eraser proteins remove the modification, and reader proteins recognize the modification and regulate the metabolism of the target RNA. The discovery of both writer and eraser proteins indicates that many RNA modifications are likely reversible. Many recent studies have suggested that abnormally regulated RNA modification may lead to tumorigenesis and cancer development (2, 15, 16).

VARIOUS RNA MODIFICATIONS DRIVE TUMORIGENESIS

In this review, we focus on six internal RNA modifications most closely linked to tumorigenesis and discuss the RNA species in which they have been identified, their molecular mechanisms, and evidence of their involvement in cancer (Fig. 1).

N6-methyladenosine (m⁶A)

N6-methyladenosine (m⁶A) is one of the best-characterized RNA modifications whose importance has recently been highlighted in various biological studies. m⁶A modification has been identified in several RNA species, including messenger







Modification	m ⁶ A	ψ	m ¹ A	Nm	m ⁵ C	m ⁷ G	
Structure							
Writer	METTL3/METTL14	PUSs(1-10), DKC1	TRMT6/TRMT61A	FTSJ1, FTSJ2, FBL	NSUNs(1-7), TRDMT1	METTL1/WDR4	
Eraser	FTO, ALKBH5	NR	ALKBH1, ALKBH3	NR	NR	NR	
Reader	YTHDFs(1-3), YTHDCs(1, 2) IGF2BPs(1-3), hnRNPC hnRNPG, eIF3, METTL3	NR	YTHDF2, YTHDF3	NR	Aly/REF, YBX1	NR	
Molecular function	mRNA	Splicing, Export, Translation, Stability	NR	Stability	NR	Export, Stability	NR
	ncRNA	Processing, Stability	Processing, Stability	Processing	Translation	Stability	Processing
RNA species	mRNAs (34-43, 45, 46), miR221/222 miRNA (44), LINC00958 lincRNA (15), XIST lincRNA (47)	18S rRNA (61)	tRNA ^{Met} (67), tRNA-derived small RNAs (70)	rRNAs (80)	HDGF mRNA (89)	let-7e miRNA (93, 97)	
Positive regulation	Gastric Cancer (40), CRC (41), HCC (15), Lung Cancer (34, 35), BLC (42, 44)	BrC (55), PC (56, 59), HCC (57)	GBM (67)	PC (78), BrC (79, 80)	Squamous-Cell Carcinoma (85), BLC (89)	HCC (6)	
Negative regulation	CRC (45, 47), BLC (46)	BrC (80), HCC (61)	HCC, PC (70)	NR	NR	Lung Cancer (93), Colon Cancer (97)	

Fig. 1. Role of RNA modification in cancer. Six representative cancer-associated RNA modifications are listed with their chemical structures, molecular functions, target RNA species, and binding proteins. Positive or negative regulation indicates the correlation between each RNA modification and the corresponding cancer progression. Blue boxes indicate general information about each RNA modification regardless of cancer association, while the yellow boxes represent information about RNA modification dependent-cancer regulation. NR, not reported.

RNA (mRNA), transfer RNA (tRNA), ribosomal RNA (rRNA), non-coding RNA (ncRNA), and viral RNA genomes (12, 17). It has been reported that m⁶A modification is the most prevalent mRNA modification, and approximately 25% of eukaryotic mRNAs harbor at least one m⁶A modified base (8, 18). Although m⁶A modification sites vary among different mRNA species and tissues, these sites have been shown to be generally located near translation stop codons in the 3' untranslated region (UTR) of mRNA (8, 18, 19). However, a study showed that m⁶A modifications in ncRNA are dispersed across the gene body without any apparent preferred location (20). Nevertheless, both coding and ncRNAs share a common set of m⁶A writer protein complexes. Methyltransferase-like protein 3 (METTL3, also known as MT-A70), which catalyzes m⁶A modification, forms a catalytic core complex with METTL14, which recognizes the DRACH motif (D = A, G, or U; R = G or A; and H = A, C, or U) (18, 21). Although several different components of the complex have been discovered, most reports consistently suggested that additional proteins, which lack methyltransferase activity, are required for proper m⁶A modification (22-24). m⁶A modification is known to be reversed by the action of m⁶A eraser proteins. Unlike m⁶A writer protein complexes, only two m⁶A eraser proteins, fat mass and obesity-associated protein (FTO) and α-ketoglutarate-dependent dioxygenase alk B homolog 5 (ALKBH5), have been identified to date (25, 26). These two eraser proteins do not form a

complex but are independently responsible for demethylation of m⁶A (25, 26). While several previous studies have reported that FTO depletion leads to global demethylation of m⁶A, more recent data indicate that FTO preferentially demethylates 2'-O-dimethyladenosine (m⁶Am) or N1-methyladenosine (m¹A) in tRNA (27). In contrast, ALKBH5 is considered a better m⁶A demethylase candidate as it recognizes the m⁶A demethylation consensus sequence (26). The m⁶A reader proteins have been studied more extensively. m⁶A modification is recognized by proteins belonging to the YT521-B homology (YTH) domain family (YTHDF1, YTHDF2, and YTHDF3), YTH domain-containing proteins (YTHDC1 and YTHDC2), heterogeneous nuclear ribonucleoproteins (hnRNPC and hnRNPG), and insulin-like growth factor 2 mRNA-binding proteins (IGF2BP1, IGF2BP2, and IGF2BP3), which regulate almost all the steps of mRNA metabolism, including splicing, export, translation, and stability (28-32). Interestingly, eukaryotic translation initiation factor 3 (eIF3) has also been shown to function as an m⁶A reader protein by directly binding m⁶A modification sites in the 5' UTR of mRNA, resulting in the recruitment of the 43S complex and initiation of translation independently of the cap-binding protein, eIF4E (33-35).

Several studies have investigated the association between m⁶A modification and cancer. These studies have revealed that the levels of m⁶A mRNA modification in cancer cells are generally elevated and closely correlated with the develop-

ment of several cancers (36-39). In colorectal cancer (CRC) and gastric cancer, for example, a high degree of m⁶A modification is associated with mRNA stability (40, 41). m⁶A reader proteins, IGF2BP1, IGF2BP2, and IGF2BP3, recognize m⁶A modifications in oncogene mRNAs, preventing mRNA degradation and ultimately promoting cancer development (40, 41). In bladder cancer (BLC) and lung cancer, m⁶A modification increases the translation efficiency of oncogenes without affecting mRNA abundance (34, 35, 42). Integrin alpha-6 (*ITGA6*) mRNA exhibits a high level of m⁶A modification at the 3' UTR of mRNA, which is recognized by the reader proteins, YTHDF1 and YTHDF3 in BLC, promoting the translation of *ITGA6* mRNA (42). In lung cancer, METTL3 has been shown to function as both an m⁶A writer and a reader protein. METTL3 recognizes m⁶A modifications at the 3' UTR of mRNAs near the translation stop codon and then interacts directly with eIF3 subunit h (eIF3h) at the 5' end, promoting translation via ribosome recycling by forming an mRNA loop. Consequently, METTL3-dependent translation enhancement of a large subset of mRNAs triggers tumorigenesis in lung cancer (34, 35). Many studies have revealed a correlation between high levels of m⁶A mRNA modifications and development of several cancers, including acute myeloid leukemia (AML), lung cancer, breast cancer (BrC), BLC, and glioblastoma (GBM) (36, 38, 39, 43). However, the m⁶A reader proteins involved in specific cancers and their molecular mechanisms are yet to be elucidated.

Cancer development has also been reported to be influenced by m⁶A modification of ncRNA (15, 44). MicroRNA (miRNA) m⁶A modification can alter the abundance of miRNA in cells, which in turn regulates the stability of the target mRNA associated with tumorigenesis in BLC and hepatocellular carcinoma (HCC) (15, 44). Specifically, METTL3 has been shown to interact with the microprocessor protein, DGCR8, affecting primary-miRNA (pri-miRNA) processing in BLC (44). A study showed that the knockdown of METTL3 in BLC induces the accumulation of *pri-miR221/222*, whereas the overexpression of METTL3 increases the level of mature *miR221/222* (44). The study further showed that METTL3 accelerates the maturation of *pri-miR221/222*, resulting in the suppression of the transcription of a tumor suppressor gene, phosphatase and tensin homolog (*PTEN*) (44). In addition, m⁶A modification of long intergenic non-coding RNA (lincRNA) has been implicated in cancer development. In HCC, METTL3-mediated m⁶A modification increases *LINC00958* stability (15). Consequently, accumulation of *LINC00958* acts as an miRNA sponge by binding and inhibiting *miR-3619-5p*. The loss of *miR-3619-5p* enhances the expression of its target oncogene, hepatoma-derived growth factor (*HDFG*) mRNA, which in turn promotes lipogenesis, cell proliferation, and migration in HCC cells (15).

However, it has been reported that a high level of m⁶A modifications may also be inhibitory in the same types of cancer, depending on the target mRNAs. Recently, low levels

of m⁶A modifications were identified in CRC and BLC (45-47). Mechanistically, the increase in m⁶A modification induced by the overexpression of METTL14 is thought to lead to rapid YTHDF2-mediated mRNA degradation of the essential developmental transcription factor, SRY-related high-mobility-group box 4 (*SOX4*). Loss of *SOX4* expression inhibits CRC malignancy via *SOX4*-mediated epithelial-mesenchymal transition (EMT) and PI3K/Akt signal transduction (45). In BLC, METTL14 expression is low, resulting in low m⁶A modification levels. This, in turn, leads to increased mRNA stability of *Notch1*, which has been shown to play a crucial role in cell proliferation, self-renewal, and metastasis of BLC (46).

The development of several cancers is also influenced by m⁶A modification of ncRNA. In CRC, an oncogenic long non-coding RNA (lncRNA), the X inactive-specific transcript (*XIST*), has been shown to be highly methylated and degraded by YTHDF2 (47). Thus, m⁶A-modification-mediated loss of *XIST* expression leads to the inhibition of cell proliferation, metastasis, and tumorigenesis (47).

Pseudouridine (ψ)

Pseudouridine (also known as 5-ribosyluracil or ψ) was first discovered in the early 1950s (48). Initially identified in tRNA and rRNA, pseudouridine has also been found in mRNA, lncRNA, and small nuclear RNA (snRNA) (49, 50). Pseudouridine is the most abundant RNA modification occurring in tRNAs and rRNAs (50). In tRNAs, pseudouridine is generally localized to the anticodon stem-loop in the D stem, and to the nucleotide position 55 in the T loop, and thus contributes to the stabilization of the tertiary structure of tRNA (51). Pseudouridines are generated post-transcriptionally via C5-ribosyl isomerization of one or a few target uridines catalyzed by pseudouridine synthases (PUSs) (52). PUSs employ two mechanisms of pseudouridine modification: guide RNA-dependent H/ACA box small nucleolar RNA (snoRNA) and guide RNA-independent pseudouridylation. In guide RNA-dependent pseudouridylation, the H/ACA box snoRNA forms a complex with dyskerin pseudouridine synthase 1 (DKC1), which recognizes specific sequences for pseudouridylation on target RNAs, including rRNA, snRNA, and snoRNA (53). In contrast, in guide RNA-independent pseudouridylation, modification of the target RNA is directly catalyzed by stand-alone PUSs (54). PUS enzymes are classified into six families: TruA, TruB, TruD, RluA, RsuA, and Pus10 (54). PUS1, which belongs to the TruA family, was originally thought to pseudouridylate tRNA alone; however, recent studies have identified that PUS1 also pseudouridylates rRNA, snRNA, and mRNA (49, 50). In addition, PUS4 and PUS7 were found to target mRNAs for pseudouridylation in HEK293T cells (50).

Unlike other known RNA modifications, neither reader nor eraser proteins for pseudouridine have been identified to date. Moreover, the functional role of pseudouridylation in mRNA remains unclear.

Upregulation of pseudouridine has been shown to be asso-

ciated with the progression of various cancers, including prostate cancer (PC), BrC, and HCC (55-58). The nucleolar protein DKC1 plays an important role in two separate cell proliferation pathways: the pseudouridylation of rRNAs, which is necessary for their processing, and the stabilization of the telomerase RNA component that is necessary for telomerase activity (55). Similarly, DKC1 expression is generally upregulated in PC (56, 59). The knockdown of DKC1 by siRNA has been shown to inhibit the proliferation of the e22Rv1, LNCaP, PC3, and Du145 in PC cell lines; however, the knockdown had no significant effects on apoptosis or senescence (56). Moreover, HCC patients with high DKC1 expression have been found to exhibit shorter overall survival rates when compared to those with low DKC1 expression (57). Moreover, elevated DKC1 expression has been shown to positively correlate with MYC oncogene expression, which triggers the expression of target genes to induce cell proliferation and cell survival. In addition, DKC1 expression was shown to induce MKI67 expression, which is considered a marker for cell proliferation (57, 58).

However, pseudouridine modification has also been shown to exert negative effects on the development of several types of cancer. Downregulated pseudouridine has been associated with BrC and HCC development (60, 61). The impairment of DKC1 protein can lead to the inactivation of p53, a well-known anti-tumor development factor that induces cell cycle arrest or apoptosis, due to abnormal p53 mRNA translation (60). However, low expression of a snoRNA, SNORA24, has been associated with poor survival of patients with steatohepatic HCC (61). As SNORA24 is known to mediate the pseudouridylation of 18S rRNA, the depletion of SNORA24 increases the rate of translational miscoding and stop codon read-through frequencies in human HCC (61).

N1-methyladenosine (m¹A)

N1-methyladenosine (m¹A) was first identified in yeast tRNA^{Phe} and later found to be widespread in tRNAs (62, 63). In human cells, the methyltransferases TRMT10C, TRMT61B, and TRMT6/61A catalyze m¹A at positions 9 and 58 of mitochondrial and cytoplasmic tRNA, respectively (63). m¹A modification of tRNA plays a pivotal role in maintaining its proper structure, stability, and function (64-67). Although m¹A modification is abundant in tRNA, relatively low levels of modification, ranging from 0.015% to 0.054%, have been found in mammalian mRNA (68). The majority of m¹A modification sites in mRNA are located in the 5' UTR near the translation start codon (68). TRMT6/61A has been found to localize in the cytoplasm, and contributes to m¹A modification in select mRNAs that harbor a tRNA-like motif, GUUCRA (63). As the TRMT6/61A complex seems to play a minor role in m¹A mRNA modification, other m¹A methyltransferases likely remain to be discovered (63).

Similar to m⁶A modification, m¹A modification is reversible. Demethylation of m¹A modification is mediated by ALKBH1 and ALKBH3 (65, 66). Interestingly, the m⁶A reader proteins, YTHDF2 and YTHDF3, have also been implicated in the

recognition of m¹A mRNA modifications (16, 69). Reminiscent of m⁶A modification, m¹A-modified mRNA undergoes rapid degradation upon binding with YTHDF2 or YTHDF3 (16, 69).

To date, several studies have reported a positive correlation between m¹A tRNA modification and cancer development. Elevated expression levels of both TRMT6/61A and initiator methionine tRNA (tRNA^{iMet}) have been detected in highly aggressive GBM compared with grade II/III gliomas (67). Depletion of the TRMT6/61A complex suppresses proliferation and promotes cell death in C6 glioma cells, which can be rescued in part by tRNA^{iMet} overexpression (67). Conversely, the ectopic expression of TRMT6/61A has been shown to upregulate the translation of oncogenic mRNAs, leading to increased colonization of C6 glioma cells (67). Similarly, ALKBH1-mediated demethylation of m¹A modified tRNA attenuates translation initiation and elongation in HeLa cells, thereby reducing cell proliferation (66).

Intriguingly, another m¹A demethylase, ALKBH3, has been shown to induce an opposite effect to that of ALKBH1 in m¹A tRNA modification (70). ALKBH3-mediated m¹A tRNA demethylation increases the susceptibility of tRNA to angiogenin cleavage and generates tRNA-derived small RNAs in various cancer cells during cancer cell proliferation, migration, and invasion (70).

2'-O-methylation (Nm)

A rather unique RNA modification is mediated via 2'-O-methylation (also known as Nm or 2'-O-me). Nm can occur in any nucleotide via the addition of a methyl group to the 2' hydroxyl of the ribose moiety (71). Nm is an abundant RNA modification that occurs in multiple RNA species, including tRNA, rRNA, snRNA, and mRNA (71-73). Nm modification sites in mRNA are mostly located within the coding sequence (CDS) regions (73). Interestingly, Nm modification is also found in approximately 16.2% of introns, suggesting that Nm modification occurs in the nucleus prior to splicing events (73). In higher eukaryotes, Nm modification occurs specifically in the first nucleotide [N1] and/or second nucleotide [N2] next to the m⁷G-cap of mRNA (74). While N1 methylation occurs in almost all mRNA molecules, N2 methylation accounts for about half of the capped mRNA. Additional m⁶A methylation following Nm leads to m⁶Am formation. Two conserved Nm writer proteins have been discovered for N1 and N2 methylation. FTSJ2 (also known as hMTr1) is responsible for the methylation of N1, whereas FTSJ1 (also known as hMTr2) contributes to N2 methylation (75, 76). FTSJ2 also methylates mitochondrial 23S rRNA (77). Another Nm writer protein, fibrillarlin (FBL), is an essential nucleolar protein that catalyzes the Nm of rRNA by interacting with a C/D box family member of U3 snoRNA that mediates the processing of precursor rRNA (74). Given that more than 70% of all Nm occurs in the CDS regions of mRNA and that known Nm writers have not been associated with CDS methylation, further studies are needed to identify additional Nm methyltransferases. Nm readers and

erasers also remain unknown.

Many studies suggest that FBL expression is abnormally high in various cancers, including PC and BrC (78-80). One report suggested that a p53 mutation fails to suppress transcription of the *FBL* gene, resulting in the overexpression of FBL (80). Another report suggested that the MYC protein directly regulates the transcription of *FBL* by binding to the promoter region of this gene (78). The elevated level of FBL expression has been shown to be associated with a higher transcriptional activity of RNA polymerase I, thus increasing rRNA production (74, 80). As increased rRNA biogenesis has been observed in various cancer cells, it is not surprising that many cancers appear to be related to FBL-mediated rRNA production. Indeed, depletion of FBL has been shown to significantly inhibit tumorigenesis in PC and BrC (79, 80). In contrast, increased FBL expression not only induces high levels of rRNA synthesis, but also promotes rRNA Nm that regulates ribosomal quality (74, 80). Altered rRNA Nm patterns have been shown to be associated with a decrease in translational fidelity (*i.e.*, increased nonsense suppression, frame-shifts, and mis-incorporation) and increased internal ribosome entry site (IRES)-dependent translation initiation of key cancer genes (74, 80). In case of another Nm methyltransferase, only one report indicated that the increased expression of FTSJ2 was positively correlated with cell proliferation in lung cancer cell lines (77). To date, a negative correlation between Nm and cancer has not been reported. Furthermore, since reversible Nm has not been reported, it may be important for cells to maintain Nm levels below the cancer initiation threshold.

N5-methylcytosine (m⁵C)

N5-methylcytosine was originally found in ncRNA, including tRNA and rRNA, and subsequently in most RNA species (81, 82). More than 10,000 m⁵C modification sites have been discovered in human mRNA and ncRNA (81). m⁵C modifications have been found to be enriched in the 5' UTRs and 3' UTRs of mRNA (81). Recently, a study reported that m⁵C modifications are enriched in CG-rich regions and immediately downstream of translation initiation regions of mRNA in multiple mice tissues (82). In rRNA, m⁵C modification sites have been shown to be fairly conserved, and both the small and large subunits of rRNA in bacteria and eukaryotes harbor m⁵C methylation sites at similar positions (83). Although several m⁵C locations have been identified in human 28S rRNA, the functional roles of these sites remain unclear (83). Several m⁵C modification writer proteins have been identified, including members of the NOL1/NOP2/Sun (NSUN)-domain-containing family (from NSUN1 to NSUN7) and DNA methyltransferase 2 (DNMT2) families (84). Initially, NSUN2 was thought to be responsible for m⁵C modification of tRNA, but was later found to also methylate many RNA species, including tRNA, mRNA, and miRNA (81, 82, 85). tRNA methyltransferase 1 (TRDMT1), a member of the DNMT2 family, is primarily responsible for the m⁵C modification of tRNA and mRNA (86). tRNA m⁵C

methylation mediated by either TRDMT1 or NSUN2 increases tRNA stability and protein synthesis (87). Other members of the NSUN family have been associated with m⁵C modification in enhancer RNA, rRNA, mitochondrial tRNA (mt-tRNA), or mt-rRNA (88). In the two protein families, NSUN2 and DNMT2 are known to be associated with malignant cancer. Although an m⁵C eraser protein has not been identified, Aly/REF, an mRNA export adapter protein complex, has been found to act as a novel m⁵C reader protein that methylates mRNA in CG-rich sequences and in translation initiation regions (82). A recent study identified Y-box-binding protein 1 (YBX1) as a reader protein that recognizes m⁵C modification and maintains the stability of the target mRNA (89).

A few studies have investigated the effects of m⁵C on tumorigenesis. It has been identified that NSUN2 protein is a downstream target of MYC that methylates RNA polymerase III transcripts (85). Elevated expression of NSUN2 has been shown to mediate MYC-induced cell proliferation and growth in squamous cell carcinoma and BLC (85, 89). Moreover, *HDGF* mRNA has been reported to undergo m⁵C modification in the 3' UTR and is stabilized by YBX1, increasing HDGF protein synthesis (89). This finding was consistent with those of another study showing that reduced survival of BLC patients was correlated with increased expression of NSUN2 and HDGF (89). Another m⁵C methyltransferase, TRDMT1, is also correlated with cell proliferation and migration and gene ontology (GO) analyses have shown that the depletion of TRDMT1 leads to altered expression of genes associated with the regulation of the cell cycle, RNA transport, and RNA degradation (90).

While few studies have reported a correlation between m⁵C modification and cancer, it will be informative to consider the effects of m⁵C on the regulation of translation, given that m⁵C modifications occur mainly in 5' or 3' UTRs.

Internal N7-methylguanosine (m⁷G)

N7-methylguanosine (m⁷G) is the best-known modification in the form of mRNA 5' cap (91). Recently, an internal m⁷G modification was also identified in rRNAs, tRNAs, miRNAs, and mRNAs (92-94). m⁷G modification is found at the nucleotide position 46 in several tRNA variable loops known to stabilize the tRNA tertiary structure (95, 96). In mammalian cells, the internal m⁷G/G ratio of mRNA ranges from 0.02% to 0.05% (94). Internal m⁷G modification has been found to be preferentially enriched in either AG-rich or GA-rich regions in mRNA (94, 95) and is mediated by METTL1 and its co-factor, WD repeat domain 4 (WDR4) in mRNA, miRNA, and tRNA (6, 96). Specifically, the knockdown of METTL1 has been shown to lead to a global decrease in internal m⁷G modification by approximately 54% and 61% in HeLa cells and HepG2 cells, respectively (94). To date, no readers or erasers for m⁷G modification have been identified.

A positive association between internal m⁷G and cancer has yet to be identified. However, internal m⁷G modification of primary miRNA transcripts has been shown to have negative

effects on lung cancer and colon cancer development. Specifically, the *pri-let-7e* transcript has been found to be targeted by METTL1 during internal m⁷G modification and generates a mature form of *let-7e* via efficient processing (93, 97). The increased expression of mature *let-7e* downregulates its target, the high-mobility group AT-hook 2 (*HMG A2*) mRNA, which is known to drive cancer metastasis (93, 97). The association with m⁷G modification is unclear; however, the m⁷G writer protein, METTL1, itself inhibits the PTEN signaling pathway in HCC to promote cell proliferation and migration, resulting in tumor growth and greater tumor vascular invasion (6).

DISCUSSION

Given the plethora of signaling pathways converging upon gene expression regulatory pathways to satisfy the increased anabolic demands in cancer, a better understanding of the gene regulatory mechanisms controlling tumorigenesis will facilitate their therapeutic exploitation (5, 6, 11). The field of epitranscriptomics has attracted the attention of various biological investigators in recent years; however, the molecular players and mechanisms underlying epitranscriptome regulation remain to be elucidated. In particular, several studies on the relationship between RNA modification and cancer have been published, many of which either lack data regarding the detailed mechanisms involved or often report contrasting results. For instance, pseudouridine, m¹A, Nm, and internal m⁷G modifications have been found in mRNAs across the gene body; however, their roles in mRNA metabolism and their effects on cancer progression remain unknown (49, 65, 94, 95). Instead, most published studies have merely investigated the effects of the enzymes catalyzing the modifications on cancers, without confirming the effects of RNA modification. With the exception of m⁶A and m¹A, the reversibility of RNA modifications and their specific reader proteins also remain unclear. Moreover, the molecular functions and cellular consequences of m⁶A or pseudouridine modification often differ across studies, depending on the degree of methylation in the specific target RNAs (Fig. 1).

In addition to the extensive interest in the roles of RNA modification in cancer, the development of potential drugs to treat cancer by modulating RNA modification has also attracted attention (98). However, there are currently no inhibitors or antagonists targeting writer and reader proteins, or the RNA modifications discussed in this review (98). Instead, several inhibitors of demethylases have been suggested. The ALKBH family and FTO share a common structure required for the binding of Fe²⁺ as a co-factor and 2-oxoglutarate (2OG) as a co-substrate (98). Therefore, most of the known compounds, including 2OG competitors such as N-oxalylglycine and its cell-penetrating derivative dimethyl oxalylglycine, succinate, fumarate, and 2-hydroxyglutarate, or metal chelators such as hydroxamic acids and flavonoids, were designed to target either Fe²⁺ or 2OG binding sites (98, 99). However, these

compounds are still far from being used in anticancer drugs because they nonspecifically inhibit various demethylases. Some specific demethylase inhibitors have also been discovered, such as the ALKBH3 inhibitor, 1-(5-methyl-1H-benzimidazol-2-yl)-4-benzyl-3-methyl-1H-pyrazol-5-ol (HUHS015), and FTO inhibitor, Rhein. However, the efficacy of these drugs remains doubtful because of their nonspecificity for target RNAs (98, 99). More recently, clustered regularly interspaced short palindromic repeats (CRISPR) based RNA-editing technology has been suggested for modulating target mRNA specific modifications (100). The CRISPR-associated nuclease Cas13 has been shown to cleave the targeted single-stranded RNA (100). A catalytically inactive mutant of Cas13 (dCas13) fused with m⁶A methyltransferases METTL3 or METTL14 has been found to bind to the target mRNA specifically directed by the guide RNA, without the cleavage of the mRNA (100). Targeting of these fusion proteins has been shown to specifically methylate adenosine within a small range of target sites, regardless of the m⁶A consensus sequence (100). Thus, CRISPR-based approaches can be applied not only in the modulation of m⁶A modifications, but also to other types of RNA modifications. Given the abnormal up/downregulation of modified RNA transcripts and their regulatory proteins in cancer, the development of this technology will shed more light on the feasibility of controlling the modification of RNA targets for cancer treatment. Taken together, the analysis presented in this review highlights the need to further elucidate the mechanisms of RNA modification-mediated regulation of gene expression to improve cancer therapy.

ACKNOWLEDGEMENTS

This work was supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government (Ministry of Science and ICT) (NRF-2020R1C1C1009842 and NRF-2020R1A4A1018398).

CONFLICTS OF INTEREST

The authors have no conflicting interests.

REFERENCES

1. Zhou S, Treloar AE and Lupien M (2016) Emergence of the noncoding cancer genome: a target of genetic and epigenetic alterations. *Cancer Discovery* 6, 1215-1229
2. Chiba T, Marusawa H and Ushijima T (2012) Inflammation-associated cancer development in digestive organs: mechanisms and roles for genetic and epigenetic modulation. *Gastroenterology* 143, 550-563
3. Yu K, Xiang L, Li S, Wang S, Chen C and Mu H (2019) HIF1 α promotes prostate cancer progression by increasing ATG5 expression. *Anim Cells Syst (Seoul)* 23, 326-334
4. Heard E and Martienssen RA (2014) Transgenerational

- epigenetic inheritance: myths and mechanisms. *Cell* 157, 95-109
5. Zhao Y, Zhao Q, Kaboli PJ et al (2019) m1A regulated genes modulate PI3K/AKT/mTOR and ErbB pathways in gastrointestinal cancer. *Transl Oncol* 12, 1323-1333
 6. Tian QH, Zhang MF, Zeng JS et al (2019) METTL1 overexpression is correlated with poor prognosis and promotes hepatocellular carcinoma via PTEN. *J Mol Med* 97, 1535-1545
 7. Yu FX, Zhao B and Guan KL (2015) Hippo pathway in organ size control, tissue homeostasis, and cancer. *Cell* 163, 811-828
 8. Meyer KD, Saletore Y, Zumbo P, Elemento O, Mason CE and Jaffrey SR (2012) Comprehensive analysis of mRNA methylation reveals enrichment in 3' UTRs and near stop codons. *Cell* 149, 1635-1646
 9. Helm M and Motorin Y (2017) Detecting RNA modifications in the epitranscriptome: predict and validate. *Nat Rev Genet* 18, 275-291
 10. Zhao BS, Roundtree IA and He C (2018) Publisher correction: Post-transcriptional gene regulation by mRNA modifications. *Nat Rev Mol Cell Biol* 19, 808
 11. Fu Y, Dominissini D, Rechavi G and He C (2014) Gene expression regulation mediated through reversible m(6)A RNA methylation. *Nat Rev Genet* 15, 293-306
 12. Boccaletto P, Machnicka MA, Purta E et al (2018) MODOMICS: a database of RNA modification pathways. 2017 update. *Nucleic Acids Res* 46, D303-D307
 13. Zhang ZJ, Park E, Lin L and Xing Y (2018) A panoramic view of RNA modifications: exploring new frontiers. *Genome Biol* 19, 11
 14. Shi H, Wei J and He C (2019) Where, when, and how: context-dependent functions of RNA methylation writers, readers, and erasers. *Mol Cell* 74, 640-650
 15. Zuo XL, Chen ZQ, Gao W et al (2020) m6A-mediated upregulation of LINC00958 increases lipogenesis and acts as a nanotherapeutic target in hepatocellular carcinoma. *J Hematol Oncol* 13, 5
 16. Zheng QL, Gan HL, Yang FL et al (2020) Cytoplasmic m(1)A reader YTHDF3 inhibits trophoblast invasion by downregulation of m(1)A-methylated IGF1R. *Cell Discov* 6, 12
 17. Nachtergaele S and He C (2018) Chemical modifications in the life of an mRNA transcript. *Annu Rev Genet* 52, 349-372
 18. Dominissini D, Moshitch-Moshkovitz S, Schwartz S et al (2012) Topology of the human and mouse m6A RNA methylomes revealed by m6A-seq. *Nature* 485, 201-206
 19. Linder B, Grozhik AV, Olarerin-George AO, Meydan C, Mason CE and Jaffrey SR (2015) Single-nucleotide-resolution mapping of m6A and m6Am throughout the transcriptome. *Nat Methods* 12, 767-772
 20. Xiao S, Cao S, Huang Q et al (2019) The RNA N(6)-methyladenosine modification landscape of human fetal tissues. *Nat Cell Biol* 21, 651-661
 21. Wang P, Doxtader KA and Nam Y (2016) Structural basis for cooperative function of Mettl3 and Mettl14 methyltransferases. *Mol Cell* 63, 306-317
 22. Wen J, Lv R, Ma H et al (2018) Zc3h13 regulates nuclear RNA m(6)A methylation and mouse embryonic stem cell self-renewal. *Mol Cell* 69, 1028-1038 e1026
 23. Ping XL, Sun BF, Wang L et al (2014) Mammalian WTAP is a regulatory subunit of the RNA N6-methyladenosine methyltransferase. *Cell Res* 24, 177-189
 24. Patil DP, Chen CK, Pickering BF et al (2016) m(6)A RNA methylation promotes XIIST-mediated transcriptional repression. *Nature* 537, 369-373
 25. Jia G, Fu Y, Zhao X et al (2011) N6-methyladenosine in nuclear RNA is a major substrate of the obesity-associated FTO. *Nat Chem Biol* 7, 885-887
 26. Zheng G, Dahl JA, Niu Y et al (2013) ALKBH5 is a mammalian RNA demethylase that impacts RNA metabolism and mouse fertility. *Mol Cell* 49, 18-29
 27. Wei J, Liu F, Lu Z et al (2018) Differential m(6)A, m(6)Am, and m(1)A demethylation mediated by FTO in the cell nucleus and cytoplasm. *Mol Cell* 71, 973-985 e975
 28. Wang Y, Li Y, Toth JI, Petroski MD, Zhang Z and Zhao JC (2014) N6-methyladenosine modification destabilizes developmental regulators in embryonic stem cells. *Nat Cell Biol* 16, 191-198
 29. Huang H, Weng H, Sun W et al (2018) Recognition of RNA N(6)-methyladenosine by IGF2BP proteins enhances mRNA stability and translation. *Nat Cell Biol* 20, 285-295
 30. Wang X, Zhao BS, Roundtree IA et al (2015) N(6)-methyladenosine modulates messenger RNA translation efficiency. *Cell* 161, 1388-1399
 31. Shi H, Wang X, Lu Z et al (2017) YTHDF3 facilitates translation and decay of N(6)-methyladenosine-modified RNA. *Cell Res* 27, 315-328
 32. Liu N, Dai Q, Zheng G, He C, Parisien M and Pan T (2015) N(6)-methyladenosine-dependent RNA structural switches regulate RNA-protein interactions. *Nature* 518, 560-564
 33. Meyer KD, Patil DP, Zhou J et al (2015) 5' UTR m(6)A promotes cap-independent translation. *Cell* 163, 999-1010
 34. Choe J, Lin S, Zhang W et al (2018) mRNA circularization by METTL3-eIF3h enhances translation and promotes oncogenesis. *Nature* 561, 556-560
 35. Lin S, Choe J, Du P, Triboulet R and Gregory RI (2016) The m(6)A methyltransferase METTL3 promotes translation in human cancer cells. *Mol Cell* 62, 335-345
 36. Han SH and Choe J (2020) Diverse molecular functions of m(6)A mRNA modification in cancer. *Exp Mol Med* 52, 738-749
 37. Fazi F and Fatica A (2019) Interplay between N(6)-methyladenosine (m(6)A) and non-coding RNAs in cell development and cancer. *Front Cell Dev Biol* 7, 116
 38. Sun T, Wu R and Ming L (2019) The role of m6A RNA methylation in cancer. *Biomed Pharmacother* 112, 108613
 39. Chen XY, Zhang J and Zhu JS (2019) The role of m(6)A RNA methylation in human cancer. *Mol Cancer* 18, 103
 40. Wang Q, Chen C, Ding Q et al (2020) METTL3-mediated m(6)A modification of HDGF mRNA promotes gastric cancer progression and has prognostic significance. *Gut* 69, 1193-1205
 41. Li T, Hu PS, Zuo Z et al (2019) METTL3 facilitates tumor progression via an m(6)A-IGF2BP2-dependent mechanism in colorectal carcinoma. *Mol Cancer* 18, 112
 42. Jin H, Ying X, Que B et al (2019) N(6)-methyladenosine

- modification of ITGA6 mRNA promotes the development and progression of bladder cancer. *EBioMedicine* 47, 195-207
43. Barbieri I, Tzelepis K, Pandolfini L et al (2017) Promoter-bound METTL3 maintains myeloid leukaemia by m(6)A-dependent translation control. *Nature* 552, 126-131
 44. Han J, Wang JZ, Yang X et al (2019) METTL3 promote tumor proliferation of bladder cancer by accelerating pri-miR221/222 maturation in m6A-dependent manner. *Mol Cancer* 18, 110
 45. Chen XX, Xu M, Xu XN et al (2020) METTL14-mediated N6-methyladenosine modification of SOX4 mRNA inhibits tumor metastasis in colorectal cancer. *Mol Cancer* 19, 106
 46. Gu CH, Wang ZY, Zhou NC et al (2019) Mettl14 inhibits bladder TIC self-renewal and bladder tumorigenesis through N6-methyladenosine of Notch1. *Mol Cancer* 18, 168
 47. Yang X, Zhang S, He C et al (2020) METTL14 suppresses proliferation and metastasis of colorectal cancer by down-regulating oncogenic long non-coding RNA XIST. *Mol Cancer* 19, 46
 48. Cohn WE and Volkin E (1951) Nucleoside-5'-phosphates from ribonucleic acid. *Nature* 167, 483-484
 49. Schwartz S, Bernstein DA, Mumbach MR et al (2014) Transcriptome-wide mapping reveals widespread dynamic-regulated pseudouridylation of ncRNA and mRNA. *Cell* 159, 148-162
 50. Li X, Zhu P, Ma S et al (2015) Chemical pulldown reveals dynamic pseudouridylation of the mammalian transcriptome. *Nat Chem Biol* 11, 592-597
 51. Hoang C and Ferre-D'Amare AR (2001) Cocystal structure of a tRNA^{Psi55} pseudouridine synthase: nucleotide flipping by an RNA-modifying enzyme. *Cell* 107, 929-939
 52. De Zoysa MD and Yu YT (2017) Posttranscriptional RNA Pseudouridylation. *Enzymes* 41, 151-167
 53. Duan J, Li L, Lu J, Wang W and Ye K (2009) Structural mechanism of substrate RNA recruitment in H/ACA RNA-guided pseudouridine synthase. *Mol Cell* 34, 427-439
 54. Rintala-Dempsey AC and Kothe U (2017) Eukaryotic stand-alone pseudouridine synthases - RNA modifying enzymes and emerging regulators of gene expression? *RNA Biol* 14, 1185-1196
 55. Montanaro L, Brigotti M, Clohessy J et al (2006) Dyskerin expression influences the level of ribosomal RNA pseudo-uridylation and telomerase RNA component in human breast cancer. *J Pathol* 210, 10-18
 56. Sieron P, Hader C, Hatina J et al (2009) DKC1 overexpression associated with prostate cancer progression. *Br J Cancer* 101, 1410-1416
 57. Liu B, Zhang J, Huang C and Liu H (2012) Dyskerin overexpression in human hepatocellular carcinoma is associated with advanced clinical stage and poor patient prognosis. *PLoS One* 7, e43147
 58. Alawi F, Lin P, Ziober B and Patel R (2011) Correlation of dyskerin expression with active proliferation independent of telomerase. *Head Neck* 33, 1041-1051
 59. Stockert JA, Gupta A, Herzog B, Yadav SS, Tewari AK and Yadav KK (2019) Predictive value of pseudouridine in prostate cancer. *Am J Clin Exp Urol* 7, 262-272
 60. Montanaro L, Calienni M, Bertoni S et al (2010) Novel dyskerin-mediated mechanism of p53 inactivation through defective mRNA translation. *Cancer Res* 70, 4767-4777
 61. McMahon M, Contreras A, Holm M et al (2019) A single H/ACA small nucleolar RNA mediates tumor suppression downstream of oncogenic RAS. *Elife* 8, e48847
 62. RajBhandary UL, Stuart A, Faulkner RD, Chang SH and Khorana HG (1966) Nucleotide sequence studies on yeast phenylalanine sRNA. *Cold Spring Harb Symp Quant Biol* 31, 425-434
 63. Li X, Xiong X, Zhang M et al (2017) Base-resolution mapping reveals distinct m(1)A methylome in nuclear- and mitochondrial-encoded transcripts. *Mol Cell* 68, 993-1005 e1009
 64. Roundtree IA, Evans ME, Pan T and He C (2017) Dynamic RNA modifications in gene expression regulation. *Cell* 169, 1187-1200
 65. Li X, Xiong X, Wang K et al (2016) Transcriptome-wide mapping reveals reversible and dynamic N(1)-methyladenosine methylome. *Nat Chem Biol* 12, 311-316
 66. Liu FG, Clark W, Luo GZ et al (2016) ALKBH1-mediated tRNA demethylation regulates translation. *Cell* 167, 816-828. e16
 67. Macari F, El-Houfi Y, Boldina G et al (2016) TRM6/61 connects PKC α with translational control through tRNAⁱ(Met) stabilization: impact on tumorigenesis. *Oncogene* 35, 1785-1796
 68. Dominissini D, Nachtergaele S, Moshitch-Moshkovitz S et al (2016) The dynamic N(1)-methyladenosine methylome in eukaryotic messenger RNA. *Nature* 530, 441-446
 69. Seo KW and Kleiner RE (2020) YTHDF2 Recognition of N-1-methyladenosine (m(1)A)-modified RNA is associated with transcript destabilization. *Acs Chemical Biol* 15, 132-139
 70. Chen Z, Qi M, Shen B et al (2019) Transfer RNA demethylase ALKBH3 promotes cancer progression via induction of tRNA-derived small RNAs. *Nucleic Acids Res* 47, 2533-2545
 71. Somme J, Van Laer B, Roovers M, Steyaert J, Versee W and Droogmans L (2014) Characterization of two homologous 2'-O-methyltransferases showing different specificities for their tRNA substrates. *RNA* 20, 1257-1271
 72. Rebane A, Roomere H and Metspalu A (2002) Locations of several novel 2'-O-methylated nucleotides in human 28S rRNA. *BMC Mol Biol* 3, 1
 73. Dai Q, Moshitch-Moshkovitz S, Han DL et al (2017) Nm-seq maps 2'-O-methylation sites in human mRNA with base precision (vol 14, pg 695, 2017). *Nat Methods* 15, 226-227
 74. Dimitrova DG, Teysset L and Carre C (2019) RNA 2'-O-methylation (Nm) modification in human diseases. *Genes (Basel)* 10, 117
 75. Belanger F, Stepinski J, Darzynkiewicz E and Pelletier J (2010) Characterization of hMT1, a human Cap1 2'-O-ribose methyltransferase. *J Biol Chem* 285, 33037-33044
 76. Werner M, Purta E, Kaminska KH et al (2011) 2'-O-ribose methylation of cap2 in human: function and evolution in a horizontally mobile family. *Nucleic Acids Res*

- 39, 4756-4768
77. Ching YP, Zhou HJ, Yuan JG, Qiang BQ, Kung Hf HF and Jin DY (2002) Identification and characterization of FTSJ2, a novel human nucleolar protein homologous to bacterial ribosomal RNA methyltransferase. *Genomics* 79, 2-6
 78. Koh CM, Gurel B, Sutcliffe S et al (2011) Alterations in nucleolar structure and gene expression programs in prostatic neoplasia are driven by the MYC oncogene. *Am J Pathol* 178, 1824-1834
 79. Su H, Xu T, Ganapathy S et al (2014) Elevated snoRNA biogenesis is essential in breast cancer. *Oncogene* 33, 1348-1358
 80. Marcel V, Ghayad SE, Belin S et al (2013) p53 acts as a safeguard of translational control by regulating fibrillarin and rRNA methylation in cancer. *Cancer Cell* 24, 318-330
 81. Squires JE, Patel HR, Nusch M et al (2012) Widespread occurrence of 5-methylcytosine in human coding and non-coding RNA. *Nucleic Acids Res* 40, 5023-5033
 82. Yang X, Yang Y, Sun BF et al (2017) 5-methylcytosine promotes mRNA export-NSUN2 as the methyltransferase and ALYREF as an m(5)C reader. *Cell Res* 27, 606-625
 83. Motorin Y, Lyko F and Helm M (2010) 5-methylcytosine in RNA: detection, enzymatic formation and biological functions. *Nucleic Acids Res* 38, 1415-1430
 84. Bohnsack KE, Hobartner C and Bohnsack MT (2019) Eukaryotic 5-methylcytosine (m(5)C) RNA methyltransferases: mechanisms, cellular functions, and links to disease. *Genes (Basel)* 10, 102
 85. Frye M and Watt FM (2006) The RNA methyltransferase Misu (NSun2) mediates Myc-induced proliferation and is upregulated in tumors. *Current Biol* 16, 971-981
 86. Goll MG, Kirpekar F, Maggert KA et al (2006) Methylation of tRNAAsp by the DNA methyltransferase homolog Dnmt2. *Science* 311, 395-398
 87. Tuorto F, Liebers R, Musch T et al (2012) RNA cytosine methylation by Dnmt2 and NSun2 promotes tRNA stability and protein synthesis. *Nat Struct Mol Biol* 19, 900-905
 88. Trixl L and Lusser A (2019) The dynamic RNA modification 5-methylcytosine and its emerging role as an epitranscriptomic mark. *Wiley Interdiscip Rev RNA* 10, e1510
 89. Chen X, Li A, Sun BF et al (2019) 5-methylcytosine promotes pathogenesis of bladder cancer through stabilizing mRNAs. *Nat Cell Biol* 21, 978-990
 90. Xue S, Xu H, Sun Z et al (2019) Depletion of TRDMT1 affects 5-methylcytosine modification of mRNA and inhibits HEK293 cell proliferation and migration. *Biochem Biophys Res Commun* 520, 60-66
 91. Ramanathan A, Robb GB and Chan SH (2016) mRNA capping: biological functions and applications. *Nucleic Acids Res* 44, 7511-7526
 92. Tomikawa C (2018) 7-methylguanosine modifications in transfer RNA (tRNA). *Int J Mol Sci* 19, 4080
 93. Pandolfini L, Barbieri I, Bannister AJ et al (2019) METTL1 promotes let-7 microRNA processing via m7G methylation. *Mol Cell* 74, 1278-1290 e1279
 94. Zhang LS, Liu C, Ma HH et al (2019) Transcriptome-wide mapping of internal N7-methylguanosine methylome in mammalian mRNA. *Mol Cell* 74, 1304-1316.e8
 95. Malbec L, Zhang T, Chen YS et al (2019) Dynamic methylome of internal mRNA N7-methylguanosine and its regulatory role in translation. *Cell Res* 29, 927-941
 96. Alexandrov A, Martzen MR and Phizicky EM (2002) Two proteins that form a complex are required for 7-methylguanosine modification of yeast tRNA. *RNA* 8, 1253-1266
 97. Liu Y, Zhang YS, Chi Q, Wang Z and Sun BS (2020) Methyltransferase-like 1 (METTL1) served as a tumor suppressor in colon cancer by activating 7-methylguanosine (m7G) regulated let-7e miRNA/HMGA2 axis. *Life Sci* 249, 117480
 98. Uddin MB, Wang Z and Yang C (2020) Dysregulations of functional RNA modifications in cancer, cancer stemness and cancer therapeutics. *Theranostics* 10, 3164-3189
 99. Niu Y, Wan A, Lin Z, Lu X and Wan G (2018) N(6)-methyladenosine modification: a novel pharmacological target for anti-cancer drug development. *Acta Pharm Sin B* 8, 833-843
 100. Wilson C, Chen PJ, Miao Z and Liu DR (2020) Programmable m(6)A modification of cellular RNAs with a Cas13-directed methyltransferase. *Nat Biotechnol* 38, 1431-1440