

RIEMANN SOLITONS ON CERTAIN TYPE OF KENMOTSU MANIFOLD

MANOJ RAY BAKSHI*, KANAK KANTI BAISHYA, AND ASHOKE DAS

ABSTRACT. The object of the present paper is to investigate the nature of Riemann solitons on generalized D -conformally deformed Kenmotsu manifold with hyper generalized pseudo symmetric curvature conditions.

1. Introduction

Let the symbols ∇ and ∇^d stand for the Riemann connection and the generalized D -conformally deformed connection respectively. Also, let R, S, Q, r and R^d, S^d, Q^d, r^d respectively stand for curvature tensor, Ricci tensor, Ricci operator, scalar curvature with respect to ∇ and ∇^d respectively. In this study, we consider an almost contact metric manifold $(M^{2n+1}, \phi, \xi, \eta, g)$ that consists of a $(1, 1)$ -tensor field ϕ , a vector field ξ and a 1-form η called respectively the structure endomorphism, the characteristic vector field and the contact form. In a recent paper, the authors ([2]) has introduced a new type of space called hyper generalized weakly symmetric manifold. Then the authors studied ([8]) hyper generalized pseudo Q -symmetric semi-Riemannian manifold. In Section 3 of this paper we extend this concept to generalized D -conformally deformed structure of a $(2n + 1)$ -dimensional Kenmotsu manifold.

A $(2n + 1)$ -dimensional Kenmotsu manifold is said to be hyper generalized pseudo symmetric (which will be abbreviated hereafter as $[H(GPS)_n, \nabla]$) if it admits the equation

$$\begin{aligned}
 & (\nabla_X \bar{R})(Y, U, V, W) \\
 = & 2\alpha(X)\bar{R}(Y, U, V, W) + \alpha(Y)\bar{R}(X, U, V, W) \\
 & + \alpha(U)\bar{R}(Y, X, V, W) + \alpha(V)\bar{R}(Y, U, X, W) \\
 & + \alpha(W)\bar{R}(Y, U, V, X) + 2\beta(X)(g \wedge S)(Y, U, V, W) \\
 & + \beta(Y)(g \wedge S)(X, U, V, W) + \beta(U)(g \wedge S)(Y, X, V, W) \\
 (1) \quad & + \beta(V)(g \wedge S)(Y, U, X, W) + \beta(W)(g \wedge S)(Y, U, V, X),
 \end{aligned}$$

Received January 20, 2021. Revised March 30, 2021. Accepted April 6, 2021.

2010 Mathematics Subject Classification: 53C15, 53C25.

Key words and phrases: Almost contact metric structure, generalized D -conformally deformation, Riemann solitons, η -Einstein manifolds.

* Corresponding author.

© The Kangwon-Kyungki Mathematical Society, 2021.

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/3.0/>) which permits unrestricted non-commercial use, distribution and reproduction in any medium, provided the original work is properly cited.

where

$$(2) \quad \begin{aligned} (g \wedge S)(Y, U, V, W) &= g(Y, W)S(U, V) + g(U, V)S(Y, W) \\ &\quad - g(Y, V)S(U, W) - g(U, W)S(Y, V), \end{aligned}$$

and α, β being non-zero 1-forms defined as $\alpha(X) = g(X, \theta_1)$ and $\beta(X) = g(X, \theta_2)$.

Ricci flow was first introduced by R. S. Hamilton ([17]) in 1982 which generalizes the notion of Riemann flow ([20], [19]). Keeping the tune with Ricci soliton, Hirica and Udriste ([18]) introduced and studied Riemann soliton. The Riemann flow is an evolution equation for metrics on a Riemannian manifold defined as follows

$$\frac{\partial}{\partial t} G(t) = -2R(g(t)), \quad t \in [0, I],$$

where $G = \frac{1}{2}g \otimes g$, \otimes is the Kulkarni-Nomizu product and R is the Riemann curvature tensor associated to the metric g . For $(0, 2)$ -tensors A and B , the Kulkarni-Nomizu product $(A \otimes B)$ is given by

$$(3) \quad \begin{aligned} (A \otimes B)(Y, U, V, Z) &= A(Y, Z)B(U, V) + A(U, V)B(Y, Z) \\ &\quad - A(Y, V)B(U, Z) - A(U, Z)B(Y, V). \end{aligned}$$

Recently, the present authors studied the Riemann solitons in the frame of $(LCS)_n$ -manifolds ([4]) and α -cosymplectic manifolds ([5]). The Riemann soliton is a smooth manifold M together with Riemannian metric g that satisfies

$$(4) \quad 2R + (g \otimes \mathcal{L}_W g) = 2\kappa(g \otimes g),$$

where W is a potential vector field, \mathcal{L}_W denotes the Lie-derivative along the vector field W and κ is a constant. A Riemann soliton is called expanding, steady and shrinking when $\kappa < 0$, $\kappa = 0$ and $\kappa > 0$ respectively.

We organize our present paper as follows: After Introduction, in Section 2, we briefly recall some known results for Kenmotsu manifolds and generalized D -conformally deformed of a Kenmotsu manifold and established some properties of the deformed Kenmotsu manifold. In Section 3, we discuss the properties of a generalized D -conformally deformed Kenmotsu manifold under hyper generalized pseudo symmetric curvature condition equipped with Riemann solitons. Finally, we determine a necessary condition for shrinking, steady and expanding of the soliton.

2. Preliminaries

According to the definition of Blair ([11]), an *almost contact structure* (ϕ, ξ, η) on a $(2n + 1)$ -dimensional Riemannian manifold satisfies the following conditions

$$(5) \quad \phi^2 = -I + \eta \otimes \xi,$$

$$(6) \quad \eta(\xi) = 1,$$

$$(7) \quad \phi\xi = 0, \quad \eta \circ \phi = 0, \quad \text{rank } \phi = n - 1.$$

Moreover, if g is a Riemannian metric on M^{2n+1} satisfying

$$(8) \quad g(\phi X, \phi Y) = g(X, Y) - \eta(X)\eta(Y),$$

$$(9) \quad g(X, \xi) = \eta(X),$$

$$(10) \quad g(\phi X, Y) = -g(X, \phi Y),$$

for any vector fields X, Y on M^{2n+1} , then the manifold M^{2n+1} ([11]) is said to admit an *almost contact metric structure* (ϕ, ξ, η, g) .

DEFINITION 2.1. [15] If in an almost contact metric structure (ϕ, ξ, η, g) on M^{2n+1} , the Riemann connection ∇ of g satisfies $(\nabla_X \phi)Y = g(\phi X, Y)\xi - \eta(Y)\phi X$, for any vector fields X, Y on M^{2n+1} , then the structure is called Kenmotsu.

PROPOSITION 2.2. [3,9,15] If $(M^{2n+1}, \phi, \xi, \eta, g)$ is a Kenmotsu manifold, then for any vector fields X, Y, Z on M^{2n+1} , the following relations hold

$$(11) \quad \nabla_X \xi = X - \eta(X)\xi,$$

$$(12) \quad (\nabla_X \eta)Y = g(X, Y)\xi - \eta(X)\eta(Y),$$

$$(13) \quad S(X, \xi) = -2n\eta(X),$$

$$(14) \quad \eta(R(X, Y)Z) = g(X, Z)\eta(Y) - g(Y, Z)\eta(X),$$

$$(15) \quad R(\xi, X)Y = \eta(Y)X - g(X, Y)\xi,$$

$$(16) \quad R(X, Y)\xi = \eta(X)Y - \eta(Y)X.$$

DEFINITION 2.3. [1] If a contact metric manifold M^{2n+1} with the almost contact metric structure (ϕ, ξ, η, g) is transformed into $(\phi^d, \xi^d, \eta^d, g^d)$, where

$$(17) \quad \phi^d = \phi, \quad \xi^d = \frac{1}{p}\xi, \quad \eta^d = p\eta, \quad g^d = qg + (p^2 - q)\eta \otimes \eta,$$

where p and q are constants such that $p \neq 0$ and $q > 0$, then the transformation is called a generalized D -conformal deformation.

Note that the generalized D -conformal deformation give rise to conformal deformation (for $p^2 = q$) and D -homothetic deformation (for $p = q = \text{constant}$) ([16], [6], [10]). The generalized D -conformal deformation are studied by various authors in ([22], [23], [21], [24]).

The relation between the Levi-Civita connections ∇ of g and ∇^d of g^d is given by ([1])

$$(18) \quad \nabla_X^d Y = \nabla_X Y + \frac{(p^2 - q)}{p^2}g(\phi X, \phi Y)\xi,$$

for any vector fields X, Y on M^{2n+1} .

In view of (17), (18) and definition of Riemannian curvature tensor, Ricci tensor, scalar curvature, we get the following:

PROPOSITION 2.4. [1] If a Kenmotsu structure (ϕ, ξ, η, g) on M^{2n+1} is transformed into $(\phi^d, \xi^d, \eta^d, g^d)$ under a generalized D -conformal deformation, then R, R^d, S, S^d, r and r^d are related by

$$(19) \quad R^d(X, Y)Z = R(X, Y)Z + \frac{(p^2 - q)}{p^2}[g(\phi Y, \phi Z)X - g(\phi X, \phi Z)Y],$$

$$(20) \quad S^d(X, Y) = S(X, Y) + \frac{2n(p^2 - q)}{p^2}g(\phi X, \phi Y),$$

$$(21) \quad r^d = \frac{r}{q} + \frac{2n(2n+1)(p^2 - q)}{p^2},$$

for any vector fields X, Y, Z on M^{2n+1} .

Now we shall bring out some properties of a generalized D -conformally deformed structure $(\phi^d, \xi^d, \eta^d, g^d)$ of a Kenmotsu manifold M^{2n+1} as follows:

PROPOSITION 2.5. *Under a generalized D -conformal deformation of a Kenmotsu structure (ϕ, ξ, η, g) on M^{2n+1} is transformed into $(\phi^d, \xi^d, \eta^d, g^d)$, then for any vector fields X, Y, Z on M^{2n+1} , we have*

$$(22) \quad \phi^d = -I + \eta^d \otimes \xi^d,$$

$$(23) \quad \eta^d(\xi^d) = 1,$$

$$(24) \quad \phi^d \xi^d = 0, \quad \eta^d \circ \phi^d = 0,$$

$$(25) \quad g^d(\phi^d X, \phi^d Y) = g^d(X, Y) - \eta^d(X)\eta^d(Y),$$

$$(26) \quad g^d(X, \xi^d) = \eta^d(X),$$

$$(27) \quad \nabla_X^d \xi^d = \frac{1}{p}[X - \eta^d(X)\xi^d],$$

$$(28) \quad (\nabla_X^d \eta^d)Y = \frac{1}{p}[g^d(X, Y) - \eta^d(X)\eta^d(Y)],$$

$$(29) \quad S^d(X, \xi^d) = -\frac{2n}{p^2}\eta^d(X),$$

$$(30) \quad \eta^d(R^d(X, Y)Z) = \frac{1}{p^2}[g^d(X, Z)\eta^d(Y) - g^d(Y, Z)\eta^d(X)],$$

$$(31) \quad R^d(\xi^d, X)Y = \frac{1}{p^2}[\eta^d(Y)X - g^d(X, Y)\xi^d],$$

$$(32) \quad R^d(X, Y)\xi^d = \frac{1}{p^2}[\eta^d(X)Y - \eta^d(Y)X].$$

Now using (28) and (29), we obtain

$$(33) \quad (\nabla_X^d S^d)(Y, \xi^d) = -\frac{1}{p}\left[\frac{2n}{p^2}g^d(X, Y) + S^d(X, Y)\right],$$

for any vector fields X, Y and Z on M^{2n+1} .

3. Main results

In the beginning, we shall define a hyper generalized pseudo symmetric space on a generalized D -conformally deformed structure $(\phi^d, \xi^d, \eta^d, g^d)$ of a Kenmotsu manifold M^{2n+1} .

3.1. Hyper generalized pseudo symmetric deformed Kenmotsu manifold.

DEFINITION 3.1. A generalized D -conformally deformed structure $(\phi^d, \xi^d, \eta^d, g^d)$ of a Kenmotsu manifold M^{2n+1} is said to be hyper generalized pseudo symmetric if it satisfies the condition

$$\begin{aligned}
 & (\nabla_X^d \bar{R}^d)(Y, U, V, W) \\
 = & 2\alpha^d(X)\bar{R}^d(Y, U, V, W) + \alpha^d(Y)\bar{R}^d(X, U, V, W) \\
 & + \alpha^d(U)\bar{R}^d(Y, X, V, W) + \alpha^d(V)\bar{R}^d(Y, U, X, W) \\
 & + \alpha^d(W)\bar{R}^d(Y, U, V, X) + 2\beta^d(X)(g^d \wedge S^d)(Y, U, V, W) \\
 & + \beta^d(Y)(g^d \wedge S^d)(X, U, V, W) + \beta^d(U)(g^d \wedge S^d)(Y, X, V, W) \\
 (34) \quad & + \beta^d(V)(g^d \wedge S^d)(Y, U, X, W) + \beta^d(W)(g^d \wedge S^d)(Y, U, V, X).
 \end{aligned}$$

where

$$\begin{aligned}
 (g^d \wedge S^d)(Y, U, V, W) = & g^d(Y, W)S^d(U, V) + g^d(U, V)S^d(Y, W) \\
 (35) \quad & - g^d(Y, V)S^d(U, W) - g^d(U, W)S^d(Y, V),
 \end{aligned}$$

and A_i^d are non-zero 1-forms defined by $A_i^d(X) = g^d(X, \sigma_i)$, for $i = 1, 2$.

In this section, we consider a Kenmotsu manifold (M^{2n+1}, g) $n \geq 1$ which is hyper generalized pseudo symmetric. Now, making use of (35) in (34), we find

$$\begin{aligned}
 & (\nabla_X \bar{R}^d)(Y, U, V, W) \\
 = & 2\alpha^d(X)\bar{R}^d(Y, U, V, W) + \alpha^d(Y)\bar{R}^d(X, U, V, W) \\
 & + \alpha^d(U)\bar{R}^d(Y, X, V, W) + \alpha^d(V)\bar{R}^d(Y, U, X, W) \\
 & + \alpha^d(W)\bar{R}^d(Y, U, V, X) + 2\beta^d(X)[g^d(Y, W)S^d(U, V) \\
 & + g^d(U, V)S^d(Y, W) - g^d(Y, V)S^d(U, W) \\
 & - g^d(U, W)S^d(Y, V)] + \beta^d(Y)[g^d(X, W)S^d(U, V) \\
 & + g^d(U, V)S^d(X, W) - g^d(X, V)S^d(U, W) \\
 & - g^d(U, W)S^d(X, V)] + \beta^d(U)[g^d(Y, W)S^d(X, V) \\
 & + g^d(X, V)S^d(Y, W) - g^d(Y, V)S^d(X, W) \\
 & - g^d(X, W)S^d(Y, V)] + \beta^d(V)[g^d(Y, W)S^d(U, X) \\
 & + g^d(U, X)S^d(Y, W) - g^d(Y, X)S^d(U, W) - g^d(U, W)S^d(Y, X)] \\
 & + \beta^d(W)[g^d(Y, X)S^d(U, V) + g^d(U, V)S^d(Y, X) \\
 (36) \quad & - g^d(Y, V)S^d(U, X) - g^d(U, X)S^d(Y, V)].
 \end{aligned}$$

Now, contracting over Y and W in (36), we get

$$\begin{aligned}
& (\nabla_X^d S^d)(U, V) \\
&= 2\alpha^d(X)S^d(U, V) + \alpha^d(U)S^d(X, V) \\
&\quad + \alpha^d(R^d(X, U)V) + 2\beta^d(Q^d X)g^d(U, V) \\
&\quad + \alpha^d(R^d(X, V)U) + \alpha^d(V)S^d(X, U) \\
&\quad + 2\beta^d(X)[2nS^d(U, V) + r^d g^d(U, V)] \\
&\quad + \beta^d(U) [(2n - 2)S^d(X, V) + r^d g^d(X, V)] \\
&\quad + \beta^d(V) [(2n - 2)S^d(X, U) + r^d g^d(X, U)] \\
(37) \quad & - \beta^d(Q^d U)g^d(X, V) - \beta^d(Q^d V)g^d(X, U).
\end{aligned}$$

Now setting $V = \xi^d$ and using (29), (31), (32) in the foregoing equation, we obtain

$$\begin{aligned}
& (\nabla_X^d S^d)(U, \xi^d) \\
&= -\frac{2n}{p^2}[2\alpha^d(X)\eta^d(U) + \alpha^d(U)\eta^d(X)] + \alpha^d(\xi^d)S^d(X, U) \\
&\quad + \frac{1}{p^2}[g^d(X, U)\alpha^d(\xi^d) - 2\eta^d(U)\alpha^d(X) + \eta^d(X)\alpha^d(U)] \\
&\quad + 2\beta^d(X)(r^d - \frac{4n^2}{p^2})\eta^d(U) + \beta^d(U)(r^d - \frac{4n(n-1)}{p^2})\eta^d(X) \\
&\quad + \beta^d(\xi^d) [2(n-1)S^d(U, X) + r^d g^d(U, X)] \\
(38) \quad & + 2\beta^d(Q^d X)\eta^d(U) - \beta^d(Q^d U)\eta^d(X) + \frac{2n}{p^2}\beta^d(\xi^d)g^d(U, X)
\end{aligned}$$

which yields by using (33)

$$\begin{aligned}
& -\frac{1}{p}[\frac{2n}{p^2}g^d(X, U) + S^d(X, U)] \\
&= [-\frac{2(2n+1)}{p^2}\alpha^d(X) + 2\beta^d(X)(r^d - \frac{4n^2}{p^2}) + 2\beta^d(Q^d X)]\eta^d(U) \\
&\quad + [-\frac{(2n-1)}{p^2}\alpha^d(U) + (r^d - \frac{4n(n-1)}{p^2})\beta^d(U) - \beta^d(Q^d U)]\eta^d(X) \\
&\quad + \frac{1}{p^2}\alpha^d(\xi^d)g^d(X, U) + \alpha^d(\xi^d)S^d(X, U) + \frac{2n}{p^2}\beta^d(\xi^d)g^d(U, X) \\
(39) \quad & + \beta^d(\xi^d) [2(n-1)S^d(U, X) + r^d g^d(U, X)].
\end{aligned}$$

Putting $X = U = \xi^d$ in succession, we obtain from (39) that

$$(40) \quad [r^d - \frac{2n(2n-1)}{p^2}]\beta^d(\xi^d) = \frac{2n}{p^2}\alpha^d(\xi^d).$$

$$\begin{aligned}
& -\frac{2(2n+1)}{p^2}\alpha^d(X) + 2\beta^d(X)(r^d - \frac{4n^2}{p^2}) + 2\beta^d(Q^d X) \\
(41) \quad & = [\frac{2(2n-1)}{p^2}\alpha^d(\xi^d) - 2\beta^d(\xi^d)(r^d - \frac{4n(n-1)}{p^2}) + 2\beta^d(Q^d \xi^d)]\eta^d(X).
\end{aligned}$$

and

$$\begin{aligned}
 & -\frac{(2n-1)}{p^2}\alpha^d(U) + (r^d - \frac{4n(n-1)}{p^2})\beta^d(U) - \beta^d(Q^dU) \\
 (42) \quad & = [\frac{(6n+1)}{p^2}\alpha^d(\xi^d) - (3r^d - \frac{12n^2-2n}{p^2})\beta^d(\xi^d)]\eta^d(U).
 \end{aligned}$$

By virtue of (40), (41) and (42), the equation (39) yields

$$\begin{aligned}
 & S^d(X, U) \\
 (43) \quad & = -\left(\frac{\frac{1}{p^2}\alpha^d(\xi^d) + (r^d + \frac{2n}{p^2})\beta^d(\xi^d) + \frac{2n}{p^3}}{\frac{1}{p} + \alpha^d(\xi^d) + 2(n-1)\beta^d(\xi^d)}\right)g^d(U, X) \\
 & -\left(\frac{\frac{(10n-1)}{p^2}\alpha^d(\xi^d) - (5r^d - \frac{20n^2-14n}{p^2})\beta^d(\xi^d)}{\frac{1}{p} + \alpha^d(\xi^d) + 2(n-1)\beta^d(\xi^d)}\right)\eta^d(U)\eta^d(X).
 \end{aligned}$$

and (40) gives

$$(44) \quad r^d = \frac{2n}{p^2}\left[\frac{\alpha^d(\xi^d)}{\beta^d(\xi^d)} + (2n-1)\right].$$

Next using (44) in (43) we have

$$\begin{aligned}
 & S^d(X, U) \\
 (45) \quad & = -\left(\frac{\frac{(2n+1)}{p^2}\alpha^d(\xi^d) + \frac{4n^2}{p^2}\beta^d(\xi^d) + \frac{2n}{p^3}}{\frac{1}{p} + \alpha^d(\xi^d) + 2(n-1)\beta^d(\xi^d)}\right)g^d(U, X) \\
 & +\left(\frac{\frac{1}{p^2}\alpha^d(\xi^d) + \frac{4n}{p^2}\beta^d(\xi^d)}{\frac{1}{p} + \alpha^d(\xi^d) + 2(n-1)\beta^d(\xi^d)}\right)\eta^d(U)\eta^d(X).
 \end{aligned}$$

This motivate us to state:

THEOREM 3.2. *Let $(\phi^d, \xi^d, \eta^d, g^d)$ be a generalized D -conformally deformed hyper generalized pseudo symmetric Kenmotsu manifold M^{2n+1} . Then such a space is conformally flat provided $\frac{1}{p} + \alpha^d(\xi^d) + 2(n-1)\beta^d(\xi^d) \neq 0$.*

THEOREM 3.3. *The scalar curvature of a hyper generalized pseudo symmetric generalized D -conformally deformed Kenmotsu manifold is constant.*

COROLLARY 3.4. *Let $(\phi^d, \xi^d, \eta^d, g^d)$ be a generalized D -conformally deformed pseudo symmetric Kenmotsu manifold M^{2n+1} . Then such a space is conformally flat provided $p\alpha^d(\xi^d) \neq -1$.*

4. Generalized D -conformally deformed hyper generalized pseudo symmetric Kenmotsu manifold and the case of Riemann soliton

In this section, we consider a generalized D -conformally deformed Kenmotsu manifold $(\phi^d, \xi^d, \eta^d, g^d)$ admitting a Riemann soliton. Then with the aid of (3) and (4),

we obtain

$$\begin{aligned}
 & 2R^d(Y, U, V, Z) + g^d(Y, Z) (\mathcal{L}_{\xi^d} g^d)(U, V) \\
 & + g^d(U, V) (\mathcal{L}_{\xi^d} g^d)(Y, Z) \\
 & - g^d(Y, V) (\mathcal{L}_{\xi^d} g^d)(U, Z) - g^d(U, Z) (\mathcal{L}_{\xi^d} g^d)(Y, V) \\
 (46) \quad & = 2\kappa [g^d(Y, Z) g^d(U, V) - g^d(Y, V) g^d(U, Z)].
 \end{aligned}$$

Now by contraction over Y and Z we get

$$(47) \quad \frac{1}{2} (\mathcal{L}_{\xi^d} g^d)(U, V) + \frac{1}{2n-1} S^d(U, V) = \frac{2n\kappa - \text{div}(\xi^d)}{2n-1} g^d(U, V).$$

and then using (47) in (46), we get

$$(48) \quad r^d = 2n[(2n+1)\kappa - \frac{4n}{p}].$$

Comparing (44) with (48) we have

$$(49) \quad (2n+1)p^2\kappa = 4np + (2n-1) + \frac{\alpha^d(\xi^d)}{\beta^d(\xi^d)}.$$

This leads to the following:

THEOREM 4.1. *Assume that a Kenmotsu structure (ϕ, ξ, η, g) on M^{2n+1} is transformed into $(\phi^d, \xi^d, \eta^d, g^d)$ under a generalized D -conformally deformation which is a hyper generalized pseudo symmetric space. Then the Riemann soliton is expanding, steady and shrinking as $\frac{\alpha^d(\xi^d)}{\beta^d(\xi^d)} + 2n(2p+1) <=> 1$.*

Acknowledgement. The first named author gratefully acknowledges to UGC, F.No. 16-6(DEC.2018)/2019(NET/CSIR) and UGC-Ref.No. 1147/(CSIR-UGC NET DEC. 2018) for financial assistance.

References

- [1] P. Alegre, D. E. Blair, and A. Carriazo, *Generalized Sasakian-space-forms*, Israel J. Math. **141** (1) (2004), 157–183.
- [2] K K Baishya, F. Ozen Zengin† and J Mike, *On hyper generalised weakly symmetric manifolds*, Nineteenth International Conference on Geometry, Integrability and Quantization June 02–07, 2017, Varna, Bulgaria Ivaïlo M. Mladenov and Akira Yoshioka, Editors Avangard Prima, Sofia 2018, pp 1–10 doi:10.7546/giq-19-2018-1-10
- [3] K. K. Baishya, P. Peska, and P. R. Chowdhury, *On almost generalized weakly symmetric Kenmotsu manifolds*, Acta Univ. Palacki. Olomuc., Fac. rer. nat. Mathematica **55** (2) (2016), 5–15.
- [4] M. R. Bakshi and K. K. Baishya, *Certain types of $(LCS)_n$ -manifolds and the case of Riemann solitons*, Differential Geometry-Dynamical Systems **22** (2020), 11–25.
- [5] M. R. Bakshi and K. K. Baishya, *Four classes of Riemann solitons on alpa-cosymplectic manifolds*, Afrika Matematika, <https://doi.org/10.1007/s13370-020-00846-6>
- [6] M. R. Bakshi, K. K. Baishya, D. G. Prakasha and P. Veerasha, *Ricci solitons in a hyper generalized pseudo symmetric D -homothetically deformed Kenmotsu manifold*, submitted
- [7] A. Biswas, A. Das, K. K. Baishya and M. R. Bakshi, *η -Ricci solitons on Kenmotsu manifolds admitting General connection*, Korean J. Math. **28** (2020) (4), 803–817, <http://dx.doi.org/10.11568/kjm.2020.28.4.803>
- [8] A. M. Blaga, M. R. Bakshi, and K. K. Baishya, *Hyper generalized pseudo Q -symmetric semi-Riemannian manifold*, Cubo, A Mathematical Journal, Vol. **23** (1) (2021), 87–96,
- [9] K. K. Baishya and P. R. Chowdhury, *On Generalized Weakly Symmetric Kenmotsu Manifolds*, Bol. Soc. Paran. Mat, (3s.) v. 39 6 (2021): 211–222.

- [10] A. M. Blaga, K. K. Baishya and N. Sarkar, *Ricci solitons in a generalized weakly (Ricci) symmetric D-homothetically deformed Kenmotsu manifold*, Ann. Univ. Paedagog. Crac. Stud. Math. **18** (2019), 123–136
- [11] D. E. Blair, *Contact Manifolds in Riemannian Geometry*, Lect. Notes in Math. 509, Springer-Verlag, New York (1976).
- [12] M. C. Chaki, *On pseudo symmetric manifolds*, Analele Stiintifice ale Universitatii "Al I. Cuza" din Iasi **33** (1987), 53–58.
- [13] R. S. Hamilton, *The Ricci flow on surfaces*, Contemp. Math. **71** (1988), 237–261.
- [14] R. S. Hamilton, *Three-manifolds with positive Ricci curvature*, J. Diff. Geom. **17** (1982), 255–306.
- [15] K. Kenmotsu, *A class of almost contact Riemannian manifolds*, Tohoku Math. J. **24** (1972), 93–103
- [16] Tanno, S., *The topology of contact Riemannian manifolds*, Illinois J. Math. **12** (1968), 700–717.
- [17] R. S. Hamilton, *The Ricci flow on surfaces*, Mathematics and general relativity, Contemp. Math. **71**, American Math. Soc. (1988), 237–262.
- [18] I.E. Hirićă, C. Udriste, *Ricci and Riemann solitons*, Balkan J. Geom. Applications. **21** (2) (2016), 35–44.
- [19] C. Udriște, *Riemann flow and Riemann wave via bialternate product Riemannian metric*. preprint, arXiv.org/math.DG/1112.4279v4 (2012).
- [20] C. Udriste, *Riemann flow and Riemann wave*, Ann. Univ. Vest, Timisoara. Ser. Mat. Inf. **48** (1-2) (2010), 265–274.
- [21] Nülifer Özdemir, Sirin Aktay, Mehmet Solgun, *On generalized D-conformal deformations of certain almost contact metric manifolds*, Mathematics 2019, 0700168.
- [22] Nagaraja, H.G., Kiran Kumar, D.L., *Ricci Solitons in Kenmotsu Manifold under Generalized D-Conformal Deformation*. Lobachevskii J Math **40** (2019), 195–200. <https://doi.org/10.1134/S1995080219020112>.
- [23] HG Nagaraja, DL Kiran Kumar, VS Prasad, *Ricci solitons on Kenmotsu manifolds under D-homothetic deformation*, Khayyam J. Math. **4** (1) (2018), 102–109.
- [24] T. Suguri and S. Nakayama, *D-conformal deformations on almost contact metric structure*, Tensor (N.S.) **28** (1974), 125–129.

Manoj Ray Bakshi

Department of Mathematics, Raiganj University, Uttar Dinajpur,
West Bengal, India
E-mail: raybakshimanoj@gmail.com

Kanak Kanti Baishya

Department of Mathematics, Kurseong College, Kurseong, Darjeeling,
West Bengal, India
E-mail: kanakkanti.kc@gmail.com

Ashoke Das

Department of Mathematics, Raiganj University, Uttar Dinajpur,
West Bengal, India
E-mail: ashoke.avik@gmail.com