

Erratum to: Evaluation of adenosine triphosphate testing for on-farm cleanliness monitoring compared to microbiological testing in an empty pig farrowing unit

Seung-Won Yi, Ara Cho, Eunju Kim, Sang-Ik Oh, Jae Hee Roh, Young-Hun Jung, Changyong Choe, Jae Gyu Yoo and Yoon Jung Do*

Division of Animal Diseases & Health, National Institute of Animal Science, Rural Development Administration, Wanju 55365, Korea



*Corresponding author

Yoon Jung Do
 Division of Animal Diseases & Health,
 National Institute of Animal Science,
 Rural Development Administration,
 Wanju 55365, Korea.
 Tel: +82-63-238-7222
 E-mail: clonea@korea.kr

Copyright © 2022 Korean Society of Animal Sciences and Technology. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0/>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

ORCID

Seung-Won Yi
<https://orcid.org/0000-0001-5545-2969>
 Ara Cho
<https://orcid.org/0000-0001-5309-7721>
 Eunju Kim
<https://orcid.org/0000-0003-4040-0474>
 Sang-Ik Oh
<https://orcid.org/0000-0003-0877-9170>
 Jae Hee Roh
<https://orcid.org/0000-0003-4223-1096>
 Young-Hun Jung
<https://orcid.org/0000-0002-8094-0304>
 Changyong Choe
<https://orcid.org/0000-0003-4222-3360>
 Jae Gyu Yoo
<https://orcid.org/0000-0002-8542-9193>
 Yoon Jung Do
<https://orcid.org/0000-0003-3207-3514>

ERRATUM

In the published article “Evaluation of adenosine triphosphate testing for on-farm cleanliness monitoring compared to microbiological testing in an empty pig farrowing unit. *J Anim Sci Technol* 2020;62(5):682-91. <https://doi.org/10.5187/jast.2020.62.5.682>,” fecal indicator bacteria listed second should be modified to total aerobic bacteria in the legend of Fig. 2a [1]. The editorial office would like to correct it.

REFERENCES

1. Yi SW, Cho A, Kim E, Oh SI, Roh JH, Jung YH, et al. Evaluation of adenosine triphosphate testing for on-farm cleanliness monitoring compared to microbiological testing in an empty pig farrowing unit. *J Anim Sci Technol*. 2020;62:682-91. <https://doi.org/10.5187/jast.2020.62.5.682>




Fig. 2. Comparison of adenosine triphosphate (ATP) test and microbial status results from plate counting (total aerobic bacteria, fecal indicator bacteria) according to the cleaning status of the (a) floor center, (b) floor corner, and (c) feeding trough of the pig farrowing unit ($p < 0.05$, $^{**}p < 0.01$ vs BC). BC, before cleaning; AC, after cleaning; AD, after disinfection.