



Research Article

Non-destructive quality prediction of domestic, commercial red pepper powder using hyperspectral imaging

Sang Seop Kim¹, Ji-Young Choi², Jeong Ho Lim¹, Jeong-Seok Cho^{1*}

¹Food Safety and Distribution Research Group, Korea Food Research Institute, Wanju 55365, Korea

²Practical Technology Research Group, World Institute of Kimchi, Gwangju 61755, Korea

Abstract We analyzed the major quality characteristics of red pepper powders from various regions and predicted these characteristics nondestructively using shortwave infrared hyperspectral imaging (HSI) technology. We conducted partial least squares regression analysis on 70% (n=71) of the acquired hyperspectral data of the red pepper powders to examine the major quality characteristics. R_c^2 values of ≥ 0.8 were obtained for the ASTA color value (0.9263) and capsaicinoid content (0.8310). The developed quality prediction model was validated using the remaining 30% (n=35) of the hyperspectral data; the highest accuracy was achieved for the ASTA color value ($R_p^2=0.8488$), and similar validity levels were achieved for the capsaicinoid and moisture contents. To increase the accuracy of the quality prediction model, we conducted spectrum preprocessing using SNV, MSC, SG-1, and SG-2, and the model's accuracy was verified. The results indicated that the accuracy of the model was most significantly improved by the MSC method, and the prediction accuracy for the ASTA color value was the highest for all the spectrum preprocessing methods. Our findings suggest that the quality characteristics of red pepper powders, even powders that do not conform to specific variables such as particle size and moisture content, can be predicted via HSI.



OPEN ACCESS

Citation: Kim SS, Choi JY, Lim JH, Cho JS. Non-destructive quality prediction of domestic, commercial red pepper powder using hyperspectral imaging. Korean J Food Preserv, 30(2), 224-234 (2023)

Received: March 15, 2023

Revised: April 07, 2023

Accepted: April 10, 2023

***Corresponding author**

Jeong-Seok Cho
Tel: +82-63-219-9140
E-mail: jscho@kfri.re.kr

Copyright © 2023 The Korean Society of Food Preservation. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<http://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Keywords red pepper powder, chemical components, hyperspectral imaging, capsaicinoids, water contents

1. Introduction

Red pepper (*Capsicum annuum* L.), which has a characteristic spicy flavor and red color, is one of the most widely consumed vegetables worldwide (FAO STAT, 2012). Most red peppers are stored in a dried state to be processed into red pepper powder (Ku et al., 2001) through simple grinding. Red pepper powder is the main ingredient of widely consumed foods in South Korea, including *kimchi* and *gochujang* (red pepper paste). It is also the spice with the highest level of consumption (Kim et al., 2019). The components of red pepper powder that affect its quality include capsaicin, vitamin C, free sugars, organic acids, and pigments, whose contents vary according to the particle size in each use (Oh et al., 2011). For red peppers, the content of capsaicinoids, which are responsible for the hot

taste, can vary for a given cultivar depending on the time of harvest and the cultivation conditions, e.g., daylight exposure, precipitation, and soil properties. Thus, quantifying and rating the hot taste of red pepper powder are critical (Mo et al., 2012). Numerous studies have been performed on the prediction of the particle size or the contents of capsaicinoids and other compounds of red pepper powder; in the research of Mo et al. (2011) and Mo et al. (2012), the quality characteristics of red pepper powder were analyzed. Nonetheless, most of the previous studies were conducted with fixed states of the particle size, the water content, and other factors of red pepper powder. The Korean Industrial Standard (KS, H 2157:2021) for red pepper powder limits the permitted water (≤ 13.0 wt.%) and ash (≤ 7.0 wt.%) contents. However, owing to the wide range of limits on the particle size and water content, there is a lack of uniformity in the quality characteristics of commercial red pepper powders across regions and producers.

The quality of red pepper powder is evaluated mainly according to the hot-taste compounds as an intrinsic factor and the color values as an extrinsic factor.

The hot taste of red pepper powder is reported to enhance the flavor of food and increase the appetite (Perucka and Oleszek, 2000). The main compounds responsible for the hot taste are capsaicinoids composed of several saturated and unsaturated moieties such as capsaicin and dihydrocapsaicin (Constant et al., 1995; Iwai et al., 1979). The redness [American Spice Trade Association (ASTA) color value] of red pepper powder is the most powerful influencing factor of consumer choice of the product (Hornero-Mendez and Minguez-Mosquera, 2001) and significantly affects the quality evaluation and product values. An increase in the

ASTA color value an international unit defined by the ASTA to describe the color of red pepper powder was shown to lead to an increase in the content of red pigments (Yoon and Lee, 2004). At present, the level of hot taste, which is indicated by the label of commercial red pepper powder, varies across regions and producers, and compliance with legal guidelines is inadequate. Hence, there is a challenge in standardizing the hot taste as well as other quality characteristics of red pepper powder, and the criteria for consumer use remain ambiguous. It is thus necessary to develop a standard technique for measuring the quality characteristics of red pepper powder that can be applied by different regions and producers (Mo et al., 2012).

Various techniques are currently available for high-quality measurements of agricultural products through chemical analyses. However, such techniques are expensive and time-consuming and pose challenges in conducting exhaustive research. The most significant disadvantage is the destruction of the sample being analyzed (Gomez et al., 2006). In contrast, the use of nondestructive spectral analysis allows rapid high-quality measurements without preprocessing or a negative impact on the target agricultural product (Jha and Matsuoka, 2004). Among the nondestructive analytical methods, the reflectance method is widely used to measure the quality of agricultural products. It involves the irradiation of the sample with light and the reaction of the penetrating light with the particles close to the surface; the light is dispersed in varying directions, producing reflectance that is used for quality measurements (Lee et al., 2004).

Therefore, in this study, random samples of different commercial red pepper powders that varied in particle size and water content were purchased, and hyperspectral data were obtained from each

sample using the hyperspectral imaging (HSI) technology of the shortwave infrared (SWIR) range, which was applied to the surface of each sample. The contents of capsaicinoids, water, and total sugar and the ASTA value were measured as the criteria of the quality characteristics of red pepper powder to evaluate the potential use of the SWIR HSI technique for nondestructive analyses of the quality characteristics of red pepper powder.

2. Materials and methods

2.1. Test materials

The red pepper powders used in this study were random samples of commercial red pepper powders with varying particle sizes and hot-taste levels. Six samples were purchased from each of the following 17 regions: Jinan-gun (produced in October 2021), Imsil-gun, Gochang-gun, Sintaein-eup of Jeongeup-si, Gwancheon-myeon of Imsil-gun, Jeong-eup-si, Andong-si, Yongin-si, Yeongju-si, Jinan-gun (produced in November 2021), Osu-myeon of Imsil-gun, Bugwi-myeon of Jinan-gun, Jeongcheon-myeon of Jinan-gun, Gyebuk-myeon of Jangsu-gun, Maryeong-myeon of Jinan-gun, Mujang-myeon of Gochang-gun, and Wonpyeong-ri of Geumsan-myeon of Gimje-si. Additionally, four samples were purchased from Uiseong-gun. In total, 106 samples were used to obtain hyperspectral images and spectroscopic data.

2.2. Analysis of quality characteristics of red pepper powders

2.2.1. Capsaicinoid content

To precisely weigh 2 g of red pepper powder, 10 mL of methanol was added with five boiling tips for thorough mixing. After 1 h extraction on a Dri-Block Heater (MaXtable H10, DaiHan, Incheon,

Korea) at 90°C, the solution was cooled at the ambient temperature for 1 h. The cooled extraction solvent was filtered using filter paper (Whatman No. 1) and then a 0.2- μm syringe filter, and the resulting capsaicinoid extract was used in subsequent analyses. For quantification, 20 μL of the extract was injected into a high-performance liquid chromatography (HPLC) system (Agilent 1260 infinity II, Agilent Technology, Agilent Co., Santa Clara, CA, USA). For the XTerraTMRP18 (5 μm , 4.6 \times 150 mm id., Waters, Milford, MA, USA) column set at 35°C, 1% acetic acid and acetonitrile were used as the mobile phase. At a flow rate of 1 mL/min, the separation was based on the gradient method, as shown in Table 1. Ultraviolet light with a wavelength of 280 nm was used for absorbance measurements. The reference compounds were capsaicin standard (Sigma-Aldrich, Co. Ltd., St. Louis, MO, USA) and dihydrocapsaicin standard (Sigma-Aldrich, Co. Ltd.).

2.2.2. Water content

The water content of red pepper powder was measured as the difference between the weights before and after drying for approximately 3 g of red pepper powder on an aluminum plate of a known weight. For the analysis, the sample was dried for 6 h in a 70°C vacuum oven dryer (vacuum dryer OV-11, Jeio Tech, Daejeon, Korea) in accordance with the ASTA analytical official method (1997).

Table 1. Capsaicinoid analytical conditions for HPLC

Time (min)	(A) 1% acetic acid in water (%)	(B) Acetonitrile (%)
0-3	60	40
15	38	62
16	20	80
20	20	80

2.2.3. ASTA color value

The ASTA color value was measured in accordance with the ASTA analytical official method. In a 100-mL volumetric flask, accurately weighed 0.1 g of red pepper powder was placed, and acetone was added to adjust the volumes to a uniform level. After 1 min of shaking, the mixture was left in a dark room for 16 h to prepare the test solution. The absorbance was measured at 460 nm, and the following equation was used for the calculation.

$$\text{ASTA value} = \frac{A \times 16.4}{W}$$

A: Absorbance at 460 nm

W: Sample weight (g)

2.2.4. Total sugar

To accurately weigh 2 g of red pepper powder, 40 mL of 80% ethanol was added for 1-min extraction on a vortex mixer. After filtration using a 0.2- μm membrane filter, 20 μL of the extract was injected into the HPLC system (Agilent 1260 infinity II, Agilent Technology). YMC-Pack Polyamine II (4.6 \times 250 mm², 5 μm , YMC Company Limited, Kyoto, Japan) was used as the column. The temperatures of the column and RI detector were set at 30 and 35°C. The solvent was acetonitrile: water = 75:25, and the flow rate was 1 mL/min. The standard compounds were fructose, glucose, and sucrose (Sigma-Aldrich, Co., Ltd.), and the sum of the contents of these compounds was used as the total sugar content.

2.3. Hyperspectral data analysis

2.3.1. Hyperspectral data collection

The sample was placed against a black background plate, and the SWIR HSI system (IMEC Kapeldreef 75, 3001 Heverlee, Belgium) with a spectral range of 900–1,700 nm and image resolution of 2,048 \times 1,088

was used. The line scan method was used to obtain the hyperspectral images. The number of bands was 256. The spectrometer had two halogen lamps on either side. For the spectral acquisition, the frame rate was 10, and the exposure time during the measurements was 10 s. For the measurements, 4 g of each sample was placed in a clear cell culture plate with a diameter of 2.5 cm and depth of 2 cm (SPL Life Sciences Co., Pocheon, Korea), and the mean of 10 measurements was calculated. Each spectral image was extracted using the Region of Interest function of the ENVI software (version 5.4, Exelis Visual Information Solutions, Harris Co., Boulder, Co, USA).

2.3.2. Partial least squares analysis

To develop a model for predicting the internal quality according to the hyperspectral reflectance spectrum, partial least squares (PLS) regression analysis was used. In the PLS analysis, the spectral data of nondestructive measurements were applied as the independent variables, and the data of destructive measurements, i.e., the capsaicinoid, water, and total sugar contents and ASTA color value, were used as the dependent variables. Thus, a linear regression model was constructed. The multivariate independent variables were reduced to new variables, whereby a regression equation was obtained to develop a prediction model. For the dependent variables, 3 or 4 samples of red pepper powder from each of the 18 regions were used. Ten measurements were performed for each sample. The samples were divided at a ratio of 7:3; i.e., 71 were used for model development, and 35 were used for model validation. The accuracies of the calibration and prediction models were expressed by the determination coefficient of calibration (R_c^2) and the determination coefficient of prediction

(R_p^2), respectively. The Unscrambler X software (10.2, Camo, Oslo, Norway) was used to construct the prediction model for the quality characteristics of red pepper powder.

2.3.3. Spectral preprocessing

For the extracted data, various preprocessing methods, including the standard normal variate (SNV), multiple scatter correction (MSC), and Savitzky-Golay derivative (SG), were applied to remove noise and reduce the resulting errors in the spectral data according to the characteristics of the analyzed samples. The SNV and MSC were used to remove the light scattering in the spectrum. The SG was used to eliminate the shift of the baseline due to the difference in the optical path or changes in

analytical conditions and to highlight the micro-characteristics of the spectrum (Han et al., 2015).

3. Results and discussion

3.1. Quality characteristics of red pepper powders

Table 2 presents the results of analyzing the main compounds that influence the quality of red pepper powders in the samples randomly purchased from different regions in South Korea. The contents of capsaicinoids distributed across the regions ranged from 42.44 to 840.03 mg/kg, with a large regional deviation at 61.36 mg/kg. This agreed with the studies of Shin and Lee (1991), Park (2004), and Kim et al. (2006), wherein the contents of capsaicinoids as the main compounds responsible for the hot taste

Table 2. Physicochemical properties of various types of red pepper powders

No.	Samples	Hot taste level ¹⁾	Capsaicinoids (mg/kg)	Water content (%)	ASTA color value	Total sugar (%)
1	Jinan-gun (2021. 10)	Medium hot	314.83±15.92	11.23±1.13	86.95±2.64	13.45±2.70
2	Imsil-gun	Slight hot	195.00±13.03	9.31±1.08	65.97±1.68	13.43±3.62
3	Gochang-gun	Slight hot	163.63±13.28	10.13±0.93	121.71±3.87	11.74±1.99
4	Jeongeup-si, Sintaein-eup	Slight hot	269.50±17.64	9.54±1.09	88.82±1.48	17.15±1.54
5	Imsil-gun, Gwanchon-myeon	Very hot	525.65±35.69	6.89±0.68	79.14±1.85	15.22±2.06
6	Jeongeup-si	Very hot	816.69±27.60	9.08±1.39	101.38±2.38	14.86±2.41
7	Andong-si	Mild hot	126.04±12.71	8.15±0.87	75.09±2.55	16.57±1.13
8	Yongin-si	Medium hot	423.80±27.35	6.44±0.64	77.15±2.53	15.38±1.59
9	Yeongju-si	Mild hot	91.42±20.61	6.47±0.89	99.99±5.07	10.85±1.08
10	Jinan-gun (2021. 11)	Very hot	840.03±29.61	7.74±2.39	87.97±5.28	7.04±1.44
11	Imsil-gun, Osu-myeon	Mild hot	42.44±14.05	10.10±0.64	70.87±5.14	13.27±2.25
12	Jinan-gun, Bugwi-myeon	Mild hot	46.40±9.29	9.11±0.59	73.65±1.38	15.01±1.15
13	Jinan-gun, Jeongcheon-myeon	Mild hot	122.13±11.09	8.35±0.58	70.98±2.09	14.44±1.49
14	Jangsu-gun, Gyebuk-myeon	Slight hot	168.05±61.36	7.35±1.40	78.22±6.46	13.97±2.52
15	Jinan-gun, Maryeong-myeon	Medium hot	404.22±17.60	6.19±0.33	63.10±3.10	15.82±1.25
16	Uiseong-gun	Mild hot	148.71±31.86	7.60±1.16	66.30±4.33	12.38±0.87
17	Gochang-gun, Mujang-myeon	Slight hot	192.25±46.07	9.58±0.05	110.57±2.86	11.99±0.97
18	Gimje-si, Geumsan-myeon, Wonpyeong-ri	Slight hot	215.89±58.12	6.17±0.53	66.51±8.10	13.10±3.15

¹⁾Very hot, 1,000–500 mg/kg; medium hot, 500–300 mg/kg; slight hot, 300–150 mg/kg; mild hot, less than 150 mg/kg.

of red pepper were shown to vary among different cultivars and considerably according to the region and conditions of cultivation even within an identical cultivar. For the water content, the range of the distribution was 6.17%-11.23%, satisfying the quality standard criterion of $\leq 15\%$ for commercial red pepper powders (Ku et al., 2004). The range also satisfied the KS criterion of $\leq 13.0\%$. The ASTA color value ranged from 63.10 to 121.71. This coincided with the results of Kim et al. (2002) and Kim et al. (2008), who reported that the ASTA color value can be affected in different ways by variables such as the time of fruit harvest, position of fruiting, and time of sunlight exposure. The range of the total sugar content was 7.04%-17.15%. All the quality indicators analyzed in this study exhibited large deviations in the measured values. This is attributed to the differences in the red pepper cultivar and region of cultivation, as well as the lack of uniformity across variables such as the particle size and water

content.

3.2. Analysis of hyperspectral reflectance spectra

The mean spectra at 900-1,700 nm obtained for the red pepper powder samples produced in 18 regions are presented in Fig. 1. Mo et al. (2011) reported that capsaicinoids were associated with the 1,100-1,120, 1,320-1,350, 1,510-1,530, and 1,620-1,640-nm ranges. Krepper et al. (2018) reported that the water content was associated with the 1,440-1,450 nm range, and another study indicated that the spectrum at 935 nm was associated with free sugars (Shao et al., 2021). In the present study, the hyperspectral reflectance spectra of red pepper powder exhibited varying patterns according to the region of production. According to the differences in the reflectance spectra in the areas associated with capsaicinoids and water, a model for the nondestructive prediction of the actual capsaicinoid, water, and total sugar contents was developed.




Fig. 1. Mean spectra of various types of red pepper powders using HSI.

3.3. Prediction of quality indicators of red pepper powders using partial least squares regression

The PLSR-based model for predicting the capsaicinoid, water, and total sugar contents and ASTA color value for red pepper powder is presented in Fig. 2. The R_c^2 of the model was the largest (0.9263) for the ASTA color value, followed by 0.8310 for the capsaicinoid content and 0.7408 for the total sugar content, and was the smallest (0.6910) for the water content. The small R_c^2 for the water content is attributed to the large deviation in the particle size of red pepper powder across regions and the resulting variation in surface area, which prevented uniformity in the rate of water evaporation among the samples. The particle size of red pepper powder leads to the variation of the reflectance intensity, as indicated by previous studies such as those of Barajas et al. (2007) and Gupta et al. (2005), who reported that physical

properties such as the particle size, distribution, and density were associated with the reflectance spectra. In a follow-up study, these factors should be adjusted to obtain larger R_p^2 values.

The hyperspectral reflectance spectra of red pepper powders from different regions were used to validate the prediction model for the main quality indicators. For the validation, 30% ($n=35$) of the samples were used to test the PLSR model developed using 70% ($n=71$) of the samples, and the results are presented in Fig. 3. The R_p^2 was the largest (0.8488) for the ASTA color value. The R_p^2 was similar between the capsaicinoid content (0.6690) and the water content (0.6688). The R_p^2 was the smallest (0.5611) for the total sugar content. For both R_c^2 and R_p^2 , the ASTA color value, which indicated the content of red pepper pigments, and the content of capsaicinoids as the hot-taste compounds exhibited large values. This was consistent with the results of




Fig. 2. Calibration results of the PLSR model ($n=71$) for various types of red pepper powders obtained using HSI.




Fig. 3. Prediction accuracy of quality indicators for various types of red pepper powders obtained using HSI.

Ananthan et al. (2018), Barbero et al. (2014), and Contreras-Padilla and Yahia (1998), who reported that the capsaicinoid concentrations increased as red peppers matured with the color change from green to red and that the capsaicinoid concentration was correlated with the red color. These findings verified the possibility of predicting the hot taste and the ASTA color value as the main quality characteristics of red pepper powder even without fixing factors such as the region of production, particle size, and water content. Additionally, as reported by Lim et al. (2012), changes in the water content during sample collection are not likely to induce a significant variation in the capsaicinoid content or the ASTA color value.

3.4. Optimization of prediction model according to spectral preprocessing method

Table 3 presents the results of model validation

using the PLSR model to which various preprocessing methods were applied for increasing the accuracy of quality analysis. The R_p^2 was 0.8892 and 0.8984 when the SNV and MSC preprocessing methods, respectively, were applied for preventing the impact of light scattering, which indicated the highest validity of the model for the ASTA color value. The R_p^2 was 0.8413 and 0.6025 when the SG primary and secondary gradient preprocessing methods, respectively, were applied for preventing the spectral baseline shift caused by the optical path or analytical conditions or for highlighting the micro-characteristics, which also indicated that model had the highest validity for the ASTA color value. For all the spectral preprocessing methods, the prediction performance was outstanding for the capsaicinoid content and the ASTA color value. Among the preprocessing methods, the highest accuracy was achieved with the MSC without fixing

Table 3. Comparison of models using different spectral preprocessing methods to evaluate the PLSR model

	Pre-processing method ¹⁾							
	SNV		MSC		SG-1		SG-2	
	R _p ^{2 2)}	RMSEP ³⁾	R _p ²	RMSEP	R _p ²	RMSEP	R _p ²	RMSEP
Capsaicinoids	0.7560	113.2623	0.7809	109.504	0.7788	117.6174	0.3275	210.8753
Water contents	0.6634	1.0433	0.6811	0.9995	0.6979	0.9982	0.5475	1.2035
ASTA color value	0.8892	5.4664	0.8984	5.2210	0.8413	6.5242	0.6025	10.6200
Total sugar	0.5799	1.9382	0.5677	1.9662	0.5244	2.0767	0.2208	2.6644

¹⁾SNV, standard normal variate; MSC, multiple scatter correction; SG-1, Savitzky-Golay 1st derivative; SG-2, Savitzky-Golay 2nd derivative.

²⁾R², coefficient of the prediction.

³⁾RMSEP, root mean square error of prediction.

factors such as the particle size and water content.

4. Summary

The main quality characteristics of domestic commercial red pepper powders from different regions in South Korea were analyzed, and SWIR HSI technology was applied for nondestructive quality prediction. For 106 red pepper powder samples from 18 regions, the contents of capsaicinoids, water, and total sugar and the ASTA color value were measured. The measured values exhibited large deviations due to various factors, such as the region of cultivation, particle size, and water content. The acquired hyperspectral data were used to conduct PLSR analyses on 70% (n=71) of the red pepper powder samples regarding the main quality characteristics. The R_c² was ≥0.8 for the ASTA color value (0.9263) and capsaicinoid content (0.8310). The R_p² from the model validation using 30% (n=35) of the hyperspectral data was the largest (0.8488) for the ASTA color value. The validity was similar for the contents of capsaicinoids and water. To increase the accuracy of the quality prediction model, spectral preprocessing was performed using the SNV, MSC, SG-1, and SG-2 methods, and the resulting model accuracy was increased to the

largest extent with the MSC method. For all the spectral preprocessing methods, the prediction accuracy was the highest for the ASTA color value. Our results confirm the possibility of predicting the quality characteristics of red pepper powder using HSI technology, even for commercial red pepper powders with varying particle sizes and water contents.

Acknowledgements

This work was funded by the Ministry of Agriculture, Food and Rural Affairs and supported by the High Value added Food Technology Development Project of the Korea Institute of Food Technology Planning and Evaluation.

Conflict of interests

The authors declare no potential conflicts of interest.

Author contributions

Conceptualization : Cho JS. Formal analysis: Choi JY. Validation: Lim JH. Writing - original draft: Kim SS. Writing - review & editing: Cho JS.

Ethics approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

ORCID

Sang Seop Kim (First author)

<https://orcid.org/0000-0001-5653-7788>

Ji-Young Choi

<https://orcid.org/0000-0002-7854-9277>

Jeong Ho Lim

<https://orcid.org/0000-0002-4806-2046>

Jeong-Seok Cho (Corresponding author)

<https://orcid.org/0000-0002-9513-4014>

References

- Ananthan R, Subhash K, Longvah T. Capsaicinoids, amino acid and fatty acid profiles in different fruit components of the world hottest Naga king chilli (*Capsicum chinense Jacq*). Food Chem, 238, 51-57 (2018)
- Barajas MJ, Cassiani AR, Vargas W, Conde C, Roper J, Figueroa J, Romanach RJ. Near-infrared spectroscopic method for real-time monitoring of pharmaceutical powders during voiding. Appl Spectrosc, 61, 490-496 (2007)
- Barbero GF, Ruiz AG, Liazid A, Palma M, Vera JC, Barroso CG. Evolution of total and individual capsaicinoids in peppers during ripening of the Cayenne pepper plant (*Capsicum annum L.*). Food Chem, 153, 200-206 (2014)
- Constant HL, Cordell GA, West DP, Johnson JH. Separation and quantification of capsaicinoids using complexation chromatography. J Nat Prod, 58, 1925-1928 (1995)
- Contreras-Padilla M, Yahia EM. Changes in capsaicinoids during development, maturation, and senescence of chile peppers and relation with peroxidase activity. J Agric Food Chem, 46, 2075-2079 (1998)
- Gomez AH, He Y, Pereira AG. Non destructive measurement of acidity, soluble solids and firmness of Satsuma mandarin using vis/NIR spectroscopy techniques. J Food Eng, 77, 313-319 (2006)
- Gupta A, Peck GE, Miller RW, Morris KR. Real-time near-infrared monitoring of content uniformity, moisture content, compact density, tensile strength, and young's modulus of roller compacted powder blends. J Pharm Sci, 94, 1589-1597 (2005)
- Han K, Lee H, Kang JH, Choi E, Oh SJ, Lee YJ, Cho BK, Kang BC. A simple method for evaluation of pepper powder color using vis/NIR hyperspectral system. Kor J Hort Sci Technol, 33, 403-408 (2015)
- Hornero-Mendez D, Minguez-Mosquera MI. Rapid spectrophotometric determination of red and yellow isochromic carotenoid fractions in paprika and red pepper oleoresins. J Agric Food Chem, 49, 3584-3588 (2001)
- Iwai K, Suzuki T, Fujiwake H, Oka S. Simultaneous microdetermination of capsaicin and its four analogues by using high performance liquid chromatography and gas chromatography mass spectrometry. J Chromatogr, 172, 303-311 (1979)
- Jha SN, Matsuoka T. Non destructive determination of acid brix ratio of tomato juice using near infrared spectroscopy. Int J Food Sci Technol, 39, 425-430 (2004)
- Kim S, Baek SG, Hwang IJ, Kim SR, Jung G, Roh E, Jang JY, Kim J, Lee T. Fungal occurrence in fresh and dried red pepper. J Food Hyg Saf, 34, 571-575 (2019)
- Kim S, Ha TY, Park J. Characteristics of pigment composition and colour value by the difference of harvesting times in Korean red pepper varieties (*Capsicum annum L.*). Int J Food Sci Technol, 43, 915-920 (2008)
- Kim S, Park JB, Hwang IK. Quality attributes of various varieties of Korean red pepper powders (*Capsicum annum L.*) and color stability during sunlight exposure. Food Chem Toxicol, 67, 2957-2961 (2002)
- Kim SA, Kim KS, Park JB. Changes of various chemical components by the difference of the degree of ripening and harvesting factors in two single harvested peppers (*Capsicum annum*

- L.). Korean J Food Sci Technol, 38, 615-620 (2006)
- Krepper G, Romeo F, Fernandes DDS, Diniz PHGD, Araujo MCU, Nezio MSD, Pistonesi MF, Centurion ME. Determination of fat content in chicken hamburgers using NIR spectroscopy and the successive projections algorithm for interval selection in PLS regression (iSPA-PLS). Spectrochim Acta A Mol Biomol Spectrosc, 189, 300-306 (2018)
- Ku KH, Kim NY, Park JB, Park WS. Characteristics of color and pungency in the red pepper for Kimchi. Korean J Food Sci Technol, 33, 231-237 (2001)
- Ku KH, Park JB, Park WS. Effects of red peppers on the its pungency and color during Kimchi fermentation. J Korean Soc Food Sci Nutr, 33, 1034-1042 (2004)
- Lee KJ, Kang SW, Choi KH. Non destructive quality measurement of fruits and vegetables using near infrared spectroscopy. Food Eng Prog, 8, 158-169 (2004)
- Lim Y, Kyung Y, Jeong HS, Kim HY, Hwang IG, Yoo SM, Lee J. Effects of drying methods on quality of red pepper powder. J Korean Soc Food Sci Nutr, 41, 1315-1319 (2012)
- Mo C, Kang S, Lee K, Lim JG, Cho BK, Lee HD. Development of prediction model for capsaicinoids content in red pepper powder using near infrared spectroscopy particle size effect. Food Eng Prog, 15, 48-55 (2011)
- Mo C, Lee K, Lim JG, Kang S, Lee HD, Cho BK. Development of non destructive pungency measurement technique for red pepper powder produced in different domestic origins. J Agric Sci, 39, 603-612 (2012)
- Oh SH, Hwang IG, Kim HY, Hwang CR, Park SM, Hwang Y, Yoo SM, Kim HR, Kim HY, Lee J, Jeong HS. Quality characteristics by particle size of red pepper powders for pepper paste and kimchi. J Korean Soc Food Sci Nutr, 40, 725-730 (2011)
- Park MJ. Research about ascorbic acid and capsaicin of red pepper from Korea. MS Thesis, Yeungnam University, Gyeongsan, Korea (2004)
- Perucka I, Oleszek W. Extraction and determination of capsaicinoids in fruit of hot pepper (*Capsicum annuum* L.) by spectrophotometry and high-performance liquid chromatography. Food Chem, 71, 287-291 (2000)
- Shao Y, Wang K, Xuan G, Gao C, Hu Z. So soluble solids content monitoring for shelf-life assessment of table grapes coated with chitosan using hyperspectral imaging. Infrared Physics Tech, 115, 103725 (2021)
- Shin HH, Lee SR. Quality attributes of Korean red pepper according to cultivars and growing areas. Korean J Food Sci Technol, 23, 296-300 (1991)
- Yoon WM, Lee JY. Effect of drying method on the fruit and powder color of red pepper. J Natural Sci, 15, 97-109 (2004)