



REVIEW

Effect of Starter Cultures on Quality of Fermented Sausages

Jungeun Hwang¹, Yujin Kim¹, Yeongeun Seo², Miseon Sung¹, Jei Oh¹,
and Yohan Yoon^{1,2,*}

¹Department of Food and Nutrition, Sookmyung Women's University, Seoul 04310, Korea

²Risk Analysis Research Center, Sookmyung Women's University, Seoul 04310, Korea

OPEN ACCESS

Received December 16, 2022

Revised December 22, 2022

Accepted December 22, 2022

*Corresponding author : Yohan Yoon
Department of Food and Nutrition,
Sookmyung Women's University,
Seoul 04310, Korea
Tel: +82-2-2077-7585
Fax: +82-2-710-9479
E-mail: yyoon@sookmyung.ac.kr

*ORCID

Jungeun Hwang
<https://orcid.org/0000-0001-9909-9490>
Yujin Kim
<https://orcid.org/0000-0002-0903-9871>
Yeongeun Seo
<https://orcid.org/0000-0003-4986-9770>
Miseon Sung
<https://orcid.org/0000-0002-1430-692X>
Jei Oh
<https://orcid.org/0000-0002-7215-1495>
Yohan Yoon
<https://orcid.org/0000-0002-4561-6218>

Abstract The expansion and advancement of the meat product market have increased the demand for fermented sausages. A typical method for manufacturing high-quality fermented sausages is using a starter culture, which improves the taste, aroma, and texture. Currently, the starter culture for manufacturing fermented sausages is mainly composed of microorganisms such as lactic acid bacteria, yeast, and fungi, which generate volatile compounds by the oxidation of fatty acids. In addition, protein decomposition and changes in pH occur during the fermentation period. It can positively change the texture of the fermented sausage. In this review, we discuss the requirements (improving food safety, the safety of starter culture, enzyme activity, and color) of microorganisms used in starter cultures and the generation of flavor compounds (heptanal, octanal, nonanal, hexanal, 2-pentylfuran, 1-penten-3-ol, and 2-pentanone) from lipids. Furthermore, quality improvement (hardness and chewiness) due to texture changes after starter culture application during the manufacturing process are discussed.

Keywords starter culture, lactic acid bacteria, quality control, fermented sausages

Introduction

Meat is a rich source of protein, vitamins, and minerals. Due to the perishable nature of meat, it has been processed in various ways for preservation. Sausage fermentation may be one of the earliest forms of meat processing. The meat is preserved long-term by fermentation and drying after filling the casing with a mixture of minced meat, fat, salt, color retention agent, sugar, and spices (Campbell-Platt, 1995). Fermented sausages have been produced in Southern and Central Europe, especially in Germany, Italy, Spain, and France, with the highest production and consumption rates per capita (Lücke, 1998). Fermented sausage production is approximately 200,000 tons annually in Spain and 110,000 tons in France (Safa et al., 2015). In 2014, fermented sausage production in Norway and Finland was 7,300 tons and 7,000 tons, respectively (Holck

et al., 2014). According to the Ministry of Food and Drug Safety, fermented sausage production increased by 77% from 272 tons in 2018 to 614 tons in 2021, and sales during the same period increased by 67% from 266 tons to 445 tons (MFDS, 2019, 2022). However, most fermented sausages in Korea are still imported, even though there is a significant increase in the production of fermented sausages (Jung and Yoon, 2020).

The main ingredients of fermented sausage are meat, fatty tissue, carbohydrates, curing agents, spices, and starter culture (Lücke, 1998). The microorganisms that are primarily involved in sausage fermentation include species of lactic acid bacteria, gram-positive and catalase-positive cocci (GCC), molds, and yeasts, mainly including *Lactobacillus sakei*, *Lactobacillus curvatus*, *Lactobacillus plantarum*, *Lactobacillus brevis*, *Lactobacillus buchneri*, *Lactobacillus paracasei*, *Staphylococcus xylosum*, *Staphylococcus carnosus*, *Staphylococcus saprophyticus*, *Kocuria varians*, *Penicillium nalgiovense*, *Penicillium chrysogenum*, and *Debaryomyces hansenii* (Leroy et al., 2006). During fermentation, the fermented sausage starter cultures should adapt to environmental conditions, such as high salt concentrations and low temperatures (Lee et al., 2006). Moreover, they are required to compete with other microorganisms by rapid growth and pH lowering so that various harmful microorganisms can be suppressed, securing microbiological safety (Lee et al., 2006). Furthermore, the starter cultures should be resistant to sodium-nitrite scavengers and sodium nitrite, which is added for fermented sausage preservation (Leistner, 1995). The starter cultures are responsible for the functions mentioned above and the sensual quality of the fermented sausage. During the short fermentation period, the starter cultures generate distinctive tastes, flavors, and colors of the sausages (Kunz and Lee, 2003). Likewise, the starter cultures play a role in improving the texture and flavor of the fermented sausages.

In this review, the requirement and role of starter cultures in sausage fermentation are intensively reviewed, and fermented sausage quality improvement by starter cultures is discussed.

Requirement for Starter Culture in Fermented Sausages

Improving food safety of fermented sausage

The growth of harmful microorganisms, such as foodborne pathogens, in fermented sausages can be inhibited by applying starter cultures to them, lowering their pH (Laranjo et al., 2019). The starter culture should be able to inhibit toxin production by other microorganisms. *P. nalgiovense*, widely used for starter culture, can inhibit the production of toxin-producing fungi, ensuring the safety of fermented sausages (Bernáldez et al., 2013).

Safety of starter culture

The starter culture should grow well under the physical and chemical conditions of the sausages and be harmless to humans. To ensure the safety of starter cultures, hemolysis, toxin production, and biogenic amine production by the microorganisms in the starter culture must be evaluated (Jeong and Lee, 2015; Sirini et al., 2022). Some starter cultures produce biogenic amines in fermented sausages, and thus, starter cultures producing no biogenic amines need to be developed, and processing methods such as packaging, additives, hydrostatic pressure, and smoking can also reduce biogenic amines (Doeun et al., 2017; Özogul and Hamed, 2018).

Enzyme activity related to sensory property

During the ripening of fermented sausages, enzyme activities derived from starter culture can positively affect the sensual elements of the fermented sausages (Wang et al., 2019). Thus, the microorganisms of the starter culture should be evaluated

for degradation of carbohydrates, proteins, peptides, and lipids to produce a pleasant aroma and taste in fermented sausage.

Maintenance of color

Starter culture plays a role in maintaining the color of fermented sausages by producing nitrate reductase and nitrite reductase, which reduce nitrate and nitrite, respectively. For example, *S. carnosus* has been used to maintain the color of the fermented sausage by forming red nitrosomyoglobin, which is produced by nitrate reductase (Götz, 1990; Löfblom et al., 2017; Neubauer and Götz, 1996).

Changes in Free Fatty Acids and Volatile Compounds by Starter Cultures

Water content and pH in fermented sausages decrease during fermentation, and several biochemical reactions, including lipolysis, occur (Fernández et al., 2000; Lücke, 2000). Lipids in fermented sausages can be oxidized to free fatty acids (Chen et al., 2017; Gandemer, 2002), enhancing the flavor of fermented sausages (Flores and Toldrá, 2011). Lactic acid bacteria, yeasts, and molds are known to hydrolyze triglycerides and increase the free fatty acids content in fermented sausages (Chen et al., 2017; Ordóñez et al., 1999; Sanz et al., 1988). Subsequently, free fatty acids undergo enzymatic and non-enzymatic oxidation, yielding alcohol, aldehydes, carboxylic acids, and other flavor compounds (Casaburi et al., 2007; Fernández et al., 2000; Melgar et al., 1990). Thus, applying microorganisms with high lipolytic activity to the manufacturing process might improve the sensory quality of fermented sausages (Casaburi et al., 2007).

During sausage fermentation, *Pediococcus pentosaceus* and *S. xylosus* increased the contents of total free fatty acids from 0.6% to 6.8% (Johansson et al., 1994). Karsloğlu et al. (2014) investigated free fatty acid contents and lipolytic changes in fermented sausages inoculated with commercial starter cultures, including *Lactobacillus sakei*, *S. carnosus*, *S. xylosus*, and *P. pentosaceus*. Oleic acid, linoleic acid, linolenic acid, and palmitic acid were the predominant fatty acids in the fermented sausage stored for 120 days. The oleic acid content had increased from 1.77%–1.99% to 6.56%–13.01%. Other researchers compared the changes in saturated fatty acid (SFA), monounsaturated fatty acid (MUFA), and polyunsaturated fatty acid (PUFA) levels in fermented sausages. When fermented sausages were inoculated with *L. sakei* and *S. xylosus*, the final SFA and PUFA contents increased, whereas the MUFA content was lower than that in sausages without a starter culture (Hu et al., 2007). In boar sausages, *D. hansenii* increased MUFA and PUFA contents, and it supports the lipolysis in fermented sausages (Patrignani et al., 2007). All these findings indicate that the use of starter culture contributes degradation of lipids in fermented sausages.

Volatile compounds may be produced from the degradation of lipid in fermented sausages when starter cultures are used (Table 1). The inoculation of *D. hansenii* in *Salchichón* changed the volatile compound profile at 54 days of fermentation, and the inoculated sausages had higher volatile compounds (58 volatile compounds) than those (41 volatile compounds) in the uninoculated sausages. Moreover, heptanal, octanal, and nonanal, known as cured-ham-like flavor compounds, were only detected in the inoculated batches (Andrade et al., 2010; Carrapiso et al., 2002). Corral et al. (2015) manufactured fermented sausages with low fat and salt content, and inoculated with *D. hansenii* in the fermented sausages. Compared to the uninoculated fermented sausages, the sausages with *D. hansenii* showed a higher content of 2-pentylfuran, which has a metallic odor. When fermented sausages were inoculated with *L. sakei* and combinations of different *Staphylococcus* species, the changes in volatile compounds differed with each *Staphylococcus* strain. In addition, only hexanal significantly decreased with the addition of starter cultures, reflecting decreased rancid flavor in the fermented sausages (Fonseca et al., 2013).

Few studies analyzed volatile compounds in fermented sausages inoculated with mold starters. Bruna et al. (2001)

Table 1. Changes in contents of volatile compounds in the sausages inoculated with starter cultures compared to the uninoculated sausages

Starter culture	Major volatile compounds	Minor volatile compounds	References
<i>Debaryomyces hansenii</i>	Decane, 1-propanol, 1-butanol, 2-propanone, 2-pentanone, 2-heptanone, pentanal, heptanal, octanal, nonanal, propanoic acid, hexanoic acid, hexadecenoic acid	Hexane, 2-methylpentane, 3-methylpentane	Andrade et al. (2010)
	Propanal, 1-propanol, 2-methylfuran, 2-pentylfuran, hexanoic acid, 2-ethyl-1-hexanol, decanoic acid	Tetrahydrofuran, 2,5-dimethylfuran, heptanoic acid	Corral et al. (2015)
<i>Lactobacillus sakei</i> , <i>Staphylococcus equorum</i>	Hexane, nonanoic acid	Octane, hexanal, 3-ethyl-heptane, nonane, 3-ethyl-4-methyl-hexane, 3-methyl-nonane, 2,2,6-trimethyl-octane, decane, 3,7-dimethyl-nonane, 3,7-dimethyl-decane, undecane, 3,3-dimethyl-hexane, 2,2,5-trimethyl-decane, 5-ethyl-5-methyl-decane, dodecane, tridecane	Fonseca et al. (2013)
<i>L. sakei</i> , <i>Staphylococcus epidermidis</i>	Hexane, 2,2,6-trimethyl-octane, 5-ethyl-5-methyl-decane, nonanoic acid	Octane, hexanal, 3-ethyl-heptane, nonane, 3-ethyl-4-methyl-hexane, 3-methyl-nonane, decane, 3,7-dimethyl-nonane, 3,7-dimethyl-decane, undecane, 3,3-dimethyl-hexane, 2,2,5-trimethyl-decane, dodecane, tridecane	
<i>L. sakei</i> , <i>Staphylococcus saprophyticus</i>	Hexane, 3-ethyl-heptane, 3-ethyl-4-methyl-hexane, 3-methyl-nonane, decane, 3,7-dimethyl-decane, 5-ethyl-5-methyl-decane	Octane, hexanal, nonane, 2,2,6-trimethyl-octane, 3,7-dimethyl-nonane, undecane, 3,3-dimethyl-hexane, 2,2,5-trimethyl-decane, dodecane, tridecane, nonanoic acid	
<i>Penicillium aurantiogriseum</i> (intracellular cell free extract)	2-Butanone, 2-methylfuran, 2-ethylfuran	1-Penten-3-ol, 2-pentanone	Bruna et al. (2001)
<i>P. aurantiogriseum</i> (spore suspension)	-	1-Penten-3-ol, 2-butenal, pentanal, hexanal, 2-hexanal, heptanal, nonanal, decanal, 2-butanone, 2-pentanone, 2-heptanone, 2-methylfuran	
<i>P. aurantiogriseum</i> (intracellular cell free extract, spore suspension)	-	1-Penten-3-ol, 2-butenal, pentanal, hexanal, 2-hexanal, heptanal, nonanal, decanal, 2-butanone, 2-pentanone, 2-heptanone, 2-methylfuran	

analyzed the difference in volatile compounds between the inoculation of *Penicillium aurantiogriseum* and the intracellular cell free extract of *P. aurantiogriseum*. The volatile compounds such as 1-penten-3-ol and 2-pentanone, which have unpleasant flavor, were significantly lower in inoculated samples than in uninoculated samples, regardless of the inocula type. On the other hand, no significant differences were observed in the free fatty acids profile between starter-culture-inoculated and uninoculated fermented sausages (Corral et al., 2015; Lorenzo et al., 2014). The different lipolytic activities might cause this among strains (Patrignani et al., 2007). These results suggest that using starter cultures increases free fatty acid contents and positively affects the production of volatile compounds in fermented sausages, which improves the flavor of fermented sausage (Fig. 1).

Changes in Texture Profiles of Fermented-Sausage

The physicochemical characteristics of fermented sausages can be evaluated during ripening or storage. One of the




Fig. 1. Effects of starter culture on quality improvement of fermented sausages.

characteristics is the texture profile, which includes cohesiveness, hardness, or chewiness. The protein structure of the fermented sausage is responsible for the proper texture formation after salting, chopping, and fermentation (Katsaras and Budras, 1992). Texture formation is closely associated with the reduction of pH in fermented sausages. During ripening, the protein structure of the fermented sausage is denatured by organic acids such as lactic acid, and the moisture content decreases gradually. Moreover, unstable coagulation bonds in the protein structure are substituted with condensed bonds, which result in the transformation from the sol state to the gel state (Katsaras and Budras, 1992). Bozkurt and Bayram (2006), Casaburi et al. (2007), and Gonzalez-Fernandez et al. (2006) confirmed that decreases in pH were negatively correlated with hardness and chewiness. Thus, when the pH and moisture of the sausages decrease, muscle fiber proteins are accumulated, which induces the gel structure. This gel structure increases the hardness and elasticity of the sausages (Essid and Hassouna, 2013; Katsaras and Budras, 1992; Fig. 1 and Table 2).

Proteolysis is one of the major phenomena that occurs during the ripening of fermented sausages, and it needs endogenous enzymes in meat and microbial enzymes (Dalmış and Soyer, 2008; Roseiro et al., 2008). Proteins degrade into small molecules such as peptides, amines, aldehydes, or amino acids by proteolysis, contributing to the final texture properties and flavor of the fermented sausages (Dalmış and Soyer, 2008; Roseiro et al., 2008; Spaziani et al., 2009).

Benito et al. (2005) showed that the hardness of fermented sausages decreased because of the proteolysis by the fungal protease in the starter culture. Aro et al. (2010) compared proteolysis among fermented sausages inoculated with no starter culture, a mixture of *L. sakei* and *S. carnosus*, and a mixture of *P. pentosaceus* and *S. xylosum*. The result showed that the microorganisms in the starter cultures broke down myofibrillar and sarcoplasmic proteins into short peptides and other small molecules. Nie et al. (2014) used *L. plantarum* or *P. pentosaceus* in fermented sausages, and free amino acid concentrations

Table 2. Correlation between physico-chemical properties and texture

Physico-chemical property	Changes in texture	References
Decrease in pH and water content	Negatively correlated with hardness and chewiness	Bozkurt and Bayram (2006); Casaburi et al. (2007); Gonzalez-Fernandez et al. (2006)
Proteolysis	Formation of flavor and final texture of fermented sausages.	Dalmış and Soyer (2008); Roseiro et al. (2008)

increased with adding starter culture. Hu et al. (2022) reported that the shear force of dry fermented sausages increased significantly in the samples inoculated with *L. plantarum* compared to those inoculated with *L. sakei*, *L. culvatus*, and *Weissella hellenica*. Wang et al. (2021) used *L. sakei*, a mixture of *L. sakei* and *S. xylosum*, and a mixture of *L. sakei*, *S. xylosum* and *S. carnosus* in fermented sausages. The result from this study implies that amino acid contents were higher in the sausages inoculated with mixed cultures than those inoculated with a single strain. Ayyash et al. (2020) compared the hardness and proteolysis effect of the dried fermented camel sausage and dried fermented beef sausage inoculated with the same starter culture. The result showed that the hardness of dried fermented camel sausages was significantly lower than that of beef. However, the proteolysis level of the fermented camel sausages was observed to be greater than that of beef. These results indicate that the proteolysis activity in fermented sausages may depend on the starter culture, and the mixed starter culture may cause higher activity than a single strain. In addition, the proteolysis activity may depend on the type of meat used in fermented sausages.

Conclusion

A starter culture in a fermented sausage should be able to inhibit the growth of harmful microorganisms, and the starter culture should be safe for humans. The starter culture plays a role in maintaining red color, breaking lipids into free fatty acid, and producing volatile compounds, which improve the flavor of fermented sausages. In addition, the texture profile of fermented sausages is affected by the proteolysis activity of the starter culture.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This research was funded by Rural Development Administration in Korea (grant number: PJ0155902022).

Author Contributions

Conceptualization: Hwang J, Yoon Y. Data curation: Hwang J, Kim Y, Seo Y, Sung M, Oh J. Formal analysis: Hwang J. Software: Hwang J, Kim Y, Seo Y, Sung M, Oh J. Validation: Kim Y, Seo Y. Writing - original draft: Hwang J, Kim Y, Seo Y, Sung M, Oh J. Writing - review & editing: Hwang J, Kim Y, Seo Y, Sung M, Oh J, Yoon Y.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

Andrade MJ, Córdoba JJ, Casado EM, Córdoba MG, Rodríguez M. 2010. Effect of selected strains of *Debaryomyces hansenii* on the volatile compound production of dry fermented sausage “salchichón”. Meat Sci 85:256-264.

- Aro JMA, Nyam-Osor P, Tsuji K, Shimada K, Fukushima M, Sekikawa M. 2010. The effect of starter cultures on proteolytic changes and amino acid content in fermented sausages. *Food Chem* 119:279-285.
- Ayyash M, Olaimat A, Al-Nabulsi A, Liu SQ. 2020. Bioactive properties of novel probiotic *Lactococcus lactis* fermented camel sausages: Cytotoxicity, angiotensin converting enzyme inhibition, antioxidant capacity, and antidiabetic activity. *Food Sci Anim Resour* 40:155-171.
- Benito MJ, Rodríguez M, Córdoba MG, Andrade MJ, Córdoba JJ. 2005. Effect of the fungal protease EPg222 on proteolysis and texture in the dry fermented sausage 'salchichón'. *J Sci Food Agric* 85:273-280.
- Bernáldez V, Córdoba JJ, Rodríguez M, Cordero M, Polo L, Rodríguez A. 2013. Effect of *Penicillium nalgiovense* as protective culture in processing of dry-fermented sausage "salchichón". *Food Control* 32:69-76.
- Bozkurt H, Bayram M. 2006. Colour and textural attributes of sucuk during ripening. *Meat Sci* 73:344-350.
- Bruna JM, Hierro EM, de la Hoz L, Mottram DS, Fernández M, Ordóñez JA. 2001. The contribution of *Penicillium aurantiogriseum* to the volatile composition and sensory quality of dry fermented sausages. *Meat Sci* 59:97-107.
- Campbell-Platt G. 1995. Fermented meats: A world perspective. In *Fermented meats*. Campbell-Platt G, Cook PE (ed). Springer, Boston, MA, USA. pp 39-52.
- Carrapiso AI, Ventanas J, García C. 2002. Characterization of the most odor-active compounds of Iberian ham headspace. *J Agric Food Chem* 50:1996-2000.
- Casaburi A, Aristoy MC, Cavella S, Di Monaco R, Ercolini D, Toldrá F, Villani F. 2007. Biochemical and sensory characteristics of traditional fermented sausages of Vallo di Diano (Southern Italy) as affected by the use of starter cultures. *Meat Sci* 76:295-307.
- Chen Q, Kong B, Han Q, Xia X, Xu L. 2017. The role of bacterial fermentation in lipolysis and lipid oxidation in Harbin dry sausages and its flavour development. *LWT-Food Sci Technol* 77:389-396.
- Corral S, Salvador A, Belloch C, Flores M. 2015. Improvement the aroma of reduced fat and salt fermented sausages by *Debaryomyces hansenii* inoculation. *Food Control* 47:526-535.
- Dalmış Ü, Soyer A. 2008. Effect of processing methods and starter culture (*Staphylococcus xylosum* and *Pediococcus pentosaceus*) on proteolytic changes in Turkish sausages (sucuk) during ripening and storage. *Meat Sci* 80:345-354.
- Doeun D, Davaatseren M, Chung MS. 2017. Biogenic amines in foods. *Food Sci Biotechnol* 26:1463-1474.
- Essid I, Hassouna M. 2013. Effect of inoculation of selected *Staphylococcus xylosum* and *Lactobacillus plantarum* strains on biochemical, microbiological and textural characteristics of a Tunisian dry fermented sausage. *Food Control* 32:707-714.
- Fernández M, Ordóñez JA, Bruna JM, Herranz B, de la Hoz L. 2000. Accelerated ripening of dry fermented sausages. *Trends Food Sci Technol* 11:201-209.
- Flores M, Toldrá F. 2011. Microbial enzymatic activities for improved fermented meats. *Trends Food Sci Technol* 22:81-90.
- Fonseca S, Cachaldora A, Gómez M, Franco I, Carballo J. 2013. Effect of different autochthonous starter cultures on the volatile compounds profile and sensory properties of Galician chorizo, a traditional Spanish dry fermented sausage. *Food Control* 33:6-14.
- Gandemer G. 2002. Lipids in muscles and adipose tissues, changes during processing and sensory properties of meat products. *Meat Sci* 62:309-321.
- Gonzalez-Fernández C, Santos EM, Rovira J, Jaime I. 2006. The effect of sugar concentration and starter culture on instrumental and sensory textural properties of chorizo-Spanish dry-cured sausage. *Meat Sci* 74:467-475.
- Götz F. 1990. *Staphylococcus carnosus*: A new host organism for gene cloning and protein production. *J Appl Bacteriol*

Symp Suppl 69:49S-53S.

- Holck A, Heir E, Johannessen TC, Axelsson L. 2014. Northern European products. In Handbook of fermented meat and poultry. Toldrá F, Hui YH, Astiasarán I, Sebranek JG, Talon R (ed). John Wiley & Sons, Hoboken, NJ, USA. pp 313-320.
- Hu Y, Li Y, Li X, Zhang H, Chen Q, Kong B. 2022. Application of lactic acid bacteria for improving the quality of reduced-salt dry fermented sausage: Texture, color, and flavor profiles. LWT-Food Sci Technol 154:112723.
- Hu Y, Xia W, Ge C. 2007. Effect of mixed starter cultures fermentation on the characteristics of silver carp sausages. World J Microbiol Biotechnol 23:1021-1031.
- Jeong DW, Lee JH. 2015. Antibiotic resistance, hemolysis and biogenic amine production assessments of *Leuconostoc* and *Weissella* isolates for kimchi starter development. LWT-Food Sci Technol 64:1078-1084.
- Johansson G, Berdagüé JL, Larsson M, Tran N, Borch E. 1994. Lipolysis, proteolysis and formation of volatile components during ripening of a fermented sausage with *Pediococcus pentosaceus* and *Staphylococcus xylosus* as starter cultures. Meat Sci 38:203-218.
- Jung YS, Yoon HH. 2020. Quantitative descriptive analysis and consumer acceptance of commercial dry fermented sausages. J East Asian Soc Diet Life 30:306-315.
- Karsloğlu B, Çiçek ÜE, Kolsarici N, Candoğan K. 2014. Lipolytic changes in fermented sausages produced with turkey meat: Effects of starter culture and heat treatment. Korean J Food Sci Anim Resour 34:40-48.
- Katsaras K, Budras KD. 1992. Microstructure of fermented sausage. Meat Sci 31:121-134.
- Kunz B, Lee JY. 2003. Production and microbiological characteristics of fermented sausages. Korean J Food Sci Anim Resour 23:361-375.
- Laranjo M, Potes ME, Elias M. 2019. Role of starter cultures on the safety of fermented meat products. Front Microbiol 10:853.
- Lee JY, Kim CJ, Kunz B. 2006. Identification of lactic acid bacteria isolated from *kimchi* and studies on their suitability for application as starter culture in the production of fermented sausages. Meat Sci 72:437-445.
- Leistner L. 1995. Stable and safe fermented sausages world-wide. In Fermented meats. Campbell-Platt G, Cook PE (ed). Springer, Boston, MA, USA. pp 160-175.
- Leroy F, Verluyten J, De Vuyst L. 2006. Functional meat starter cultures for improved sausage fermentation. Int J Food Microbiol 106:270-285.
- Löfblom J, Rosenstein R, Nguyen MT, Ståhl S, Götz F. 2017. *Staphylococcus carnosus*: From starter culture to protein engineering platform. Appl Microbiol Biotechnol 101:8293-8307.
- Lorenzo JM, Gómez M, Fonseca S. 2014. Effect of commercial starter cultures on physicochemical characteristics, microbial counts and free fatty acid composition of dry-cured foal sausage. Food Control 46:382-389.
- Lücke FK. 1998. Fermented sausages. In Microbiology of fermented foods. Wood BJB (ed). Springer, Boston, MA, USA. pp 441-483.
- Lücke FK. 2000. Utilization of microbes to process and preserve meat. Meat Sci 56:105-115.
- Melgar MJ, Sanchez-Monge JM, Bello J. 1990. A study of the changes in the chemical properties of fat during ripening of dry Spanish sausage. J Food Compos Anal 3:73-80.
- Ministry of Food and Drug Safety [MFDS]. 2019. Production performance statistics of food, etc. in 2018. Available from: https://www.mfds.go.kr/brd/m_374/view.do?seq=30198&srchFr=&srchTo=&srchWord=&srchTp=&itm_seq_1=0&itm_s

- eq_2=0&multi_itm_seq=0&company_cd=&company_nm=&page=1. Accessed at Dec 14, 2022.
- Ministry of Food and Drug Safety [MFDS]. 2022. Production performance statistics of food, etc. in 2021. Available from: https://www.mfds.go.kr/brd/m_374/view.do?seq=30207&srchFr=&srchTo=&srchWord=&srchTp=&itm_seq_1=0&itm_seq_2=0&multi_itm_seq=0&company_cd=&company_nm=&page=1. Accessed at Dec 14, 2022.
- Neubauer H, Götz F. 1996. Physiology and interaction of nitrate and nitrite reduction in *Staphylococcus carnosus*. J Bacteriol 178:2005-2009.
- Nie X, Lin S, Zhang Q. 2014. Proteolytic characterisation in grass carp sausage inoculated with *Lactobacillus plantarum* and *Pediococcus pentosaceus*. Food Chem 145:840-844.
- Ordóñez JA, Hierro EM, Bruna JM, de la Hoz L. 1999. Changes in the components of dry-fermented sausages during ripening. Crit Rev Food Sci Nutr 39:329-367.
- Özogul F, Hamed I. 2018. The importance of lactic acid bacteria for the prevention of bacterial growth and their biogenic amines formation: A review. Crit Rev Food Sci Nutr 58:1660-1670.
- Patrignani F, Iucci L, Vallicelli M, Guerzoni ME, Gardini F, Lanciotti R. 2007. Role of surface-inoculated *Debaryomyces hansenii* and *Yarrowia lipolytica* strains in dried fermented sausage manufacture. Part 1: Evaluation of their effects on microbial evolution, lipolytic and proteolytic patterns. Meat Sci 75:676-686.
- Roseiro LC, Santos C, Sol M, Borges MJ, Anjos M, Gonçalves H, Carvalho AS. 2008. Proteolysis in *Painho de Portalegre* dry fermented sausage in relation to ripening time and salt content. Meat Sci 79:784-794.
- Safa H, Gatellier P, Lebert A, Picgirard L, Mirade PS. 2015. Effect of combined salt and animal fat reductions on physicochemical and biochemical changes during the manufacture of dry-fermented sausages. Food Bioproc Technol 8:2109-2122.
- Sanz B, Selgas D, Parejo I, Ordóñez JA. 1988. Characteristics of lactobacilli isolated from dry fermented sausages. Int J Food Microbiol 6:199-205.
- Sirini N, Munekata PES, Lorenzo JM, Stegmayer MÁ, Pateiro M, Pérez-Álvarez JÁ, Sepúlveda N, Sosa-Morales ME, Teixeira A, Fernández-López J, Frizzo L, Rosmini M. 2022. Development of healthier and functional dry fermented sausages: Present and future. Foods 11:1128.
- Spaziani M, Del Torre M, Stecchini ML. 2009. Changes of physicochemical, microbiological, and textural properties during ripening of Italian low-acid sausages. Proteolysis, sensory and volatile profiles. Meat Sci 81:77-85.
- Wang D, Hu G, Wang H, Wang L, Zhang Y, Zou Y, Zhao L, Liu F, Jin Y. 2021. Effect of mixed starters on proteolysis and formation of biogenic amines in dry fermented mutton sausages. Foods 10:2939.
- Wang D, Zhao L, Su R, Jin Y. 2019. Effects of different starter culture combinations on microbial counts and physicochemical properties in dry fermented mutton sausages. Food Sci Nutr 7:1957-1968.