



ARTICLE

Bactericidal Effect of Combination of Atmospheric Pressure Plasma and Nisin on Meat Products Inoculated with *Escherichia coli* O157:H7

Hag Ju Lee¹, Yeseul Heo¹, Hye-Jin Kim¹, Ki Ho Baek², Dong-Gyun Yim¹, Anand Kumar Sethukali¹, Dongbin Park¹, and Cheorun Jo^{1,3,4,*}

¹Department of Agricultural Biotechnology, Center for Food and Bioconvergence, and Research Institute of Agriculture and Life Science, Seoul National University, Seoul 08826, Korea

²Department of Nano-Bio Convergence, Korea Institute of Materials Science, Changwon 51508, Korea

³Institute of Green Bio Science and Technology, Seoul National University, Pyeongchang 25354, Korea

⁴Department of Animal Product Technology, Faculty of Animal Husbandry, Universitas Padjadjaran, West Java 45363, Indonesia

OPEN ACCESS

Received August 31, 2022

Revised October 12, 2022

Accepted November 29, 2022

*Corresponding author : Cheorun Jo
Department of Agricultural Biotechnology,
Center for Food and Bioconvergence, and
Research Institute of Agriculture and Life
Science, Seoul National University, Seoul
08826, Korea
Tel: +82-2-880-4820
Fax: +82-2-873-2271
E-mail: cheorun@snu.ac.kr

*ORCID
Hag Ju Lee
<https://orcid.org/0000-0003-2906-7666>
Yeseul Heo
<https://orcid.org/0000-0001-8161-778X>
Hye-Jin Kim
<https://orcid.org/0000-0002-9384-6720>
Ki Ho Baek
<https://orcid.org/0000-0002-5438-9547>
Dong-Gyun Yim
<https://orcid.org/0000-0003-0368-2847>
Anand Kumar Sethukali
<https://orcid.org/0000-0003-0817-6396>
Dongbin Park
<https://orcid.org/0000-0003-4979-6049>
Cheorun Jo
<https://orcid.org/0000-0003-2109-3798>

Abstract This study was conducted to investigate the bactericidal effect of nisin (Nisin) only, atmospheric pressure plasma (APP) only, and a combination of APP and nisin (APP+Nisin) on beef jerky and sliced ham inoculated with *Escherichia coli* O157:H7, gram-negative bacteria. The bactericidal effect against *E. coli* O157:H7 and *Listeria monocytogenes* was confirmed using a nisin solution at a concentration of 0–100 ppm, and APP+Nisin was tested on beef jerky and sliced ham using 100 ppm nisin. Beef jerky and sliced ham were treated with APP for 5 min and 9 min, respectively. In the bacterial solution, 100 ppm nisin out of 0–100 ppm nisin exhibited the highest bactericidal activity against *L. monocytogenes* (gram-positive bacteria; $p < 0.05$); however, it did not exhibit bactericidal effects against *E. coli* O157:H7 (gram-negative bacteria). The APP+Nisin exhibited a 100% reduction rate in both *E. coli* O157:H7 and *L. monocytogenes* compared to the control group, and was more effective than the Nisin. The APP+Nisin decreased the number of colonies formed by 0.80 and 1.96 Log CFU/g for beef jerky and sliced ham, respectively, compared to the control, and exhibited a higher bactericidal effect compared to the Nisin ($p < 0.05$). These results demonstrate the synergistic bactericidal effect of APP and nisin, providing a possible method to improve the limitations of nisin against gram-negative bacteria. In addition, this technology has the potential to be applied to various meats and meat products to control surface microorganisms.

Keywords atmospheric pressure plasma, nisin, gram-negative bacteria, meat products, bactericidal effect

Introduction

Meat consumption per capita and total meat consumption are increasing worldwide

(Baek et al., 2020b). Meat and meat products are a good environment for the growth of microorganisms (Heredia and García, 2018), and can be easily contaminated with food-borne pathogens during their manufacture, storage, and distribution (Yoo et al., 2021). Contamination of meats can cause diseases such as food poisoning, which is caused by the ingestion of pathogens and microbial toxins. The symptoms of food poisoning include abdominal pain, diarrhea, vomiting, and fever; in severe cases, it can even cause death (Gourama, 2020). Therefore, to consume meat products safely, food safety must be ensured, for which technology to effectively control microorganisms is required.

Thermal treatment is one of the representative technologies for inactivating foodborne pathogens and ensuring food safety. However, it degrades the nutritional and sensory qualities of food (Heo et al., 2021). Therefore, there is a need for technology that can efficiently control pathogenic bacteria and minimize the negative effects on the nutritional and sensory quality of meat products. Non-thermal treatment, unlike thermal technology, can minimize the nutritional and sensory quality degradation of food (Jadhav et al., 2021). Non-thermal technologies include physical techniques, such as ultrasonication, plasma technology, irradiation, pulsed electric field, and high-pressure processing (Jadhav et al., 2021; Osae et al., 2020), as well as chemical methods, such as the use of chlorine-based sanitizers, organic acids, and bacteriocins (Demirok et al., 2013; Mani-López et al., 2012; Min et al., 2007).

Among these technologies, nisin is a peptide produced by a specific strain of *Lactococcus lactis*, which has been designated generally recognized as safe by the Food and Drug Administration (FDA). It exhibits microbial inactivation properties, and is the most commonly used bacteriocin (Zhao et al., 2020). Nisin causes leakage of intracellular bacterial components, leading to the death of bacterial cells (Tong et al., 2014). It exhibits high antibacterial activity against gram-positive bacteria. However, the lipopolysaccharide layer in the outer membrane of gram-negative bacteria confers resistance to nisin, limiting its efficacy against gram-negative bacteria (Liang et al., 2020). This limitation can be overcome by combining it with other technologies that can destroy the outer membrane (Wang et al., 2018). Some studies have been conducted to improve the antibacterial activity of nisin against gram-negative bacteria by damaging the integrity of the outer membrane in combination with an antibacterial agent (Zhao et al., 2020) or physical technology, such as electric field, high-pressure processing, ultrasound, and plasma (Costello et al., 2021; Gallo et al., 2007; Li et al., 2016; Liao et al., 2018; López-Pedemonte et al., 2003; Mok et al., 2020; Novickij et al., 2018; Pokhrel et al., 2019; Ukuku et al., 2019).

Cold atmospheric pressure plasma (APP) is a non-thermal sterilization technology that use ionized gas under quasi-neutral conditions (Lee et al., 2011). According to Yong et al. (2015), it can be seen that plasma can effectively inactivate *Escherichia coli* (*E. coli*), *Salmonella Typhimurium*, and *Listeria monocytogenes* (*L. monocytogenes*). APP generates various UV photons, electrons, positive and negative ions, atomic species, and free radicals (Kim et al., 2020). In particular, reactive species such as superoxide anions, hydroxyl radicals, nitric oxide, and ozone generated through plasma (Yong et al., 2015) cause cell membrane damage via physical and chemical changes, such as lipid peroxidation of cell membranes, which ultimately induces cell leakage and lead to cell death (Zhao et al., 2022). In addition, owing to structural differences, such as the peptidoglycan layer of gram-positive and gram-negative bacteria, APP exhibits a greater bactericidal effect against gram-negative bacteria than gram-positive bacteria (Yoo et al., 2021). Therefore, the bactericidal effect of APP seems to overcome the problems related to the resistance of gram-negative bacteria to nisin. In addition, there are few studies on improving the bactericidal effect of the combination of plasma and nisin treatment (Costello et al., 2021; Ukuku et al., 2019). However, this phenomenon needs to be investigated in more detail.

Therefore, in this study, combined treatment with APP and nisin was performed to determine their bactericidal effect on meat products. To achieve this, the effect of APP and nisin combination on the reduction of *E. coli* O157:H7 and *L.*

monocytogenes was evaluated. The APP and nisin combination was then applied beef jerky and sliced ham inoculated with *E. coli* O157:H7 to confirm its bactericidal effect.

Materials and Methods

Encapsulated atmospheric pressure plasma (APP) system

The encapsulated atmospheric pressure dielectric barrier discharge plasma described by Kang et al. (2022) was used. Briefly, the plasma device was fabricated by attaching copper electrodes and a polytetrafluoroethylene sheet to a rectangular plastic container (137×104×53 mm) and a lid. The sample (bacterial solution or inoculated meat product) was placed in a glass dish in the plasma device, and discharge was performed at 2.2 kHz and 8.4 kV using atmospheric air.

Bacterial strains and culture preparation

E. coli O157:H7 (NCCP 15739) and *L. monocytogenes* (ATCC 19111) were obtained from the National Culture Collection for Pathogens (Cheongju, Korea) and Korean Culture Center of Microorganisms (Seoul, Korea), respectively. Single colonies of *E. coli* O157:H7 and *L. monocytogenes* were transferred into 25 mL tryptic soy broth (TSB) and TSB containing yeast extract (YE), respectively, and cultured twice at 37°C for 24 and 18 h using orbital agitation at 120 rpm. Then, each broth was transferred to a 50 mL centrifuge tube and centrifuged at 4,001×g at 4°C for 10 min. The supernatant was discarded and cell pellet was completely diluted with 0.85% NaCl to adjust the initial concentration of each bacterial strain from 10⁷ to 10⁸ CFU/mL by measuring the optical density at 600 nm (OD₆₀₀=0.2).

Microbial analysis

To quantify the number of bacteria in samples, serial dilutions were performed using 0.85% NaCl. After mixing reacting the bacterial solution with the treatment solution, 1 mL of the solution was transferred to a glass tube containing 9 mL of 0.85% NaCl, followed by serial dilution. The appropriately diluted solution was spread on tryptic soy agar (TSA) and TSA+YE to count *Escherichia coli* O157:H7 and *L. monocytogenes*, respectively. The media was incubated at 37°C for 48 h. The number of colonies is expressed as Log CFU/mL or g.

Effect of nisin on the bacterial solution

A stock solution (2,500 ppm) of nisin (N5764-5G, Sigma-Aldrich, St. Louis, MO, USA) was prepared by dissolving 125 mg of nisin in 50 mL of 20 mM hydrochloric acid (7647-01-0, Duksan Pure Chem, Ansan, Korea). The stock solution was diluted with deionized (DI) water to prepare 25, 50, and 100 ppm nisin solutions. Then, 2.7 mL of each concentration of nisin solution and 0.3 mL of the bacterial solution were placed in a centrifuge tube. The centrifuge tube was vortexed for 5 s and incubated for 30 min at room temperature (25±2°C). For control, 2.7 mL DI water and 0.3 mL of the bacterial solution were used. In all treatment groups, the mixture of 2.7 mL treatment solution and 0.3 mL bacterial solution was considered as the 10⁰ state. Microbial analysis was performed after dilution with 0.85% NaCl to an appropriate concentration.

Synergistic effect of atmospheric pressure plasma (APP) and nisin on the bacterial solution

The scheme for preparing nisin (Nisin) only, APP only, and combination of APP and nisin (APP+Nisin) is shown in Fig. 1A. Control and Nisin were prepared using a bacterial solution (0.3 mL) with DI water (2.7 mL) and 100 ppm nisin solution (2.7




Fig. 1. Schematic diagram of bactericidal test treatment for bacterial solution (A) and inoculated meat products (B). DI water, deionized water; APP, cold atmospheric pressure plasma.

mL), respectively. For APP, DI water (2.7 mL) was added to the APP-treated bacterial solution (0.3 mL), which was discharged for 5 min. To determine the synergistic effect of APP+Nisin, APP was treated to the bacterial solution (0.3 mL), and 100 ppm nisin solution (2.7 mL) was added. All treated solutions were transferred to a 50 mL centrifuge tube, vortexed for 5 s, and

incubated for 30 min at room temperature ($25\pm 2^\circ\text{C}$) to match the reaction time with the Nisin. The solution in the centrifuge tube was regarded as 10^0 state. Microbial analysis was performed as described above after dilution with 0.85% NaCl to an appropriate concentration.

Effect of atmospheric pressure plasma (APP) and nisin combination treatment on *Escherichia coli* O157:H7 inoculated beef jerky and sliced ham

Beef jerky and sliced ham were purchased from a local market (Seoul, Korea). All samples were exposed to UV light for 30 min to remove surface microorganisms prior to cutting the samples and inoculating them with *E. coli* O157:H7. Beef jerky and sliced ham were cut into 5 g (35×35 mm, $n=2$) and 3 g (40×40 mm, $n=3$) pieces, respectively, and 100 μL and 50 μL of *E. coli* O157:H7 ($\text{OD}_{600}=0.20$), respectively, were spot-inoculated on the surface. The inoculated samples were then dried at room temperature ($25\pm 2^\circ\text{C}$) for 30 min. The scheme of preparing the Nisin, APP, and APP+Nisin for inoculated beef jerky and sliced ham is shown in Fig. 1B. The Nisin involved dipping the samples in nisin solution (100 ppm) for 30 min. For the APP, beef jerky and sliced ham were treated with APP for 5 and 9 min, respectively, and immersed in 45 and 27 mL of DI water, respectively. They were then vortexed for 5 s and allowed to react for 30 min at room temperature ($25\pm 2^\circ\text{C}$). For APP+Nisin, APP was directly applied to the beef jerky and sliced ham for 5 min and 9 min, respectively. Both samples were then dipped in 45 and 27 mL of nisin solution (100 ppm), respectively, vortexed for 5 s, and incubated at room temperature ($25\pm 2^\circ\text{C}$) for 30 min. After completion of the reaction, the beef jerky and sliced ham were transferred to a centrifuge tube containing 0.85% NaCl and vortexed for 2 min to detach the bacterial strain present in the sample, which was considered as the 10^0 state. Microbial analysis was performed as described above.

Statistical analysis

All the experiments were performed in triplicate, except for those involving beef jerky, which was replicated. All data were assessed using SAS (version 9.4, SAS Institute, Cary, NC, USA) with statistical significance set at $p<0.05$. Statistical analysis was conducted using one-way analysis of variance and Tukey's multiple comparison test.

Results and Discussion

Effect of nisin on reduction of *Escherichia coli* O157:H7 and *Listeria monocytogenes*

Fig. 2A shows the bactericidal effect of nisin on *E. coli* O157:H7 and *L. monocytogenes*. The initial count of *E. coli* O157:H7 was 7.11 Log CFU/mL. The number of viable *E. coli* O157:H7 cells at 25, 50, and 100 ppm nisin decreased by 0.04, 0.04, and 0.01 Log CFU/mL, respectively, compared with that at 0 ppm nisin. However, this difference was not statistically significant. This suggests that nisin concentration below 100 ppm has no bactericidal effect on *E. coli* O157:H7.

The initial count of *L. monocytogenes* was 7.66 Log CFU/mL. The number of viable *L. monocytogenes* cells at 50 and 100 ppm nisin decreased by 0.09, and 0.24 Log CFU/mL, respectively, compared to that at 0 ppm nisin ($p<0.05$). In addition, the 100 ppm treatment had the highest bactericidal effect among the nisin concentrations tested ($p<0.05$). When compared the result of the bactericidal effect of nisin at 100 ppm on *E. coli* O157:H7 and *L. monocytogenes*, there was a higher reduction rate in *L. monocytogenes* than in *E. coli* O157:H7 (3.13% vs. 0.14%). This result confirmed that nisin has a lower bactericidal effect on gram-negative bacteria (*E. coli* O157:H7) than gram-positive bacteria (*L. monocytogenes*). Unlike gram-positive bacteria, gram-negative bacteria have an outer membrane composed of phospholipids, proteins, and lipopolysaccharides




Fig. 2. Bactericidal effect of nisin by concentration (A) and the Nisin, APP, and APP+Nisin treatment (B) on the bacterial solution. Error bars represent SD. ^{A-C} Different letters indicate a significant difference ($p < 0.05$) among the treatment. Nisin, 100 ppm nisin-only treatment; APP, cold atmospheric pressure plasma-only treatment; APP+Nisin, combination of APP and nisin treatment.

(Abee et al., 1995). These outer membranes exhibited impermeability to nisin. As nisin is greater than 3 kDa, it is unable to react the cytoplasmic membrane (Abee et al., 1995; Stiles and Hastings, 1991). Therefore, to improve the bactericidal effect of nisin on *E. coli* O157:H7, the APP+Nisin was evaluated. Based on the bactericidal effect of nisin on the bacterial solution, a concentration of 100 ppm was selected for the APP+Nisin.

Synergistic effect of atmospheric pressure plasma (APP) and nisin combination against *Escherichia coli* O157:H7 and *Listeria monocytogenes*

The synergistic effect of APP+Nisin against *E. coli* O157:H7 and *L. monocytogenes* was evaluated (Fig. 2B). The Nisin exhibited the lowest reduction of *E. coli* O157:H7 among all treatments ($p < 0.05$), and there was no difference compared to the control. No viable cells were detected in *E. coli* O157:H7 cells treated with APP. Furthermore, no viable *E. coli* O157:H7 cells in the APP+Nisin were detected.

Similar to the result for *E. coli* O157:H7, Nisin exhibited the lowest bactericidal effect among all the treatment groups against *L. monocytogenes* ($p < 0.05$). No viable *L. monocytogenes* cells were detected after treatment with either APP or APP+Nisin.

In this study, APP treatment (APP and APP+Nisin) exhibited significantly higher bactericidal effects than Nisin on *E. coli* O157:H7 and *L. monocytogenes* ($p < 0.05$). However, the bactericidal effect of APP was too strong to evaluate the synergistic effects of APP and nisin. Unlike the bactericidal effect of Nisin on *E. coli* O157:H7 and *L. monocytogenes*, APP exhibited the

same bactericidal effect on both pathogens. This is because during APP, the outer membrane of *E. coli* O157:H7 is damaged by reactive species capable of destroying the DNA, proteins, lipids, and cell membranes of bacterial cells (Kim et al., 2013), resulting in increased permeability of the lipopolysaccharide layer, which would enhance the activity of nisin against bacterial cells. Similarly, according to Yoo et al. (2021), APP of *E. coli* O157:H7 for 4 min reduced viable cells by 22.31% compared to the control group. Furthermore, *E. coli* O157:H7 and *L. monocytogenes* exhibited a 100% reduction in cell viability compared with the control group when treated with APP for 90 s and 10 min, respectively (Yong et al., 2015).

Bactericidal effect of atmospheric pressure plasma (APP) and nisin combination treatment against inoculated beef jerky and sliced ham

The effect of the APP+Nisin on beef jerky and sliced ham was assessed by inoculation with *E. coli* O157:H7 (Fig. 3). The Nisin, APP, and APP+Nisin decreased the number of viable cells of *E. coli* O157:H7 in beef jerky by 0.19, 0.38, and 0.80 Log CFU/mL with a reduction rate of 3.37%, 6.75%, and 14.21%, respectively, compared to the control. Nisin and APP treatment in beef jerky did not exhibit significant differences when compared with the control. In contrast, the APP+Nisin exhibited the highest bactericidal effect among the treatments ($p < 0.05$).

Similarly, Nisin did not decrease the number of *E. coli* O157:H7 in sliced ham. However, APP and the APP+Nisin induced a decrease in *E. coli* O157:H7 in sliced ham at 1.34 and 1.96 Log CFU/g with a reduction rate of 21.72% and 31.77%, respectively, compared to the control. Both the APP and the APP+Nisin exhibited significantly higher bactericidal effects than the control and Nisin; in particular, the APP+Nisin exhibited a higher bactericidal effect than the APP ($p < 0.05$).

The APP+Nisin exhibited synergistic bactericidal effects in both beef jerky and sliced ham inoculated with *E. coli* O157:H7. However, the bactericidal effect of APP and the APP+Nisin on *E. coli* O157:H7 in meat products was lower than that in bacterial solutions. This is because changes in the physicochemical properties of plasma occurred during APP treatment for organic matters, such as beef jerky and sliced ham. Specifically, the various reactive species generated via APP could react with organic matter, which would reduce the concentration of reactive species capable of responding to bacterial cells. In addition, during the reaction with organic matters, pH may increase and oxidation-reduction potential may decrease, which may also affect the bactericidal effect of APP (Baek et al., 2020a; Xiang et al., 2019). For this reason, the degree of




Fig. 3. Bactericidal effect of Nisin, APP, and APP+Nisin treatment on beef jerky (A) and sliced ham (B) inoculated with *Escherichia coli* O157:H7. Error bars represent SD. ^{A-C} Different letters indicate a significant difference ($p < 0.05$) among the treatment. Nisin, 100 ppm nisin-only treatment; APP, cold atmospheric pressure plasma-only treatment; APP+Nisin, combination of APP and nisin treatment.

damage that APP can induce on the outer membrane of *E. coli* O157:H7 may be reduced, which may be the cause of the lower bactericidal effects of meat products compared to bacterial cells.

In addition, the degree of reduction rate of the APP+Nisin against *E. coli* O157:H7 on beef jerky and sliced ham were 14.21% and 31.77%, respectively. Differences in appearance, such as surface shape and thickness of the samples, can affect the degree of destruction of the bacterial cell membrane by APP. An irregular surface of a sample can create a physical barrier to protect the bacterial strain, which require more energy and time for agents to exert their antibacterial effect (Fernández et al., 2013). In other words, compared to sliced ham, beef jerky with its irregular surface, makes it more difficult for the APP+Nisin to exert its bactericidal effect, compared to that in solutions on *E. coli* O157:H7. This could be responsible for the lower bactericidal effect of APP on beef jerky.

Conclusion

The APP+Nisin exhibited a synergistic bactericidal effect against *E. coli* O157:H7 (gram-negative bacteria). In addition, their bactericidal effect was demonstrated in beef jerky and sliced ham inoculated with *E. coli* O157:H7. This shows that the APP+Nisin has the potential to efficiently control microorganisms on the surfaces of meat and meat products. Further research is needed to establish the optimal conditions for the APP+Nisin applicable for various meat products.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This research was supported by the Basic Science Research Program through the National Research Foundation of Korea (NRF), funded by the Ministry of Education (2021R1I1A1A01044665).

Author Contributions

Conceptualization: Baek KH, Jo C. Data curation: Lee HJ, Heo Y, Kim HJ. Formal analysis: Lee HJ, Kim HJ. Methodology: Lee HJ, Heo Y, Kim HJ, Baek KH. Software: Lee HJ, Kim HJ, Yim DG, Jo C. Validation: Lee HJ, Kim HJ, Jo C. Investigation: Lee HJ, Heo Y, Kim HJ, Sethukali AK, Park D. Writing - original draft: Lee HJ, Kim HJ. Writing – review & editing: Lee HJ, Heo Y, Kim HJ, Baek KH, Yim DG, Sethukali AK, Park D, Jo C.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

Abee T, Krockel L, Hill C. 1995. Bacteriocins: Modes of action and potentials in food preservation and control of food poisoning. *Int J Food Microbiol* 28:169-185.

- Baek KH, Heo YS, Park JY, Kang T, Lee YE, Lim J, Kim SB, Jo C. 2020a. Inactivation of *Salmonella* Typhimurium by non-thermal plasma bubbles: Exploring the key reactive species and the influence of organic matter. *Foods* 9:1689.
- Baek KH, Yong HI, Yoo JH, Kim JW, Byeon YS, Lim J, Yoon SY, Ryu S, Jo C. 2020b. Antimicrobial effects and mechanism of plasma activated fine droplets produced from arc discharge plasma on planktonic *Listeria monocytogenes* and *Escherichia coli* O157:H7. *J Phys D Appl Phys* 53:124002.
- Costello KM, Smet C, Gutierrez-Merino J, Bussemaker M, Van Impe JF, Velliou EG. 2021. The impact of food model system structure on the inactivation of *Listeria innocua* by cold atmospheric plasma and nisin combined treatments. *Int J Food Microbiol* 337:108948.
- Demirok E, Veluz G, Stuyvenberg WV, Castañeda MP, Byrd A, Alvarado CZ. 2013. Quality and safety of broiler meat in various chilling systems. *Poult Sci* 92:1117-1126.
- Fernández A, Noriega E, Thompson A. 2013. Inactivation of *Salmonella enterica* serovar Typhimurium on fresh produce by cold atmospheric gas plasma technology. *Food Microbiol* 33:24-29.
- Gallo LI, Pilosof AMR, Jagus RJ. 2007. Effect of the sequence of nisin and pulsed electric fields treatments and mechanisms involved in the inactivation of *Listeria innocua* in whey. *J Food Eng* 79:188-193.
- Gourama H. 2020. Foodborne pathogens. In *Food safety engineering*. Demirci A, Feng H, Krishnamurthy K (ed). Springer, Cham, Switzerland.
- Heo YS, Yim DG, Baek KH, Kang T, Lee YE, Kim J, Choe W, Jo C. 2021. Effect of inkjet-printed flexible dielectric barrier discharge plasma on reduction of pathogen and quality changes on sliced cheese. *LWT-Food Sci Technol* 143:111128.
- Heredia N, García S. 2018. Animals as sources of food-borne pathogens: A review. *Anim Nutr* 4:250-255.
- Jadhav HB, Annapure US, Deshmukh RR. 2021. Non-thermal technologies for food processing. *Front Nutr* 8:657090.
- Kang T, Yim D, Kim SS, Baek KH, Kim HJ, Jo C. 2022. Effect of plasma-activated acetic acid on inactivation of *Salmonella* Typhimurium and quality traits on chicken meats. *Poult Sci* 101:101793.
- Kim HJ, Yong HI, Lee BW, Park S, Baek KH, Kim TH, Jo C. 2020. Plasma-polymerized phlorotannins and their enhanced biological activities. *J Agric Food Chem* 68:2357-2365.
- Kim HJ, Yong HI, Park S, Choe W, Jo C. 2013. Effects of dielectric barrier discharge plasma on pathogen inactivation and the physicochemical and sensory characteristics of pork loin. *Curr Appl Phys* 13:1420-1425.
- Lee HJ, Jung H, Choe W, Ham JS, Lee JH, Jo C. 2011. Inactivation of *Listeria monocytogenes* on agar and processed meat surfaces by atmospheric pressure plasma jets. *Food Microbiol* 28:1468-1471.
- Li H, Xu Z, Zhao F, Wang Y, Liao X. 2016. Synergetic effects of high-pressure carbon dioxide and nisin on the inactivation of *Escherichia coli* and *Staphylococcus aureus*. *Innov Food Sci Emerg Technol* 33:180-186.
- Liang ZR, Hsiao HI, Jhang DJ. 2020. Synergistic antibacterial effect of nisin, ethylenediaminetetraacetic acid, and sulfite on native microflora of fresh white shrimp during ice storage. *J Food Saf* 40:e12794.
- Liao H, Jiang L, Cheng Y, Liao X, Zhang R. 2018. Application of nisin-assisted thermosonication processing for preservation and quality retention of fresh apple juice. *Ultrason Sonochem* 42:244-249.
- López-Pedemonte TJ, Roig-Sagués AX, Trujillo AJ, Capellas M, Guamis B. 2003. Inactivation of spores of *Bacillus cereus* in cheese by high hydrostatic pressure with the addition of nisin or lysozyme. *J Dairy Sci* 86:3075-3081.
- Mani-López E, García HS, López-Malo A. 2012. Organic acids as antimicrobials to control *Salmonella* in meat and poultry products. *Food Res Int* 45:713-721.
- Min JS, Lee SO, Jang A, Jo C. 2007. Irradiation and organic acid treatment for microbial control and the production of

- biogenic amines in beef and pork. *Food Chem* 104:791-799.
- Mok JH, Pyatkovskyy T, Yousef A, Sastry SK. 2020. Synergistic effects of shear stress, moderate electric field, and nisin for the inactivation of *Escherichia coli* K12 and *Listeria innocua* in clear apple juice. *Food Control* 113:107209.
- Novickij V, Zinkevičienė A, Stanevičienė R, Gruškieienė R, Servienė E, Vepškaitė-Monstavičė I, Krivorotova T, Lastauskienė E, Sereikaitė J, Girkontaitė I, Novickij J. 2018. Inactivation of *Escherichia coli* using nanosecond electric fields and nisin nanoparticles: A kinetics study. *Front Microbiol* 9:3006.
- Osaie R, Essilfie G, Alolga RN, Akaba S, Song X, Owusu-Ansah P, Zhou C. 2020. Application of non-thermal pretreatment techniques on agricultural products prior to drying: A review. *J Sci Food Agric* 100:2585-2599.
- Pokhrel PR, Toniazzo T, Boulet C, Oner ME, Sablani SS, Tang J, Barbosa-Cánovas GV. 2019. Inactivation of *Listeria innocua* and *Escherichia coli* in carrot juice by combining high pressure processing, nisin, and mild thermal treatments. *Innov Food Sci Emerg Technol* 54:93-102.
- Stiles ME, Hastings JW. 1991. Bacteriocin production by lactic acid bacteria: Potential for use in meat preservation. *Trends Food Sci Technol* 2:247-251.
- Tong Z, Ni L, Ling J. 2014. Antibacterial peptide nisin: A potential role in the inhibition of oral pathogenic bacteria. *Peptides* 60:32-40.
- Ukuku DO, Niemira BA, Ukanalis J. 2019. Nisin-based antimicrobial combination with cold plasma treatment inactivate *Listeria monocytogenes* on Granny Smith apples. *LWT-Food Sci Technol* 104:120-127.
- Wang Z, Bi X, Xiang R, Chen L, Feng X, Zhou M, Che Z. 2018. Inactivation of *Escherichia coli* by ultrasound combined with nisin. *J Food Prot* 81:993-1000.
- Xiang Q, Kang C, Zhao D, Niu L, Liu X, Bai Y. 2019. Influence of organic matters on the inactivation efficacy of plasma-activated water against *E. coli* O157:H7 and *S. aureus*. *Food Control* 99:28-33.
- Yong HI, Kim HJ, Park S, Alahakoon AU, Kim K, Choe W, Jo C. 2015. Evaluation of pathogen inactivation on sliced cheese induced by encapsulated atmospheric pressure dielectric barrier discharge plasma. *Food Microbiol* 46:46-50.
- Yoo JH, Baek KH, Heo YS, Yong HI, Jo C. 2021. Synergistic bactericidal effect of clove oil and encapsulated atmospheric pressure plasma against *Escherichia coli* O157:H7 and *Staphylococcus aureus* and its mechanism of action. *Food Microbiol* 93:103611.
- Zhao X, Chen L, Wu J, He Y, Yang H. 2020. Elucidating antimicrobial mechanism of nisin and grape seed extract against *Listeria monocytogenes* in broth and on shrimp through NMR-based metabolomics approach. *Int J Food Microbiol* 319:108494.
- Zhao Y, Shao L, Jia L, Meng Z, Liu Y, Wang Y, Zou B, Dai R, Li X, Jia F. 2022. Subcellular inactivation mechanisms of *Pseudomonas aeruginosa* treated by cold atmospheric plasma and application on chicken breasts. *Food Res Int* 160:111720.