



Positive rates for *Enterobius vermicularis* eggs among preschool children in Yeosu-si, Jeollanam-do, Korea (2017-2021)



Myoung-Ro Lee¹ , Hee-Eun Shin¹, Seon-Ok Back¹, Young-Ju Lee¹, Jung-Won Ju¹ , Chun Soon Park², Hee-Il Lee^{1,*}

¹Division of Vectors and Parasitic Diseases, Korea Diseases Control and Prevention Agency, Osong 28159, Korea; ²Yeosu-si Public Health Center, Yeosu 59674, Korea

Abstract

Received: 13 September 2022
Accepted: 22 December 2022

*Correspondence
(isak@korea.kr)

Citation

Lee MR, Shin HE, Back SO, Lee YJ, Ju JW, Park CS, Lee HI. Positive rates for *Enterobius vermicularis* eggs among preschool children in Yeosu-si, Jeollanam-do, Korea (2017-2021). Parasit Host Dis 2023;61(1):84-88.

This study aimed to evaluate the positive rates for *Enterobius vermicularis* eggs among preschool children in Yeosu-si, Jeollanam-do, the Republic of Korea (Korea) over a period of 5 years (2017-2021). Perianal swab samples, obtained using cellophane tape, from 10,392 preschool children in 26 districts were examined microscopically for *E. vermicularis* eggs. The test results were notified through the local health center, and the families of children who tested positive were advised to provide them anthelmintics treatment. The annual positive rates were 5.0%, 5.2%, 4.4%, 2.2%, and 1.0% in 2017, 2018, 2019, 2020, and 2021, respectively. The overall positive rate was higher in boys than in girls ($P < 0.05$), and children aged 5-7 years were at a higher risk of being infected than those aged 0-4 years ($P < 0.05$). Although the rates of infection by *E. vermicularis* in the survey area, Yeosu-si, were still in the 1% range, the results of this study suggest that they can be significantly reduced through continuous intervention centered around the test-treatment strategy.

Keywords: *Enterobius vermicularis*, cellophane tape perianal swap, preschool children, Yeosu-si

Enterobiasis, an infection caused by the human nematode parasite *Enterobius vermicularis* (pinworm), is commonly observed in children from developed and developing countries. It typically spreads through direct transmission from an infected individual to an uninfected one, but it can also be contracted through actions, such as sucking on toys, pencil biting, and playing in areas contaminated with pinworm eggs [1]. Moreover, it is commonly observed in overcrowded areas, such as preschools [2], and its estimated global prevalence in children is 5.1-22.4% [3-6]. In 2000, Park et al. [7] reported that the positive rate of *E. vermicularis* infection among children in the western and southern offshore islands of the Republic of Korea (Korea) was 18.5%, whereas other studies reported rates ranging from 3.5% to 10.0% in other parts of Korea, such as Paju-si (3.5%), Inje-gun (4.6%), Chuncheon-si (5.7%) [8], Gimhae-si (10.5%) [9], and Muan-gun (4.0%) [10]. Moreover, the prevalence rates of infection in Seoul and other large cities have been reported to range between 0.6% and 3.9% during 2008-2019 [11]. However, to the best of our knowledge, no consecutive surveys of the rates of pinworm infections among preschool children in the same local region have been conducted to date. Therefore, this study aimed to determine the incidence of *E. vermicularis* infection among preschool children in Yeosu-si.

The need for ethical approval was waived as this study was conducted to evaluate public welfare through a fact-finding survey (Infectious Disease Control and Prevention Act, Ar-

© 2023 The Korean Society for Parasitology and Tropical Medicine

This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License (<https://creativecommons.org/licenses/by-nc/4.0>) which permits unrestricted non-commercial use, distribution, and reproduction in any medium, provided the original work is properly cited.

Author contributions

Data curation: Ju JW
 Investigation: Shin HE, Back SO, Lee YJ,
 Park CS
 Supervision: Lee HI
 Writing – original draft: Lee MR

Conflict of interest

The authors declare no conflict of interest related to this study.

ORCID


Myoung-Ro Lee
 (https://orcid.org/0000-0003-3369-0063)
 Jung-Won Ju
 (https://orcid.org/0000-0001-5248-6995)
 Hee-Il Lee
 (https://orcid.org/0000-0002-1636-0639)

ticle 17). In total, 10,392 children from 26 districts (including 20 from dong, 1 from eup, and 5 from myeon) in Yeosu-si were examined for the presence of *E. vermicularis* eggs using the cello tape perianal swab method (Fig. 1). Parents of these children were instructed to press the sticky side of a cellophane tape on the perianal area of the child immediately after they woke up, and all collected samples were collected by the preschool teachers and transported to the Division of Vectors and Parasitic Diseases, Korea Disease Control and Prevention Agency. Subsequently, the samples were assessed by qualified technicians using light microscopy, with the presence of eggs being confirmed at low magnification ($\times 100$) and identification of the *E. vermicularis* eggs being carried out at high magnification ($\times 400$). Notably, the pinworm egg resembles a persimmon seed in shape, with length and width ranging from 50 to 60 μm and 20 to 30 μm , respectively.

Data analysis included comparison of categorical variables using a chi-square test, and the incidence trends of *E. vermicularis* infection were examined in terms of age and sex. All statistical analyses were performed using SPSS, v18.0 (IBM, Chicago, IL, USA), and the level of statistical significance was set at $P < 0.05$ with a 95% confidence interval.

Overall, > 2,000 samples were examined every year (except 2019), and the results revealed that the rates of *E. vermicularis* infection in Yeosu-si, Jeollanam-do, gradually decreased between 2017 and 2021 (Fig. 2; 5.0% (116/2,307 children) in 2017; 5.2% (120/2,297) in 2018; 4.4% (15/329) in 2019; 2.2% (58/2,664) in 2020; and 1.0% (27/2,821) in 2021). Notably, the rate of *E. vermicularis* infection was statistically significantly declined by 2021 ($P < 0.05$).

This study was conducted in 26 of 27 districts in Yeosu-si (excluding Samil-dong). In 2017, the rate of *E. vermicularis* infection was highest in Daegyo-dong (10.6%), followed by Guk-dong (9.4%), Dolsan-eup (9.0%), and the other regions ($< 9\%$). In 2021, the rates of infections in Hwayang-myeon (8.3%) and Samsan-myeon (25.0%) were high, although the






Fig. 2. Egg-positive rate of *E. vermicularis* in Yeosu-si, Jellanam-do, Korea, from 2017 to 2021.

Table 1. Egg positive rates of *Enterobius vermicularis* by the survey year and region

Mark*	Regions	No. positive/No. examined (%)				
		2017	2018	2019	2020	2021
1	Yulchon-myeon	0/10 (0.0)	1/28 (3.6)	0/0 (0.0)	4/42 (9.5)	2/40 (5.0)
2	Myodo-dong	0/0 (0.0)	0/0 (0.0)	0/0 (0.0)	0/15 (0.0)	0/0 (0.0)
3	Sola-myeon	13/215 (6.0)	0/12 (0.0)	0/0 (0.0)	4/268 (1.5)	2/198 (1.0)
4	Jusam-dong	2/105 (1.9)	2/107 (1.9)	0/0 (0.0)	1/50 (2.0)	4/65 (6.2)
5	Yecheon-dong	12/345 (3.5)	15/444 (3.4)	5/40 (12.5)	2/63 (3.2)	1/277 (0.4)
6	Dundeok-dong	0/16 (0.0)	2/35 (5.7)	0/0 (0.0)	1/48 (2.1)	0/66 (0.0)
7	Mipyeong-dong	11/218 (5.0)	12/220 (5.5)	0/0 (0.0)	8/239 (3.3)	1/70 (1.4)
8	Mandeok-dong	0/11 (0.0)	15/110 (13.6)	0/0 (0.0)	2/179 (1.1)	1/85 (1.2)
9	Ssangbong-dong	11/257 (4.3)	16/277 (5.8)	8/142 (5.6)	8/184 (4.3)	3/430 (0.7)
10	Sijeon-dong	21/389 (5.4)	15/211 (7.1)	0/0 (0.0)	1/248 (0.4)	0/312 (0.0)
11	Moonsu-dong	13/179 (7.3)	14/405 (3.5)	2/139 (1.4)	7/501 (1.4)	8/512 (1.6)
12	Yeoseo-dong	9/178 (5.1)	5/93 (5.4)	0/0 (0.0)	3/381 (0.8)	1/386 (0.3)
13	Gwangnim-dong	0/16 (0.0)	0/0 (0.0)	0/0 (0.0)	0/0 (0.0)	0/0 (0.0)
14	Hwayang-myeon	1/32 (3.1)	0/0 (0.0)	0/0 (0.0)	1/27 (3.7)	1/12 (8.3)
15	Wolho-dong	0/0 (0.0)	0/32 (0.0)	0/0 (0.0)	0/26 (0.0)	2/95 (2.1)
16	Guk-dong	3/32 (9.4)	2/91 (2.2)	0/0 (0.0)	3/74 (4.1)	0/7 (0.0)
17	Daegyo-dong	5/47 (10.6)	0/7 (0.0)	0/0 (0.0)	1/45 (2.2)	0/38 (0.0)
18	Seogang-dong	2/71 (2.8)	2/57 (3.5)	0/0 (0.0)	1/23 (4.3)	0/15 (0.0)
19	Chungmu-dong	0/0 (0.0)	0/0 (0.0)	0/0 (0.0)	0/56 (0.0)	0/59 (0.0)
20	Jungang-dong	0/0 (0.0)	8/32 (25.0)	0/0 (0.0)	2/32 (6.3)	0/32 (0.0)
21	Dongmun-dong	1/41 (2.4)	0/6 (0.0)	0/0 (0.0)	7/55 (12.7)	0/6 (0.0)
22	Hallyeo-dong	0/0 (0.0)	0/0 (0.0)	0/0 (0.0)	1/20 (0.0)	0/0 (0.0)
23	Hwajeong-myeon	0/0 (0.0)	0/8 (0.0)	0/0 (0.0)	0/0 (0.0)	0/1 (0.0)
24	Nam-myeon	0/0 (0.0)	1/11 (9.1)	0/0 (0.0)	0/7 (0.0)	0/22 (0.0)
25	Samsan-myeon	0/12 (0.0)	0/0 (0.0)	0/0 (0.0)	0/0 (0.0)	1/4 (25.0)
26	Dolsan-eup	12/133 (9.0)	10/111 (9.0)	0/8 (0.0)	1/81 (1.2)	0/89 (0.0)
	Total	116/2,307 (5.0)	120/2297 (5.2)	15/329 (4.6)	58/2,664 (2.2)	27/2,821 (1.0)

*The districts surveyed in this study.

Table 2. Comparison of the egg positive rates of *Enterobius vermicularis* by the gender and age-group in 2017 and 2021

Category	2017			2021		
	No. examined	No. positive (%)	<i>P</i> -value	No. examined	No. positive (%)	<i>P</i> -value
Gender						
Male	1,068	67 (6.3)	< 0.05	1,414	18 (1.3)	0.046
Female	1,037	34 (3.3)		1,374	9 (0.7)	
Age group (yr)						
0-4	408	14 (3.4)	0.07	1,922	11 (0.6)	< 0.05
5-7	1,608	86 (5.3)		898	16 (1.8)	

number of examinees (or subjects) was small (Table 1). In districts with high infection rates, including Daegyo-dong, Guk-dong, Dolsan-eup, Moonsu-dong, and Sola-myeon, the rates of *E. vermicularis* infections decreased from 6.0 to 10.6% in 2017 and from 6.0 to 1.6% in 2021. This reduction in infection rates could be attributed to the continuous management and treatment of infected children and their family members, which is considered an effective strategy for controlling *E. vermicularis* infection among preschool children. Examination of the risk of *E. vermicularis* infection by sex revealed that males exhibited 2 times higher risk (6.3%) than females (3.3%) in 2017. Similar results were observed in 2021, despite the decrease in overall rates of infection ($P < 0.05$; Table 2).

The study population was classified into age groups of 0-4 and 5-7 years to account for differences in preschool programs (i.e., children aged < 4 years: nap time and fewer educational activities; children aged > 5 years: no nap time and more opportunities to engage in outdoor activities with children from other classes). The children in the 5-7 year age group exhibited significantly higher infection rates than those in the 0-4 year age group ($P < 0.05$). Moreover, in 2017, children in the 5-7 year age group exhibited a 1.5 times higher probability of being infected with *E. vermicularis* than those in the 0-4 year age group, and this probability increased to 3 times in 2021 ($P < 0.05$; Table 2). This finding is consistent with previous studies, which have also observed a higher risk of pinworm infection in children in the 5-7 year age group than in younger children, possibly due to more frequent exposure to contaminated environments [12-15].

Hong et al. [10] reported that the *E. vermicularis* infection rate in Jeollanam-do was 1.5% in 2019. However, the infection rate in Yeosu-si at the beginning of this survey was 3 times higher (4.6%) than that observed in Jeollanam-do in 2019, indicating that it can be considered a high-risk area for pinworm infections requiring the adoption of continuous management strategies.

In conclusion, *E. vermicularis* infections are still widely prevalent among preschool children in Yeosu-si, and continuous examination and treatment of infected children and their families is required to decrease incidence rates.

Acknowledgments

We thank the person in charge of the Yeosu Health Center for participating in sample collection for this study. This work was supported by a grant (6331-311-210-13, 2021) from the Korea Disease Control and Prevention Agency (KDCA), Republic of Korea.

References

1. Wendt S, Trawinski H, Schubert S, Rodloff AC, Mössner J, Lübbert C. The diagnosis and treatment of pinworm infection. *Dtsch Arztebl Int* 2019;116(13):213-219. <https://doi.org/10.3238/arztebl.2019.0213>
2. Song HJ, Cho CH, Kim JS, Choi MH, Hong ST. Prevalence and risk factors for enterobiasis among preschool children in a metropolitan city in Korea. *Parasitol Res* 2003;91(1):46-50. <https://doi.org/10.1007/s00436-003-0836-3>
3. Ryue HS, Jung JW, Pai KS. An epidemiological study on refractory enterobiasis. *Korean J Pediatr* 2004;47(2):177-182 (in Korean).
4. Fan CK, Chuang TW, Huang YC, Yin AW, Chou CM, et al. *Enterobius vermicularis* infection: prevalence and risk factors among preschool children in kindergarten in the capital area, Republic of the Marshall Islands. *BMC Infectious Diseases* 2019 19(1);1:536. <https://doi.org/10.1186/s12879-019-4159-0>
5. Wang S, Yao Z, Hou Y, Wang D, Zhang H, et al. Prevalence of *Enterobius vermicularis* among preschool children in 2003 and 2013 in Xinxiang city, Henan province, Central China. *Parasite* 2016;23:30. <https://doi.org/10.1051/parasite/2016030>
6. Friesen J, Bergmann C, Neuber R, Fuhrmann J, Wenzel T, et al. Detection of *Enterobius vermicularis* in greater Berlin, 2007-2017: seasonality and increased frequency of detection. *Eur J Clin Microbiol Infect Dis* 2019;38(4):719-723. <https://doi.org/10.1007/s10096-019-03495-1>
7. Park JH, Han ET, Kim WH, Shin EH, Guk SM, et al. A survey of *Enterobius vermicularis* infection among children on western and southern coastal islands of the Republic of Korea. *Korean J Parasitol* 2005;43(4):129-134. <https://doi.org/10.3347/kjp.2005.43.4.129>
8. Hong SH, Lee SE, Jeong YI, Lee WJ, Cho SH. Comparison of egg positive rates of *Enterobius vermicularis* among preschool children in three Korean localities. *Korean J Parasitol* 2011; 49(4):441-443. <https://doi.org/10.3347/kjp.2011.49.4.441>
9. Lee SE, Lee JH, Ju JW, Lee WJ, Cho SH. Prevalence of *Enterobius vermicularis* among preschool children in Gimhae-si, Gyeongsangnam-do, Korea. *Korean J Parasitol* 2011;49(2): 183-185. <https://doi.org/10.3347/kjp.2011.49.2.183>
10. Hong SH, Jeong YI, Lee JH, Cho SH, Lee WJ, Lee SE. Prevalence of *Enterobius vermicularis* among preschool children in Muan-gun, Jeollanam-do, Korea. *Korean J Parasitol* 2012;50(3): 259-262. <https://doi.org/10.3347/kjp.2012.50.3.259>
11. Shin HJ, Jung BK, Ryoo SW, Hong SJ, Chang TH, et al. *Enterobius vermicularis* infection among preschool children: a 12-year (2008-2019) survey in large cities and provinces of the Republic of Korea. *Korean J Parasitol* 2021;59(4):421-426. <https://doi.org/10.3347/kjp.2021.59.4.421>
12. Rivero MR, De Angelo C, Feliziani C, Liang S, Tiranti K, et al. Enterobiasis and its risk factors in urban, rural and indigenous children of subtropical Argentina. *Parasitol* 2022;149(3):396-406. <https://doi.org/10.1017/S0031182021001955>
13. Chai JY, Yang SK, Kim JW, Choi SL, Song GY, et al. High prevalence of *Enterobius vermicularis* infection among schoolchildren in three townships around Yangon, Myanmar. *Korean J Parasitol* 2015;53(6):771-775. <https://doi.org/10.3347/kjp.2015.53.6.771>
14. Lee KJ, Lee IY, Im K. *Enterobius vermicularis* egg positive rate in a primary school in Chungchongnam-do (Province) in Korea. *Korean J Parasitol* 2000;38(3):177-178. <https://doi.org/10.3347/kjp.2000.38.3.177>
15. Song HJ, Cho CH, Kim JS, Choi MH, Hong ST. Prevalence and risk factors for enterobiasis among preschool children in a metropolitan city in Korea. *Parasitol Res* 2003;91(1):46-50. <https://doi.org/10.1007/s00436-003-0836-3>