



Original Article

# Dental plaque removal efficiency of a smart toothbrush based on augmented reality in children

Min-Ji Park<sup>1†</sup> · Suk-Bin Jang<sup>2†</sup> · Jae-Young Lee<sup>1†</sup>

<sup>1</sup>Department of Dental Hygiene, Dankook University

<sup>2</sup>Division of Infectious Diseases, Departments of Internal Medicine, Dankook University Hospital, Dankook University College of Medicine

**Corresponding Author: Jae-Young Lee**, Department of Dental Hygiene, Dankook University, 119 Dandae-ro, Dongnam-gu, Cheonan-si, Chungcheongnam-do, 31116, Korea. Tel: +82-41-550-1496, Fax: +82-504-090-7206, E-mail: dentaljy@dankook.ac.kr

## ABSTRACT

**Objectives:** Based on the strengths of augmented reality (AR), this study aimed to determine the dental plaque removal efficiency of a guided brushing program in children. **Methods:** This randomized, controlled, double-blind crossover clinical trial evaluated the plaque removal rate after regular brushing versus AR-based guided toothbrushing in 20 children aged 5-12 years. **Results:** Overall, the dental plaque removal efficiency of AR-based brushing was superior to that of regular brushing ( $p<0.05$ ). When classified in detail, no significant difference was noted in the plaque removal rate between the two brushing methods in the anterior region ( $p=0.056$ ), whereas a significant difference in the plaque removal rate was observed in the posterior region ( $p<0.05$ ). **Conclusions:** Based on these results, the efficacy of dental plaque removal for brushing using an AR-based smartphone application was confirmed; thus, this can be used for oral health education incorporating ICT technology in the future.

**Key Words:** Augmented Reality, Dental Plaque, Toothbrushing

## Introduction

Periodontal disease and dental caries were the 5th most expensive medical expenses in the first half of 2021 in Korea. Since the statistical cost index includes systemic diseases, this ranking is quite high[1]. This means that treatment-oriented dental services are being prioritized over preventive care. In addition, the cost of dental treatment is increasing due to this.

According to the 2018 Korean National Children's Oral Health Survey, More than half of 12-year-olds have experienced dental caries, and the average number of decayed tooth by 12-year-olds was 1.84, higher than the average of 1.2 in OECD member countries. These statistical results suggest that more than half of school-age children suffer from dental diseases, such as dental caries; thus, awareness of oral health requires improvement [2].

Dental plaque can lead to serious oral disease [3-6]; to prevent disease, various oral care tools are used to manage these dental plaques, but it is difficult to manage them completely [7,8]. In addition, although knowledge about health can be improved using existing health education methods [9,10], knowledge alone does not change oral health behaviors [11]. Augmented reality is being used in various fields such as education and medicine that require behavioral change, and its effectiveness is gradually increasing [12,13].

Brushing not only improves aesthetics by cleaning food residues left behind during eating and drinking, it refreshes the oral cavity and prevents dental plaque formation and acid production [14]. Its mechanical action removes already formed and attached dental plaque, and it is the most commonly used oral health care behavior in daily life [15]. Therefore, it is very important to develop correct behavioral habits to encourage lifelong oral health management, ideally from childhood [16,17]. In addition, children should form good habits before their permanent teeth erupt so that they can maintain a healthy oral condition for the rest of their lives [18,19].

Received January 09, 2023

Revised February 04, 2023

Accepted February 12, 2023

Copyright © 2023 by Journal of Korean Society of Dental Hygiene. This is an Open Access article distributed under the terms of the Creative Commons Attribution Non-Commercial License(<http://creativecommons.org/licenses/by-nc/4.0>)

<sup>†</sup>These authors contributed equally to this work.

AR-based education provides a rich learning environment [20] because it includes additional virtual learning information and enables self-directed experiential learning by allowing direct manipulation using the learner's various senses [21-23]. Tooth brushing education requires instruction on the area to be brushed accurately and education on how to brush, and education through video or real objects is effective. Toothbrushing instruction requires on tooth area to be brushed accurately and education on how to brush [24], and in non-face-to-face education, toothbrushing education is conducted through video media, or in face-to-face education, education is conducted through real objects such as tooth models and toothbrush [25,26].

Since AR encourages induces users to act naturally and actively interact with virtual images, in this case in a brushing habit-forming application, users can continuously and voluntarily develop appropriate toothbrushing and oral health care behaviors.

The purpose of this study was to evaluate whether plaque removal efficiency could be improved through toothbrushing using an AR guide program designed to help learning to brush teeth properly by utilizing the advantages of AR.

## Methods

### 1. Study subjects

The feasibility study was designed using two period cross-over design. This study included 20 subjects residing in C city for people aged by 5-12, who agreed them and their parents to participate after understanding the purpose and contents of this study. This study was approved by the Institutional Review Board (IRB approval No.: S-D20220008) and was conducted in accordance with the ethical standards of the Declaration of Helsinki. The number of samples was calculated using the standard deviation of previous studies, and the total number of samples was calculated as 20 considering the reliability of 90% and dropout rate of less than 10% [14].

### 2. Subject inclusion and exclusion criteria

#### 1) Inclusion & exclusion criteria

**Table 1.** Inclusion & exclusion criteria of this study

Criteria	Contents
Inclusion criteria	1. Participants included age 5-12 years 2. Parents who can use an augmented reality toothbrush using a smartphone
Exclusion criteria	1. Participants with systemic disease 2. Difficulty opening the mouth 3. Fixed prosthesis covering three or more tooth surfaces 4. Current or past illness that could impact the study findings or procedure

#### 2) Criteria for subject discontinuation and withdrawal


The present study could be discontinued or subjects could withdraw if they did not comply with or violated items stated in the written informed consent form, if unpredicted severe complications occurred, or if a researcher or the principal investigator deemed a subject inappropriate for continued study participation.

#### 3) Subject identification

Subjects who met the inclusion and exclusion criteria specified in the study protocol and whose parents provided written informed consent were assigned a unique identifier, formatted as YDF-18##, that was recorded on the corresponding case report form.

#### 4) Overview of study procedures

The present clinical evaluation used a crossover design as described below.



**Fig. 1.** Flow chart of the study procedure (AR: augmented reality)

## 5) The clinical study procedure

### (1) Preparations

This randomized doubleblind clinical study was designed to evaluate for smart brush. During the screening process, the subject's appropriate oral health condition and written consent were obtained, and evaluation was performed using a manual toothbrush, and brushing was standardized by the Fone's method. After that, at the second visit, dental plaque removal effect after using the smart toothbrush was evaluated, and compared with that of first visit.

### 6) Observation items and clinical exams

#### (1) Plaque index test

A disclosing solution (Trace, Disclosing Solution; Young Dental, Earth City, MO, US) was applied to all erupted teeth, and plaque deposit intensities and locations were observed.

After cleaning of all tooth surfaces dried using air syringe, the disclosing solution was applied using cotton pellets moistened with the solution. Once the surfaces of the teeth had been dried, they were evaluated using the Patient Hygiene Performance (PHP) index. The following tooth surfaces were evaluated: buccal surface of the maxillary right first molar, labial surface of the maxillary right central incisor, buccal surface of the maxillary left first molar, lingual surface of the mandibular right first molar, labial surface of the mandibular left central incisor, and lingual surface of the mandibular left first molar.

#### (2) Use of a smart toothbrush

Brush Monster Toothbrush <Fig. 2>, an augmented reality, recognizes the location of the toothbrush through a smart device, and based on this, a guide for the user's brushing area is followed on the smartphone. Through game-based content, it guides you in getting rid of germs in your mouth so that you can brush your teeth without missing parts.



**Fig. 2.** Screen view offered by Brush Monster

Tailored education was provided using a smart toothbrush. Each child used their dominant hand regardless of side and had the option of choosing a toothbrushing technique (scrub, Fones, or Bass). Based on the tailored education, the children could not only develop an effective toothbrushing technique, using a total of 16 steps, from brushing the teeth in the order of occlusal, buccal, and lingual surfaces (starting from the occlusal surface in the mandibular left and ending at the lingual surface in the maxillary right), but they were empowered to do everything on their own from the beginning to the end of the toothbrushing process (from squeezing toothpaste on the toothbrush to washing the cup). Each subject was provided a Brush Monster, and after they completed the training.

The AR-based smart toothbrush used in the present study was intended to instruct the children how to correctly brush step by step.

### 3. Statistical Analysis

The study used two period cross-over design. This study measured tooth-brushing performance and time and oral hygiene using manual toothbrush and after using smart toothbrush. The paired t-test and two groups t-test was used to determine the significant difference in tooth-brushing performance and each test-outcome variable within a group. Statistical significance was set at  $p < 0.05$ . All statistical analyses were conducted using the IBM SPSS program (ver 26.0; IBM Corp., Armonk, NY, USA).

## Results

### 1. Outcome assessment

#### 1) Study subjects

A total of 20 children without any prior toothbrushing education participated in the present study. Of them, 12 were boys and 8 were girls; 9 were 5 years old; 2 each were 7, 8, and 9 years old; and 1 was 10 years old. The mean participant age was 6.35 years.

## 2) Assessment of changes in dental plaque

### (1) Overall assessment

The visit 1 assessment of the effects of manual toothbrushing in the 20 subjects showed that the mean PHP index score decreased from 20.30 points ( $\pm 3.73$ ) before toothbrushing to 13.35 ( $\pm 5.08$ ) after. After a 6-day washout period, the effects of the AR-based vibrating toothbrush used in conjunction with the associated educational program were assessed during visit 2. The mean PHP index score decreased from 22.00 ( $\pm 3.63$ ) before toothbrushing to 6.35 ( $\pm 6.19$ ) after toothbrushing <Table 2>.

**Table 2.** Comparison of changes in dental plaque index between manual toothbrush and AR guided toothbrush

Subject	Manual toothbrush			AR guided toothbrush		
	Pre	Post	Change rate (%)	Pre	Post	Change rate (%)
1	18	8	-55.6	19	6	-68.4
2	20	12	-35.0	23	4	-82.6
3	22	9	-31.8	21	7	-66.7
4	12	8	-16.7	18	7	-61.1
5	27	18	-33.3	20	3	-85.0
6	27	27	-33.3	27	13	-51.9
7	18	10	-22.2	23	10	-56.5
8	19	7	-42.1	24	12	-50.0
9	23	16	-21.7	23	2	-91.3
10	23	6	-39.1	22	9	-59.1
11	26	13	-38.5	18	7	-61.1
12	18	9	-38.9	25	8	-68.0
13	17	17	-23.5	25	0	-100.0
14	18	15	-44.4	14	0	-100.0
15	23	20	-21.7	27	0	-100.0
16	21	15	-42.9	22	0	-100.0
17	19	13	-52.6	19	0	-100.0
18	18	13	-38.9	28	21	-25.0
19	19	15	-52.6	24	18	-25.0
20	18	16	-38.9	18	0	-100.0
Mean $\pm$ SD	20.30 $\pm$ 3.73	13.35 $\pm$ 5.08	-36.2	22.00 $\pm$ 3.63	6.35 $\pm$ 6.19	-72.6
<i>P</i> *			<0.001			<0.001

\*by paired t-test, AR: augmented reality

Comparison of plaque removal efficiency between the Brush Monster and the manual toothbrush showed that the PHP index score decreased by an average of 15.65 and 6.95 points, respectively. The mean plaque removal rate with the Brush Monster was 72.6%, approximately twice as high as that for the manual toothbrush (36.2%,  $p < 0.05$ ).

### (2) Assessment of anterior and posterior teeth

The plaque removal effects of the Brush Monster versus manual brushing were assessed separately for the anterior and posterior teeth. In the anterior teeth, the mean PHP index score was 7.35 points ( $\pm 2.32$ ) before and 3.50 ( $\pm 2.04$ ) after manual brushing, showing a decrease of 3.85 points. In the posterior teeth, the mean was 12.95 ( $\pm 3.30$ ) before and 9.85 ( $\pm 4.46$ ) after manual brushing, showing a decrease of 3.10 points <Table 3>. The plaque removal effects of the AR-based vibrating toothbrush, however, were as follows: In the anterior teeth, the mean PHP index score was 7.70 ( $\pm 2.18$ ) before and 2.30 ( $\pm 2.64$ ) after using the AR-based toothbrush, showing a decrease of 5.40. In the posterior teeth, the mean PHP index score was 14.30 ( $\pm 3.29$ ) before and 4.05 ( $\pm 5.03$ ) after using the AR-based toothbrush, showing a decrease of 10.25 <Table 4>. Thus, the difference in the plaque removal effects in the anterior teeth between the manual and AR-based smart vibrating toothbrushes was 1.55 points ( $\pm 3.41$ ), with the effect being greater for the AR-based toothbrush, although the difference was not statistically significant ( $p = 0.056$ ). In the posterior teeth, the difference was 7.15 points ( $\pm 5.65$ ), with the plaque removal effects being greater in the AR-based smart vibrating toothbrush as well, although this difference was statistically significant ( $p < 0.05$ ).

**Table 3.** Comparison of changes in dental plaque index according to anterior site and posterior site in manual toothbrush

Subject	Manual toothbrush			AR guided toothbrush		
	Pre	Post	Change rate (%)	Pre	Post	Change rate (%)
1	10	3	-70.0	8	5	-37.5
2	7	5	-28.6	13	7	-46.2
3	7	5	-28.6	15	4	-73.3
4	2	2	0.0	10	6	-40.0
5	9	3	-66.7	18	15	-16.7
6	9	9	0.0	18	18	0.0
7	4	0	-100.0	14	10	-28.6
8	8	3	-62.5	11	4	-63.6
9	5	0	-100.0	18	16	-11.1
10	9	2	-77.8	14	4	-71.4
11	10	3	-70.0	16	10	-37.5
12	7	3	-57.1	11	6	-45.5
13	4	4	0.0	13	13	0.0
14	8	5	-37.5	10	10	0.0
15	5	2	-60.0	18	18	0.0
16	9	3	-66.7	12	12	0.0
17	10	4	-60.0	9	9	0.0
18	7	3	-57.1	11	10	-9.1
19	10	6	-40.0	9	9	0.0
20	7	5	-28.6	11	11	0.0
Mean±SD	7.35±2.32	3.50±2.04	-50.6	12.95±3.30	9.85±4.46	-24.0
<i>p</i> *			<0.001			<0.001

\*by paired t-test, AR: augmented reality

**Table 4.** Comparison of changes in dental plaque index according to anterior site and posterior site in manual toothbrush in augmented guided toothbrush

Subject	Manual toothbrush			AR guided toothbrush		
	Pre	Post	Change rate (%)	Pre	Post	Change rate (%)
1	6	2	-66.7	13	4	-69.2
2	8	0	-100.0	15	4	-73.3
3	4	0	-100.0	17	7	-58.8
4	3	1	-66.7	15	6	-60.0
5	10	3	-70.0	10	0	-100.0
6	9	8	-11.1	18	5	-72.2
7	6	4	-33.3	17	6	-64.7
8	10	7	-30.0	14	5	-64.3
9	10	1	-90.0	13	1	-92.3
10	10	6	-40.0	12	3	-75.0
11	6	4	-33.3	12	3	-75.0
12	10	5	-50.0	15	3	-80.0
13	9	0	-100.0	16	0	-100.0
14	5	0	-100.0	9	0	-100.0
15	7	0	-100.0	20	0	-100.0
16	9	0	-100.0	13	0	-100.0
17	10	0	-100.0	9	0	-100.0
18	8	4	-50.0	20	17	-15.0
19	7	1	-85.7	17	17	0.0
20	7	0	-100.0	11	0	-100.0
Mean±SD	7.70±2.18	2.30±2.64	-71.3	14.30±3.29	4.05±5.03	-75.0
<i>p</i> *			<0.001			<0.001

\*by paired t-test, AR: augmented reality



## Discussion

Oral health behavior develops in early childhood; accordingly, lifelong oral health habits are influenced, and appropriate education at the relevant time is important [27]. Toothbrushes are also closely associated with tooth loss, periodontal disease, and gum disease. These conditions are closely associated with systemic diseases in adulthood, difficulty forming positive oral health management behaviors, and limited ability to perform complicated oral health management processes [28,29].

With the advanced development of extended reality such as augmented reality and virtual reality, the application of ICT technology to health maintenance programs is increasing. ICT technology can be used for tasks such as identifying and tracking oral health problems, helping with diagnoses and treatment planning, assisting with surgical procedures, and predicting patient outcomes. AR-based devices and software are also evolving to help track and analyze oral hygiene habits and provide personalized oral health recommendations. However, the use of AR in oral health is still in its early stages and further research is needed to fully understand its potential and limitations. Therefore, we conducted this study to evaluate the plaque removal effect using apps currently available in the market and to review the applicability.

Until now, AR technology has been mainly applied to the field of oral surgery, but it is gradually expanding to the field of education. Amantini et al [30] conducted a study to train children between the ages of 6 and 10 on how to brush their teeth using AR technology. And it was announced that digital tools using AR have potential effects to motivate oral hygiene practice and improve children's health awareness. In addition, in the study of a control experiment targeting people with intellectual disabilities, oral hygiene improved significantly in both the AR-based group and the visual data-based education, but the AR-based tooth brushing education was found to be more effective [31].

A comparison of the removal efficiency of the tooth surface dental plaque using the manual toothbrush and the smart toothbrush with the enhanced reality guide showed that the overall tooth surface dental plaque index was significantly reduced. This led to significant results given the fact that guide-based brushing for persons with disabilities with previous behavioral restrictions can better remove dental plaque on the tooth surface and reduce the gingival index [32]. Moreover, it has significance since electric oral management products such as electric toothbrushes and water flossers are useful supplementary oral management tools [33-35]. Regarding changes in child behavior, these AR techniques can be combined with gaming techniques to arouse children's interest, and these repetitive behaviors and education aid in the establishment of correct health habits [36].

The use of an AR-based smart toothbrush in conjunction with an educational program significantly reduced PHP index scores versus the manual toothbrush in the subjects evaluated in the present study. However, the effects of using the AR-based toothbrush on plaque removal in the anterior teeth were not significant compared to those of manual brushing, while the difference was statistically significant in the posterior teeth. Using the application to watch the person on the screen of the mobile phone, the efficiency of removing dental plaque from the tooth surfaces that are difficult to see with the same plane mirror as the oral cavity is lower than that of the anterior teeth. It is important to brush the teeth without falling into the oral cavity to remove dental plaque, and it is very important to manage the gingival margin or brush the teeth in the lingual direction to prevent oral diseases [37]. Therefore, it is necessary to reinforce special care for parts of the mouth where brushing is ineffective by increasing the time allocated to that area in the brushing guide in the future.

This study could consider the augmented reality-based dental plaque removal effect and usability, but there are some factor in interpreting it. First, the AR app guided brushing methods other than the Fones method, and other brushing methods may have affected the removal of dental plaque. Second, in children who normally brush their teeth for less than 3 minutes, the effect of removing dental plaque may be more evident due to the increased brushing time. Finally, it is difficult to generalize to a small sample. Nevertheless, it is meaningful as there are currently few studies that have confirmed oral health indicators using AR-based apps.

This study secured an appropriate number of research participants based on existing clinical studies, but to generalize our findings, reinforcement is needed further clinical studies are required.

## Conclusions

A clinical trial was conducted to investigate the removal efficiency of dental plaque on the tooth surface of an augmented reality-based smart toothbrush.

1. The reduction effect of the overall plaque index seemed to decrease more than when using the augmented reality toothbrush.
2. When the parts deviated, there was no significant difference when brushing the anterior part, but there was a significant difference in the removal of dental plaque through brushing in the posterior part.

Through the results of this study, it was possible to confirm the possibility of using augmented reality as a media for toothbrushing education, and it is expected that it can be used as a tool for oral health education as a tool for motivation.

## Acknowledgements

This research was funded by the National Research Foundation of Korea (NRF), grant number 2021R1F1A1064231. The NRF was funded by the Korean government (MSIT).

## Conflicts of Interest

The authors declared no conflicts of interest.

## Authorship

Conceptualization: JY Lee; Data collection: MJ Park, JY Lee; Formal analysis: MJ Park, SB Jang, JY Lee; Writing-original draft: MJ Park, SB Jang, JY Lee; Writing-review&editing: MJ Park, SB Jang, JY Lee

## References

1. Korea health insurance review & assessment service, Outpatient benefits by frequency of disease [Internet]. Korea health insurance review & assessment service;2023[cited 2022 Dec 31], Available from: <https://opendata.hira.or.kr/op/opc/olapHifrqSickInfo.do>.
2. Cho HJ, Ma DS, Lee KJ, Jeong SH, Jung HW, Choi YH et al. 2018 Children's oral health survey improvement plan and pre-planning research. Osong: Korea Ministry of Health and Welfare; 2018: 45.
3. Thylstrup A, Bruun C, Holmen L. In vivo caries models-mechanisms for caries initiation and arrestment. *Adv Dent Res* 1994;8(2):144-57. <https://doi.org/10.1177/08959374940080020401>
4. Kidd EAM, Fejerskov O. What constitutes dental caries? Histopathology of carious enamel and dentin related to the action of cariogenic biofilms. *J Dent Res* 2004;83(1):35-8. <https://doi.org/10.1177/154405910408301s07>
5. Starke EM, Mwatha A, Ward M, Argosino K, Jenkins W, Millemam J, et al. A comparison of the effects of a powered and manual toothbrush on gingivitis and plaque: a randomized parallel clinical trial. *J Clin Dent* 2019;30(1):24-9.
6. Colombo APV, Tanner ACR. The role of bacterial biofilms in dental caries and periodontal and peri-implant diseases: a historical perspective. *J Dent Res* 2019;98(4):373-85. <https://doi.org/10.1177/0022034519830686>
7. Sowinski J, Petrone DM, Wachs GN, Chaknis P, Kemp J, Sprosta AA, et al. Efficacy of three toothbrushes on established gingivitis and plaque. *Am J Dent* 2008;21(6):339-45.
8. Radzki D, Wilhelm-Węglarz M, Pruska K, Kusiak A, Ordyniec-Kwaśnica I. A fresh look at mouthwashes-what is inside and what is it for? *Int J Environ Res Public Health* 2022;19(7):3926. <https://doi.org/10.3390/ijerph19073926>
9. Tolvanen M, Lahti S, Hausen H. Changes in toothbrushing frequency in relation to changes in oral health-related knowledge and attitudes among children-a longitudinal study. *Eur J Oral Sci* 2010;118(3):284-9. <https://doi.org/10.1111/j.1600-0722.2010.00737.x>
10. Kay E, Locker DA. Systematic review of the effectiveness of health promotion aimed at improving oral health. *Community Dent Health* 1998;15(3):132-44.



11. Andrew R, Gay D, Maryann W, Maryann C, Randy J, Ronald L, et al. Six-month comparison of powered versus manual toothbrushing for safety and efficacy in the absence of professional instruction in mechanical plaque control. *J Periodontol* 2002;73(7):770-8. <https://doi.org/10.1902/jop.2002.73.7.770>
12. Javornik A. Augmented reality: research agenda for studying the impact of its media characteristics on consumer behaviour. *J Retail Consum Serv* 2016;30:252-61. <https://doi.org/10.1016/j.jretconser.2016.02.004>
13. Park MJ, Yoo JM. Effects of perceived interactivity of augmented reality on consumer responses: a mental imagery perspective. *J Retail Consum Serv* 2020;52(3):1-9. <https://doi.org/10.1016/j.jretconser.2019.101912>
14. Duta M, Amariei CI, Bogdan CM, Popovici DM, Ionescu N, Nuca CI. An overview of virtual and augmented reality in dental education. *Oral Health Dent Manag* 2011;10(1):42-9.
15. Jensen O, Gabre P, Skold UM, Birkhed D. Fluoride toothpaste and toothbrushing; knowledge, attitudes and behaviour among Swedish adolescents and adults. *Swed Dent J* 2011;35(4):203-13.
16. Dos Santos APP, de Oliveira BH, Nadanovsky P. A systematic review of the effects of supervised toothbrushing on caries incidence in children and adolescents. *Int J Paediatr Dent* 2018;28(1):3-11. <https://doi.org/10.1111/ipd.12334>
17. Pujar P, Subbareddy VV. Evaluation of the tooth brushing skills in children aged 6-12 years. *Eur Arch Paediatr Dent* 2013;14(4):213-9. <https://doi.org/10.1007/s40368-013-0046-3>
18. Berzinski M, Morawska A, Mitchell AE, Baker S. Parenting and child behaviour as predictors of toothbrushing difficulties in young children. *Int J Paediatr Dent* 2020;30(1):75-84. <https://doi.org/10.1111/ipd.12570>
19. Shin SJ, Ryu DY, Bae SM, Choi YK. Evaluation on the oral health promotion program effect from a part of community child center. *J Dent Hyg Sci* 2011;11(3):163-71.
20. Hwang GJ, Wu PH, Chen CC, Tu NT. Effects of an augmented reality-based educational game on students' learning achievements and attitudes in real-world observations. *Interact Learn Environ* 2016;24(8):1895-906. <https://doi.org/10.1080/10494820.2015.1057747>
21. Lai AF, Chen CH, Lee GY. An augmented reality based learning approach to enhancing students' science reading performances from the perspective of the cognitive load theory. *Br J Educ Technol* 2019;50(1):232-47. <https://doi.org/10.1111/bjet.12716>
22. Moro C, Phelps C, Redmond P, Stromberga Z. HoloLens and mobile augmented reality in medical and health science education: a randomised controlled trial. *Br J Educ Technol* 2021;52(2):680-94. <https://doi.org/10.1111/bjet.13049>
23. Logeswaran A, Munsch C, Chong YJ, Ralph N, McCrossnan J. The role of extended reality technology in healthcare education: towards a learner-centred approach. *Future Healthc J* 2021;8(1):e79-84. <https://doi.org/10.7861/fhj.2020-0112>
24. Killoy WJ, Love JW, Love J, Fedi Jr PF, Tira DE. The effectiveness of a counter-rotary action powered toothbrush and conventional toothbrush on plaque removal and gingival bleeding: a short term study. *J Periodontol* 1989;60(8):473-7. <https://doi.org/10.1902/jop.1989.60.8.473>
25. Addy M, Renton-Harper P, Warren P, Newcombe RG. An evaluation of video instruction for an electric toothbrush: comparative single-brushing cross-over study. *J Clin Perio* 1999;26(5):289-93. <https://doi.org/10.1034/j.1600-051X.1999.260505.x>
26. Yoo SY, Chon JA, Seo JH, Moon HR, Jwa SK. Oral health education effect on primary school children. *Int J Clin Prev Dent* 2012;8(4):225-9.
27. Kim HK, Bae SM, Shin SJ, Ryu DY, Son JH, Eom MR, et al. The comparison of effect in oral health education frequency for elementary school students from a part of community child center. *J Korean Soc Dent Hyg* 2011;11(5):759-71.
28. Rajchanovska D, Zafirova IB. Oral habits among pre-elementary children in Bitola. *Contributions* 2012;33(1):157-69.
29. Ekuni D, Yamanaka R, Yamamoto T, Miyauchi M, Takata T, Watanabe T. Effects of mechanical stimulation by a powered toothbrush on the healing of periodontal tissue in a rat model of periodontal disease. *J Periodontol Res* 2010;45(1):45-51. <https://doi.org/10.1111/j.1600-0765.2009.01195.x>
30. Amantini SNSR, Montilha AAP, Antonelli BC, Leite KTM, Rios D, Cruvinel T, et al. Using augmented reality to motivate oral hygiene practice in children: protocol for the development of a serious game. *JMIR Res Protoc* 2020;17;9(1):e10987. <https://doi.org/10.2196/10987>
31. Patil S, Hedad IA, Jafer AA, Abutaleb GK, Arishi TM, Arishi SA, et al. Effectiveness of mobile phone applications in improving oral hygiene care and outcomes in orthodontic patients. *J Oral Bio Craniofac Res* 2021;11(1):26-32. <https://doi.org/10.1016/j.jobcr.2020.11.004>
32. Spieler EL. Preventing toothbrush abrasion and the efficacy of the alert toothbrush: a review and patient study. *Compend Contin Educ Dent* 1996;17(5):478-80.
33. Jeon BJ, Oh JS, Son SM. Effects of tooth brushing training, based on augmented reality using a smart toothbrush, on oral hygiene care among people with intellectual disability in Korea. *Healthcare* 2021;9(3):348. <https://doi.org/10.3390/healthcare9030348>
34. Rodrigues CR, Ando T, Singer JDM, Issáo M. The effect of training on the ability of children to use dental floss. *ASDC J Dent Child* 1996;63(1):39-41.

35. Yunusovich MB, Ahadovich SA. Use of mouthwashes in the treatment of chronic catarrhal gingivitis in children. Middle European Scientific Bulletin 2020;5:96-8. <https://doi.org/10.47494/mesb.2020.5.69>
36. Ciancio, Sebastian G. The dental water jet: a product ahead of its time. Compend Contin Educ Dent 2009;30(1):7-13.
37. Sandlund M, Lindh Waterworth E, Häger C. Using motion interactive games to promote physical activity and enhance motor performance in children with cerebral palsy. Dev Neurorehabil 2011;14(1):15-21. <https://doi.org/10.3109/17518423.2010.533329>

## 증강현실 기반 스마트 칫솔의 치면세균막 제거효과 평가

### 초록

**연구목적:** 본 연구는 스마트 폰을 활용한 증강현실의 기반의 칫솔질 가이드 프로그램을 이용하여 어린이의 치면세균막 제거 효율이 어떻게 변화하는지 알아보려고 하였다. **연구방법:** 이중맹검 교차임상시험을 통하여, 5-12세 어린이 20명을 대상으로 일반 칫솔질 후 치면세균막 제거효율을 측정하고 워시아웃기간 이후, 증강현실 기반 칫솔질 수행 후 치면세균막 제거효율을 평가하여 비교하였다. **연구결과:** 전반적으로 증강현실 기반 양치질이 일반 양치질보다 플라그 제거 효율이 우수하였다( $p < 0.05$ ). 세부적으로 분류하면 전치부에서는 두 가지 칫솔질 방법 간 플라그 제거율에서 유의한 차이가 없었으나( $p = 0.056$ ), 구치부에서는 플라그 제거율에서 유의한 차이가 있음을 확인하였다( $p < 0.05$ ). **결론:** 증강현실 기반을 기반으로 스마트폰을 이용한 양치질에 대한 치태 제거 효능을 확인하였으며, 향후 ICT 기술을 접목한 구강보건교육에 활용할 수 있을 것으로 판단된다.

**색인 :** 증강현실, 치면세균막, 칫솔질