



Prevalence of chigger mites and *Orientia tsutsugamushi* strains in northern regions of Gangwon-do, Korea



Soojin Kim^{1,†}, In Yong Lee^{2,†}, Sezim Monoldorova¹, Jiro Kim^{1,3}, Jang Hoon Seo⁴, Tai-Soon Yong^{2,*},
Bo Young Jeon^{1,*}

¹Department of Biomedical Laboratory Science, College of Software and Digital Healthcare Convergence, Yonsei University, Wonju 26493, Korea; ²Department of Tropical Medicine and Institute of Tropical Medicine, Yonsei University College of Medicine, Seoul 03722, Korea; ³Department of Clinical Laboratory Science, Cheju Halla University, Jeju 63092, Korea; ⁴Department of Biomedical Laboratory Science, Shinhan University, Uijeongbu 11644, Korea

Abstract

Received: 26 October 2022
Accepted: 22 June 2023

*Correspondence
(tsy, tsyong212@yuhs.ac;
byj, bojeon@yonsei.ac.kr)

†These authors contributed equally
to this work.

Citation

Kim S, Lee IY, Monoldorova S, Kim J,
Seo JH, Yong TS, Jeon BY.
Prevalence of chigger mites and *Orientia
tsutsugamushi* strains in northern regions
of Gangwon-do, Korea.
Parasites Hosts Dis 2023;61(3):263-271.

The present study aimed to survey the prevalence of chigger mites and *Orientia tsutsugamushi* (*O. tsutsugamushi*) infection in the northern regions of Gangwon-do, Korea. From early February to early June 2015, a total of 17,050 chiggers were collected from striped field mice, *Apodemus agrarius*, in Cheorwon-gun, Hwacheon-gun, Yanggu-gun, and Goseong-gun, which are well-known endemic areas of scrub typhus in Korea. The chiggers were analyzed using molecular genomic methods, as previously described. Among the 7,964 identified chiggers, the predominant species was *Leptotrombidium pallidum* (76.9%), followed by *L. zetum* (16.4%), *L. orientale* (4.3%), *L. palpale* (0.3%), *L. tectum* (0.2%), and *Neotrombicula tamiyai* (1.8%). The chigger index (CI) was highest in Hwacheon (115.58), followed by Cheorwon (97.02), Yanggu (76.88), and Goseong (54.68). Out of the 79 *O. tsutsugamushi*-positive chigger pools, 67 (84.8%) were identified as the Boryong strain, 10 (12.7%) as the Youngworl strain, and only 2 were the Jecheon strain. Based on the high infestation of chiggers in striped field rodents and the high rate of *O. tsutsugamushi* infection in chigger mites, Hwacheon-gun and Cheorwon-gun are presumed to be high-risk areas for scrub typhus. Furthermore, *L. pallidum*, a major vector of scrub typhus, and the dominant *O. tsutsugamushi* serotype, the Boryong strain, were found in the northern regions of Gangwon-do, Korea.

Keywords: *Orientia tsutsugamushi*, chiggers, scrub typhus, strain, Korea

Introduction

Scrub typhus is caused by the obligate intracytosolic bacteria *Orientia tsutsugamushi*, which is transferred to humans by the biting of the larvae of numerous species of trombiculid mites. Scrub typhus (*tsutsugamushi* illness) is found in Asia-Pacific countries such as Korea, Japan, China, Thailand, Taiwan, and India [1-3]. Scrub typhus is the most frequent acute febrile illness in Korea in the autumn, with the number of cases peaking between October and November [4]. It occurs across Korea, particularly in the southern provinces of Jeolla-do, Gyeongsang-do, and Chungcheongnam-do [5]. In Korea, it is one of the most frequent arthropod-borne infections.

More than 20 serotypes of *O. tsutsugamushi* have been discovered based on antigenic features; the virulence of *O. tsutsugamushi* depends on the serotype or strain, and the prevalence of serotypes varies by nation. *O. tsutsugamushi* has 3 primary prototypes: Karp, Gilliam, and Kato serotypes [6]. Several strains of *O. tsutsugamushi* have been recorded in

© 2023 The Korean Society for
Parasitology and Tropical Medicine

This is an Open Access article distributed under
the terms of the Creative Commons Attribution
Non-Commercial License (https://creativecommons.org/licenses/by-nc/4.0) which permits
unrestricted non-commercial use, distribution,
and reproduction in any medium, provided the
original work is properly cited.

Author contributions

Conceptualization: Kim S
 Formal analysis: Kim S, Lee IY, Sezim M
 Investigation: Sezim M, Kim J, Seo JH
 Project administration: Sezim M, Kim J,
 Seo JH
 Resources: Kim S, Kim J
 Writing – original draft: Lee IY, Jeon BY
 Writing – review & editing: Lee IY,
 Yong TS, Jeon BY

Conflict of interest

The authors declare no conflict of interest related to this study.

ORCID

Soojin Kim
 (<https://orcid.org/0009-0000-4448-6872>)
 In Yong Lee
 (<https://orcid.org/0000-0001-9821-7504>)
 Sezim Monoldorova
 (<https://orcid.org/0000-0002-8869-7062>)
 Jiro Kim
 (<https://orcid.org/0000-0002-7352-680X>)
 Jang Hoon Seo
 (<https://orcid.org/0000-0001-5986-8590>)
 Tai-Soon Yong
 (<https://orcid.org/0000-0002-3445-0769>)
 Bo Young Jeon
 (<https://orcid.org/0000-0002-4314-216X>)

Korea, including the 3 important prototype strains Gilliam, Karp, and Kato, as well as strains (Boryong, Wonju, Jecheon, Pajoo, and Youngworl, Kanda, Oishi, OI011, Taguchi, and Shimokoshi) [7-9]. The Boryong strain is found in the southern and central parts of Korea, whereas the Gilliam, Karp, and Pajoo strains are mostly found in the central region [10]. The epidemiological features of *O. tsutsugamushi* strains are determined by the vector competence of chiggers in each region. There have been few investigations on chiggers and *O. tsutsugamushi* infections in Gangwon-do, where scrub typhus is uncommon. The investigation of the *O. tsutsugamushi* infection in Gangwon-do, Korea, which is near the Demilitarized Zone, is necessary.

We collected chigger mites from striped field mice, *Apodemus agrarius*, to explore *O. tsutsugamushi* infection in chigger mites, the vectors of scrub typhus in Gangwon-do, Korea.

Materials and Methods

Animals

The trapping of wild rodents was conducted in accordance with the guidelines of the Institutional Animal Care and Use Committee at Yonsei University Mirae Campus, Korea, and was approved by the committee (Approval No., YWC-151203). Standard procedures, as previously described [11], were followed for the collection and transportation of specimens to minimize hazards from potentially infected rodents.

Surveillance localities and periods

Wild rodents were captured in 4 locations (Cheorwon; 38° 26' 92", 127° 16' 93", Hwacheon; 38° 07' 63", 127° 52' 12", Yanggu; 38° 06' 24", 128° 16' 89", Cheorwon; 38° 26' 92", 127° 16' 93", and Goseong; 38° 29' 27", 128° 49' 57") in Gangwon-do, close to the Demilitarized Zone (Supplementary Fig. S1). Wild rats were gathered from rice paddies, canals, hills, grasslands, reservoirs, and river banks.

Collection of wild rodents

Sherman live folding traps (7.7×9×23 cm; H.B. Sherman Traps, Tallahassee, FL, USA) with peanut butter biscuit bait were used to capture wild rats. Traps were placed 4–5 meters apart from 15: 00 to 18: 00 and removed at 07: 00 the next morning. Each trap was put in a safe container and delivered to the laboratory with the rat inside. The seized wild rats were carbon dioxide killed and identified using morphological keys: gender, weight, and body length.

Collection of chiggers

Chiggers were detached from the ears of wild rats collected using delicate forceps and a needle under a dissecting microscope. Chiggers put on a slide glass using polyvinyl alcohol mounting media (BioQuip Products, Rancho Dominguez, CA, USA). Chiggers were identified using morphological standard criteria after being viewed under a stereomicroscope [12]. The number of chigger mites per rodent was used to compute the chigger index (CI).

Chigger homogenization and DNA extraction

One to 30 chiggers collected from each animal host were pooled and transferred to a reinforced homogenizing tube (Bertin Technology, Montigny-le-Bretonneux, France) containing 2.8 mm zirconium oxide beads (Bertin Technology) and 200 μ l of sterile phosphate-buffered saline. The pooled chiggers were homogenized using a bead beater, Precellys 24 (Bertin Technology), with 2 cycles of 20 sec at 6,500 rpm. The genomic DNA was extracted from chiggers using the G-spin Genomic DNA Extraction Kit for Bacteria (iNtRON Biotechnology, Seongnam, Korea) according to the manufacturer's instructions. The extracted DNA was stored at -20°C until further use.

Detection of *O. tsutsugamushi* using nested PCR

Nested PCR was used to detect the 56-kDa type-specific antigen of *O. tsutsugamushi* using the Tsutsugamushi Detection Kit (TSUTSU, iNtRON) according to the manufacturer's instructions. Outer primer set: forward primer (5'-GCAATATTGCTAGTGCAATGTCTGC-3') and reverse primer (5'-ATGCATGCATGRCGCTKCAATTTA-3'); inner primer set: forward primer (5'-ATAGGCCTATAAGTATWGCKGATCG-3') and reverse primer (5'-CATCTAGAYGCACTATTAGGCAAA-3'). The first primer set was used for amplification, which included an initial denaturation at 94°C for 5 min, 40 cycles of 94°C for 30 sec, annealing at 58°C for 30 sec, elongation at 72°C for 40 sec, and a final extension at 72°C for 5 min. Two μ l of the first PCR product served as the template for the second round of PCR, which was carried out in 30 cycles under the same circumstances as the first. The PCR products were separated on a 1.5% agarose gel and visualized using a UV transilluminator.

DNA sequencing and phylogenetic analysis

A LaboPass gel and PCR clean-up kit (Cosmogenetech Inc., Seoul, Korea) were used to purify and sequence the PCR products. BLAST (<http://ncbi.nlm.nih.gov/blastn>) identified the amplified sequence as a 56-kDa type-specific antigen (TSA) gene. The 56-kDa type-specific antigen (TSA) sequences of *O. tsutsugamushi* were aligned using MUSCLE [13] and the MEGA X program's unweighted pair group method with arithmetic mean (UPGMA). The phylogenetic analysis was carried out using the MEGA X program's bootstrap and the maximum likelihood tree [15]. MEGA 6.0 was used to determine the genetic distance, and the topologies were assessed using a bootstrap analysis with 1,000 iterations [17].

Results

Collection of wild rodents by regions

Between February 2015 and early June 2015, a total of 186 wild rodents were captured in Cheorwon, Hwacheon, Cheorwon, Yanggu, and Goseong in northern Gangwon-do, Korea. The captured wild rodents belonged to 4 species (3 genera) of the Muridae family: *A. agrarius*, *A. peninsulae*, *Micromys minutus*, and *Myodes regulus*. Among these species, *A. agrarius* was the most frequently captured, accounting for 90.8% ($n = 169$) of the total, followed by *Myodes regulus* (4.8%, $n = 9$), *Micromys minutus* (3.2%, $n = 6$), and *A. peninsulae* (1.1%, $n = 2$) (Table 1). The majority of wild rodents were captured in Hwacheon ($n = 106$, 57.0%), followed by Cheorwon ($n = 43$, 23.1%), Goseong ($n = 19$, 10.2%), and Yanggu ($n =$

Table 1. The collection results of wild rodents in northern regions of Gangwon-do from February to June 2015

Locality	<i>Apodemus agrarius</i>	<i>Apodemus peninsulae</i>	<i>Micromys minutus</i>	<i>Myodes regulus</i>	Total (%)
Hwacheon	97	2	2	5	106 (57.0)
Cheorwon	36	0	3	4	43 (23.1)
Yanggu	17	0	1	0	18 (9.7)
Goseong	19	0	0	0	19 (10.2)
Total (%)	169 (90.8)	2 (1.1)	6 (3.2)	9 (4.8)	186 (100.0)

Table 2. Chigger infestation status of *Apodemus agrarius* collected in northern regions of Gangwon-do from February to June 2015

Locality	No. of infested/ no. of captured (%)	No. of chiggers	Chigger indices (CI) ^a
Hwacheon	90/97 (92.8)	11,211	115.58
Cheorwon	25/36 (69.4)	3,493	97.02
Yanggu	17/17 (100.0)	1,307	76.88
Goseong	18/19 (94.7)	1,039	54.68
Total (%)	150,169 (88.8)	17,050	100.89

^aChigger indices (CI), the number of chigger mites per collected rodent.

Table 3. Detection results of trombiculid mites in striped field mice, *Apodemus agrarius*, by the survey regions of Gangwon-do

Locality	Chigger species (%) ^a						Total (%)
	<i>L. pallidum</i>	<i>L. orientale</i>	<i>L. palpale</i>	<i>L. tectum</i>	<i>L. zetum</i>	<i>N. tamiyai</i>	
Hwacheon	3,756	276	24	16	1,144	66	5,282 (66.3)
Cheorwon	1,304	49	0	0	145	82	1,580 (19.8)
Yanggu	537	1	0	0	0	2	540 (6.8)
Goseong	527	16	0	0	19	0	562 (7.1)
Total (%)	6,124 (76.9)	342 (4.3)	24 (0.3)	16 (0.2)	1,308 (16.4)	150 (1.8)	7,964 (100)

^aHalf of the collected chigger mites were used for the identification of chigger species.

18, 9.7%) (Table 1).

Chigger infestation of *A. agrarius*

Among the wild rodents gathered in the research region, 17,050 chiggers were obtained from *A. agrarius*. 150 (88.8%) of the 169 *A. agrarius* were infested with larval mites (Table 2). Yanggu had the greatest infestation rate (100.0%, 17 of 17), while Cheorwon had the lowest (69.4%, 25 of 36). Hwacheon has the highest chigger index (CI) (115.58 CI), followed by Cheorwon (97.02 CI), Yanggu (76.88 CI), and Goseong (54.68 CI) (Table 2).

Identification of trombiculid mites

Half of the detached chiggers were submitted for identification (Table 3). The 7,964 chigger mites included 5 *Leptotrombidium* species and one *Neotrombicula* species. *Leptotrombidium pallidum* (*L. pallidum*) was the most common species (76.9%, $n = 6,124$), followed by *L. zetum* (16.4%, $n = 1,308$), *Neotrombicula tamiyai* (*N. tamiyai*) (1.8%, $n = 150$), *L. palpale*

Table 4. Results of detection and genotyping of *Orientia tsutsugamushi* in chiggers from northern regions of Gangwon-do

Locality	No. of tested chiggers ^a	No. of tested pools	No. of positive pools	<i>O. tsutsugamushi</i> strain			Minimum positive rate (%; 95% CI)	Maximum positive rate (%; 95% CI)
				Boryong	Youngworl	Jecheon		
Hwacheon	5,929	242	38	28	10	0	0.64 (0.45–0.88)	15.70 (11.36–20.91)
Cheorwon	1,913	73	30	30	0	0	1.57 (1.06–2.23)	41.09 (29.71–53.23)
Yanggu	767	28	9	8	0	1	1.17 (0.54–2.22)	32.14 (15.88–52.35)
Goseong	477	24	2	1	0	1	0.42 (0.05–1.51)	8.33 (1.03–26.99)
Total (%)	9,086	367	79	67 (84.8)	10 (12.7)	2 (2.5)	0.87 (0.69–1.08)	21.52 (17.43–26.09)

^aHalf of the collected chigger mites were pooled, each pool consisted of 1-30 chigger mites and were used for the detection and genotyping of *O. tsutsugamushi*. Minimum positive rate, number of positive pool(s)/total number of the tested chigger mite × 100%; Maximum positive rate, number of positive pool(s)/total number of the tested chigger mite pools × 100%; 95% CI, 95% confidence interval.

(0.3%, $n = 24$), and *L. tectum* (0.2%, $n = 16$) *L. pallidum* was the most prevalent species in all research regions, with the largest geographical occurrence in Hwacheon (61.3%), Cheorwon (21.3%), Yanggu (8.8%), and Goseong (8.6%).


Detection and genotyping of *O. tsutsugamushi* in chiggers

For the identification of *O. tsutsugamushi*, 367 pools from 9,086 chiggers were submitted to nested PCR (Table 4). The bacteria were found in 79 of the 367 chigger pools (0.9% minimum positive rate, 24.5% maximum positive rate). Cheorwon had the highest minimum positive rate of 1.6% (30 out of 73), followed by Yanggu (1.2%, 9 out of 28), Hwacheon (0.6%, 38 out of 242), and Goseong (0.4%, 2 out of 24).

In 79 *O. tsutsugamushi*-positive chigger pools, 3 strains of *O. tsutsugamushi* were found. Boryong (84.8%, 67 of 79) was the most common *O. tsutsugamushi* strain, followed by Youngworl (12.7%, 10 of 79) and Jecheon strains (2.5%, 2 of 79). Boryong strain filled 73.7% of the 38 pools in Hwacheon, with 28 pools, while Youngworl strain occupied 26.3%, with 10 pools (Table 4). Boryong strains were discovered in 8 of 9 pools in Yanggu, accounting for 88.9%, while Jecheon strains were detected in 11.1% of pools in Goseong. Boryong and Jecheon strains were each detected in Goseong. In Cheorwon, however, only the Boryong strain was found. Boryong and Youngworl strains were found in the Hwacheon mountainous region, while only Boryong strains were found in the Cheorwon plains. Boryong and Jecheon strains were found in both Yanggu and Goseong, which were linked by mountain ranges (Table 4).

Phylogenetic analysis

A phylogenetic study of *O. tsutsugamushi* strains of chigger mites in northern Gangwon-do was undertaken, and it was categorized into 3 clusters, exhibiting Boryong, Jecheon, and Youngworl strains (Fig. 1). Boryong strains isolated from chigger mites in the Hwacheon, Yanggu, and Cheorwon areas correlated with the reference Boryong strain (GenBank accession numbers: AM494475). The M235 Jecheon strain (OQ656415) obtained from Goseong chiggers was similar to the reference Jecheon strain (AF430143), and the Youngworl strains M166, M148, M143, M129, and M128 (OQ656409, OQ656410, OQ656413, and OQ656414) obtained from Hwacheon chiggers were similar to the reference Youngworl strain (AF430141).



Discussion

The chigger mite is one of the most medically significant outdoor mites in Korea, and it mediates scrub typhus, which occurs annually and presents a public health hazard [1-7]. Chigger mites, *L. pallidum* and *L. scutellare*, are recognized as the primary vectors of scrub typhus in Korea.

We researched the chigger mite from collected wild rodents and the *O. tsutsugamushi* infection of the chigger mites in Hwacheon, Cheorwon, Yanggu, and Goseong in the northern regions of Gangwon-do, Korea, where scrub is prevalent. Hwacheon has the highest population of wild rodents and chigger infestations among the 4 areas. Not only was the chigger infestation high in Cheorwon, but so was the detection rate of *O. tsutsugamushi* chigger mites. These assume that Hwacheon and Cheorwon are scrub typhus hotspots. These findings show that chigger mite infestation and *O. tsutsugamushi* infection might aid in estimating the risk of scrub typhus.

During the spring season, when there are few scrub typhus cases, there is evidence of *O. tsutsugamushi* infection in chiggers. The most common chigger species was *L. pallidum* (76.9%), followed by *L. zetum* (16.4%), *L. orientale* (4.3%), *N. tamiyai* (1.8%), *L. palpale* (0.3%), and *L. tectum* (0.2%). The dominating species, *L. pallidum*, was the most common in Hwacheon (61.3%), followed by Cheorwon (21.3%), Yanggu (8.8%), and Goseong (8.6%). These findings are consistent with prior research that found *L. pallidum* to be the most common chigger mite in northern Gyeonggi-do [18] and Gangwon-do [19]. *L. pallidum* was followed by *L. orientale*, *L. scutellare*, and *L. palpale* in Gyeonggi-do, and *L. palpale*, *L. orientale*, and *L. zetum* in Gangwon-do [18,19]. In the current research, however, *L. scutellare* was not found, unlike in Gyeonggi-do, and *L. zetum* was the most prevalent after *L. pallidum*, with *N. tamiyai* and *L. tectum* also found in Gangwon-do. These variations might be attributable to regional and seasonal changes.

In Gangwon-do, the average chigger index was 100.89 and peaked in the spring. These findings seem to corroborate with a previous finding [20] that chigger indices spiked twice in the spring and fall. This might be explained by the fact that *L. pallidum* achieves its height between September and November, nearly vanishes during the winter, and returns to its peak between March and June [21]. This implies that *L. pallidum* is the most major vector of *O. tsutsugamushi* in Gangwon-do, northern Korea, in the spring, with other chiggers such as *L. zetum*, *L. orientale*, *N. tamiyai*, *L. tectum*, and *L. palpale* also implicated [21-23].

The Boryong strain accounted for 84.8% of the total strains in the *O. tsutsugamushi* genotyping study on chigger mites collected in northern regions of Gangwon-do, whereas Youngworl and Jecheon were minor strains. Boryong and Youngworl strains were dominant in the Hwacheon area, while Boryong and Jecheon strains were dominant in the Yanggu and Goseong regions. Only the Boryong strain was dominant in the Cheorwon area. Several strains of *O. tsutsugamushi* have been documented in Korea, including the 3 prototype strains, Gilliam, Karp, and Kato, as well as 3 native strains (Boryong, Pajoo, and Youngworl). Boryong is the most prominent strain and has been documented in the central and southern parts of Korea; Gilliam, Karp, Kato, Pajoo, and Youngworl strains have also been identified in the central region [10,24-26]. These results suggest that there is a correlation between the distribution of *O. tsutsugamushi* strains, and the population density and vector competence of chigger mites.

The phylogenetic analysis of *O. tsutsugamushi* strains in northern Gangwon-do was classified into three clusters: Boryong, Jecheon, and Youngworl were found to be similar to previous study results [6].

According to a report by the Korea Disease Control and Prevention Agency (KDCA; <https://is.kdca.go.kr>), the incidence of scrub typhus was low in northern Gangwon-do in the spring. The current research found that the infection rate of *O. tsutsugamushi* in chiggers was low in the spring, implying that the occurrence of scrub typhus in northern regions of Gangwon-do, was related to the rate of infection in chiggers in the spring.

Scrub typhus is most common in the southern and southwestern plains of Korea, particularly Jeonbuk (38.6%), Chungnam (32.7%), and Gyeongnam (9.8%) provinces [8]. On the other hand, since the current study's target area consists mostly of hilly areas in the country's central and northeastern regions, the prevalence of typhus looks to be low. Scrub typhus has a high frequency in Korea in the fall, especially from October to November, and

a low incidence in the spring [26–28], as seen in the current research. Extensive research on the *O. tsutsugamushi* strains and their frequency is required in the future, particularly on the relationship between vectors and people.

In conclusion, Hwacheon and Cheorwon are high-risk areas for scrub typhus, with a high prevalence of chigger infestation on striped wild rats and a high rate of *O. tsutsugamushi* infection in chiggers. *L. pallidum* was the predominant chigger species, and various *O. tsutsugamushi* strains, including the dominant Boryong strain, Youngworl strain, and Jecheon strain, were identified in chiggers from the northern regions of Gangwon-do, Korea.

Acknowledgements

We thank Namkyoung Kim and Jonguk Jeong for technical support. This study was supported by a grant from climate change vector surveillance programs of the Korea Disease Control and Prevention Agency (KDCA), and by a grant from Armed Forces Medical Research Institute, The Armed Forces Medical Command, Republic of Korea.

References

- Elliott I, Pearson I, Dahal P, Thomas NV, Roberts T, et al. Scrub typhus ecology: a systematic review of *Orientia* in vectors and hosts. *Parasit Vectors* 2019;12(1):513. <https://doi.org/10.1186/s13071-019-3751-x>
- Bhate R, Pansare N, Chaudhari Sp, Barbudde SB, Choudhary VK, et al. Prevalence and phylogenetic analysis of *Orientia tsutsugamushi* in rodents and mites from Central India. *Vector Borne Zoonotic Dis* 2017;17(11):749-754. <https://doi.org/10.1089/vdz.2017.2159>
- Ha JH, Lee DH, Park JS, Cho OH, Kim DH, et al. Isolation and genetic characterization of *Orientia tsutsugamushi* from scrub typhus patients in Gyeongsangnam-do, Korea. *J Bacteriol Virol* 2016;46(4):275-282. <https://doi.org/10.4167/jbv.2016.46.4.275>
- Roh JY, Song BG, Park WI, Shin EH, Park C, et al. Coincidence between geographical distribution of *Leptotrombidium scutellare* and scrub typhus incidence in South Korea. *PLoS One* 2014;9(12):e113193. <https://doi.org/10.1371/journal.pone.0113193>
- Chung MH, Kang JS. History of tsutsugamushi disease in Korea. *Infect Chemother* 2019;51(2):196-209. <https://doi.org/10.3947/ic.2019.51.2.196>
- Kelly DJ, Fuerst PA, Ching WM, Richards AL. Scrub typhus: the geographic distribution of phenotypic and genotypic variants *Orientia tsutsugamushi*. *Clin Infect Dis* 2009;48(suppl):203-230. <https://doi.org/10.1086/596576>
- Shim SK, Shin YK, Choi EN, Han BG, Lee HK, et al. Analysis of cellular fatty acids in *Orientia tsutsugamushi* as taxonomic markers. *Microbiol Immunol* 2005;49(4):343-347. <https://doi.org/10.1111/j.1348-0421.2005.tb03738.x>
- Choi YJ, Lee IY, Song HJ, Kim J, Park HJ, et al. Geographical distribution of *Orientia tsutsugamushi* strains in chiggers from three provinces in Korea. *Microbiol Immunol* 2018;62(9):547-553. <https://doi.org/10.1111/1348-0421.12639>
- Bahk YY, Ahn SK, Lee J, Kwon HW, Hong SJ, et al. Monitoring chigger mites for *Orientia tsutsugamushi* in field small mammals in Hwaseong-si, Gyeonggi-do, Korea, 2019-2020. *Korean J Parasitol* 2021;59(3):319-324. <https://doi.org/10.3347/kjp.2021.59.3.319>
- Shim SK, Choi EN, Yu KO, Park HJ, Kim CM, et al. Characterisation of *Orientia tsutsugamushi* genotypes from wild rodents and chigger mites in Korea. *Clin Microbiol Infect* 2009;15(suppl):311-312. <https://doi.org/10.1111/j.1469-0691.2008.02254.x>
- Mills JN, Corneli A, Young JC, Garrison LE, Khan AS, et al. Hantavirus pulmonary syndrome-United States: updated recommendation for risk reduction. *MMWR Recomm Rep* 2002;51(RR-9):1-12.
- Ree HI. Fauna and key to the chigger mites of Korea (Acarina: Trombiculidae and Leeuwenhoekidae). *Korea J Syst Zool* 1990;6:57-70.
- Edgar RC. MUSCLE: multiple sequence alignment with high accuracy and high throughput. *Nucleic Acids Res* 2004;32(5):1792-1797. <https://doi.org/10.1093/nar/gkh340>
- Kumar S, Stecher G, Li M, Nnyaz C, Tamura K. MEGA X: molecular evolutionary genetics analysis across computing platforms. *Mol Biol Evol* 2018;35(6):1547-1549. <https://doi.org/10.1093/molbev/msy096>
- Saitou N, Nei M. The neighbor-joining method: a new method for reconstructing phylogenetic trees. *Mol Biol Evol* 1987;4(4):406-425. <https://doi.org/10.1093/oxfordjournals.molbev.a040454>

16. Felsenstein J. Confidence limits on phylogenies: an approach using the bootstrap. *Evolution* 1985;39(4):783-791. <https://doi.org/10.1111/j.1558-5646.1985.tb00420.x>
17. Tamura K, Stecher G, Peterson D, Filipski A, Kumar S. MEGA6: molecular evolutionary genetics analysis Version 6.0. *Mol Biol Evol* 2013;30(12):2725-2729. <https://doi.org/10.1093/molbev/mst197>
18. Lee IY, Kim HC, Lee YS, Seo JH, Lim JW, et al. Geographical distribution and relative abundance of vectors of scrub typhus in the Republic of Korea. *Korean J Parasitol* 2009;47(4):381-386. <https://doi.org/10.3347/kjp.2009.47.4.381>
19. Lee IY, Lim JW, Seo JH, Kim HC, Lee KJ, et al. Geographical distribution and epidemiologic factors of chigger mites on *Apodemus agrarius* during autumn in Korea. *Korean J Parasitol* 2021;59(5):473-479. <https://doi.org/10.3347/kjp.2021.59.5.473>
20. Kim SY, Gill B, Song BG, Chu H, Park WI, et al. Annual fluctuation in chigger mite populations and *Orientia tsutsugamushi* infections in scrub typhus endemic regions of South Korea. *Osong Public Health Res Perspect* 2019;10(6):351-358. <https://doi.org/10.24171/j.phrp.2019.10.6.05>
21. Lee IY, Ree HI, Hong HK. Seasonal prevalence and geographical distribution of trombiculid mites (Acarina: Trombiculidae) in Korea. *Korean J Zool* 1993;36(3):408-415 (in Korean).
22. Ree HI, Chang WH, Kee SH, Lee IY, Jeon SH. Detection of *Orientia tsutsugamushi* DNA in individual trombiculids using polymerase chain reaction in Korea. *Med Entomol Zool* 1997;48(3):197-209. <https://doi.org/10.7601/mez.48.197>
23. Ree HI, Lee IY, Jeon SH, Yoshida Y. Geographical distribution of vectors and sero-strains of tsutsugamushi disease at mid-south inland of Korea. *Korean J Parasitol* 1997;35(3):171-179. <https://doi.org/10.3347/kjp.1997.35.3.171>
24. Kim HC, Lee IY, Cheong ST, Richards AL, Gu SH, et al. Sero-surveillance of scrub typhus in small mammals collected from military training sites near the DMZ, northern Gyeonggi-do, Korea and analysis of the relative abundance of chiggers from mammals examined. *Korean J Parasitol* 2010;48(3):237-243. <https://doi.org/10.3347/kjp.2010.48.3.237>
25. Shim SK, Shin YK, Choi EN, Han BG, Lee HK, et al. Analysis of cellular fatty acids in *Orientia tsutsugamushi* as taxonomic markers. *Microbiol Immunol* 2005;49(4):343-347. <https://doi.org/10.1111/j.1348-0421.2005.tb03738.x>
26. Park SW, Lee CK, Kwak YG, Moon C, Kim BN, et al. Antigenic drift of *Orientia tsutsugamushi* in South Korea as identified by the sequence analysis of a 56-kDa protein-encoding gene. *Am J Trop Med Hyg* 2010;83(4):930-935. <https://doi.org/10.4269/ajtmh.2010.09-0791>
27. Noh MS, Lee YJ, Chu CS, Gwack J, Youn SK, et al. Are there spatial and temporal correlations in the incidence distribution of scrub typhus in Korea? *Osong Public Health Res Perspect* 2013;4(1):39-44. <https://doi.org/10.1016/j.phrp.2013.01.002>
28. Chung MH, Kang JS. History of tsutsugamushi disease in Korea. *Infect Chemother* 2019;51(2):196-209. <https://doi.org/10.3947/ic.2019.51.2.196>