

REVIEW

Modern Concepts of Restructured Meat Production and Market Opportunities

Abdul Samad^{1,†}, AMM Nurul Alam^{1,†}, Swati Kumari¹, Md. Jakir Hossain¹, Eun-Yeong Lee¹, Young-Hwa Hwang², and Seon-Tea Joo^{1,2,*}

¹Division of Applied Life Science (BK21 Four), Gyeongsang National University, Jinju 52828, Korea

²Institute of Agriculture & Life Science, Gyeongsang National University, Jinju 52828, Korea


OPEN ACCESS

Received January 18, 2024

Revised February 13, 2024

Accepted February 14, 2024

***Corresponding author** : Seon-Tea Joo
 Division of Applied Life Science (BK21 Four), Gyeongsang National University, Jinju 52828, Korea
 Tel: +82-55-772-1943
 Fax: +82-55-772-1949
 E-mail: stjoo@gnu.ac.kr

***ORCID**

Abdul Samad
<https://orcid.org/0000-0002-4724-3363>
 AMM Nurul Alam
<https://orcid.org/0000-0003-3153-3718>
 Swati Kumari
<https://orcid.org/0009-0001-3330-7821>
 Md. Jakir Hossain
<https://orcid.org/0009-0008-7663-9202>
 Eun-Yeong Lee
<https://orcid.org/0000-0002-3467-7349>
 Young-Hwa Hwang
<https://orcid.org/0000-0003-3687-3535>
 Seon-Tea Joo
<https://orcid.org/0000-0002-5483-2828>

[†] These authors contributed equally to this work.

Abstract Restructured meat (RM) products are gaining importance as an essential component of the meat industry due to consumers' interest in health benefits. RM products imply the binding or holding of meat, meat by-products, and vegetable proteins together to form a meat product with meat's sensory and textural properties. RM products provide consumers with diversified preferences like the intake of low salt, low fat, antioxidants, and high dietary fiber in meat products. From the point of environmental sustainability, RM may aid in combining underutilized products and low-valued meat by adequately utilizing them instead of dumping them as waste material. RM processing technique might also help develop diversified and new hybrid meat products. It is crucial to have more knowledge on the quality issues, selection of binding agents, their optimum proportion, and finally, the ideal processing techniques. It is observed in this study that the most crucial feature of RM could be its healthy products with reduced fat content, which aligns with the preferences of health-conscious consumers who seek low-fat, low-salt, high-fiber options with minimal synthetic additives. This review briefly overviews RM and the factors affecting the quality and shelf life. Moreover, it discusses the recent studies on binding agents in processing RM products. Nonetheless, the recent advancements in processing and market scenarios have been summarized to better understand future research needs. The purpose of this review was to bring light to the ways of sustainable and economical food production.

Keywords restructured meat, hybrid meat, sensory quality, processing technique, market scenario

Introduction

Historically, meat has been a rich source of protein and an essential part of the human diet (Baugreet et al., 2018). The meat contains not only protein but also vital minerals and vitamins. Due to the growing population, Food and Agriculture Organization of the United Nations (FAO, 2014) has projected that worldwide meat consumption will double by 2050. In the next decade, the consumption growth is

assumed to be 14% (OECD and FAO, 2021). The rising income and population growth are critical factors for the increase in demand for meat (Alam et al., 2024). Fig. 1 describes the demand for meat with the increasing population. To meet the growing demand for meat in the coming years, there is a need for more resources for meat processing technology. Thus, to fulfill this meat demand, various alternatives are developed, including cultured meat (Post et al., 2020), hybrid cultured meat (Alam et al., 2024), hybrid meat (HBM; Baune et al., 2023), and plant-based meat (Kumari et al., 2024).

The processing of different ingredients (e.g., meat or vegetable protein) and transforming them into a new product with high nutritive values is known as restructured meat (RM; Poláček et al., 2021). Usually, the development of RM involves the inclusion of various additives and binding agents (Carpentieri et al., 2022). The prospect of RM is related to the sustainability of the meat industry by utilizing less used parts of meat or by-products as they are being wasted due to deficient demand. Thus, there is an opportunity to combine underutilized meat parts and conventional low-cost plant protein sources into a new restructured product (Freire et al., 2016). Various methods, such as meat restructuring and HBM, have been used to change the meat to include health benefits (functional components) from other sources, such as plants (Mireles-Arriaga et al., 2017) and other protein sources (Baugreet et al., 2018). RM could be a valuable solution for health-conscious consumers to reduce portions of meat in their diet.

Mark Post established the basis for a fundamental change in our comprehension of meat manufacturing and proposed the concept of alternative meat manufacturing in 2012 (Post, 2012). RM is such a kind of alternative processing method. In RM, products can be manufactured by reducing the size of low-value, underutilized meat particles with the help of different processing techniques, e.g., chopping, cutting, sectioning, tenderizing, flaking, and grounding. RM and HBM emulsions are often reformed into restructured steak, patties, and other reformed meat batter-like (Anna Anandh and Annal Villi, 2018). While producing these emulsions, different methods are used to turn meat into a fine slurry and mixed well with starch, fat,




Fig. 1. Global meat consumption trends. Adapted from Blaustein and Smith (2021) with permission of MIT Technology Review.

and different herbs and spices to improve the flavor of RM products. Researchers are continuously working to explore RM techniques, such as Schönfeldt and Strydom (2011), and explore the nutritional consequences of this non-traditional method. Further studies are required to improve the physiochemical and sensory qualities of meat. Nevertheless, any new technology might have pros and cons; RM is not an exception and has advantages and disadvantages (Freire et al., 2016), illustrated in Fig. 2.

RM is taking an entry to the consumers worldwide with variation according to culture and continents. For example, the production rate of RM in Korea is still low; however, in other parts of the world, people are working on restructuring technology to improve meat quality and utilize less valuable meat by combining it with different types of meat. Meat prices are increasing rapidly, and RM products would be cost-effective. For this reason, it is easy for meat consumers to adopt RM products (Gadekar et al., 2015). This review contains techniques and processes of RM and discusses various methods and approaches to overcome quality issues and binding efficiency in RM. It also includes the current developments and trends in RM.

Process of Restructured Meat

Processing of RM involves several steps. The basic principle of RM is to craft a mixture of various meat types or add some plant-based protein and fiber sources. This combination establishes the basis for attaining the end product's intended taste, consistency, and nutritional makeup (Patel et al., 2023). After meticulously choosing the meat or plant source, the vital stage is the grinding process. The mechanical process makes the mixture uniform and helps create the distinct texture of RM (Farouk, 2010).

Bio-adhesives, typically sourced from natural origins, are crucial in improving the cohesion and arrangement of RM. This stage entails carefully and precisely incorporating bio-adhesives to provide the best possible adhesion of meat particles, enhancing the end product's overall quality and stability. The production process of RM relies heavily on temperature and time factors. Ensuring the product's texture, flavor, and safety requires maintaining an ideal temperature for a certain period. The regulated environment guarantees the appropriate bio-adhesive curing and enhances the overall quality of the RM. In Fig. 3, the simple process of restructuring meat is elaborated, in which RM is made by combining meat and cereals.

Factors Associated with Qualities of Restructured Meat Products

Physical qualities of restructured meat products

As RM is frequently composed of a blend of several meat origins and occasionally includes plant-based components, it can display a wide variation in color, texture, and flavor. It is necessary to add some additives to boost the quality of RM. The




Fig. 2. Advantages and disadvantages of restructured meat.




Fig. 3. Schematic diagram showing the restructured meat manufacturing process.

selection of additives and processing procedures is essential in defining the product's ultimate color, taste, and texture. For example, adding natural colors like beetroot extract or annatto can enhance the visual appearance by creating a red or pink color that imitates the look of conventional meat. Nevertheless, the production of RM presents a complicated problem in attaining the appropriate color while preserving nutritional content and ensuring customer approval (Andrade et al., 2023).

The visual aspect of restructured steaks is a significant concern for consumers (Kumar et al., 2023). The fundamental criteria for restructured beef products are vibrant colors, meat resembling entire muscle steaks, and uniform dispersion of tiny fat particles (Cifuentes-Galindres et al., 2024). The mixing duration affects the meat's color and will accelerate the degradation of the intended color if it exceeds 12 minutes (Gómez et al., 2020). A study examined the efficacy of vacuum mixing in reducing color degradation in fresh meat. However, the final steaks had a less attractive surface color. Salt can produce discoloration in restructured steak (Mandigo and Osburn, 2019). According to Gadekar et al. (2015) higher amounts of salt lead to a drop in the desirability of color. Sodium tripolyphosphate (STP) can enhance the natural color by countering the impact of salt. Wang et al. (2021) stated that there is a correlation between discoloration and lipid oxidation of restructured steaks. The researchers discovered that the deterioration of color in beef happened earlier during storage compared to the oxidation of lipids. Furthermore, the oxidation of pigments may have facilitated the lipid oxidation process. Meanwhile, Serrano et al. (2006) found no correlation between color degradation and lipid oxidation.

Texture is crucial in determining the mouthfeel and overall sensory experience of meat products. RM is subjected to a rigorous process to replicate the fibrous composition observed in conventional cuts. Methods such as extrusion and texturization imitate the texture and tenderness commonly found in meat. The study conducted by Ribeiro et al. (2023) investigates the influence of different ingredients and processing conditions on the texture of restructured beef. The continual improvement of restructured beef products prioritizes achieving the correct texture while addressing problems such as

dryness or hardness. Hydrocolloids effectively make better meat texture (Dinani et al., 2023). Thus, they can be added to RM products to enhance meat quality.

Sensory qualities of restructured meat products

The taste of RM plays a crucial role in the culinary experience, as it aims to accurately mimic traditional meat's flavor while potentially introducing other functional qualities. The combination of various components, utilization of taste intensifiers, and the culinary procedure collectively influence the ultimate flavor composition. Chen et al. (2021) investigate techniques to augment umami, the savory flavor commonly linked to meat, in restructured food items to bring positive consumer variation. Moreover, the possibility of enhancing the sensory experience of RM by using herbs, spices, and alternative protein sources is an exciting opportunity to introduce new and varied flavors. Salts like NaCl can be used to improve the texture and taste of meat (Cornet et al., 2021). Gyrating and kneading aim to generate favorable characteristics in the final output (Zhang et al., 2024).

Tumbling, homogenizing, or proper mixing enhances the dispersion of all ingredients and augments the ionic strength and pH. This leads to increased product yield, improved water retention, tenderness, juiciness, final product appearance, and better binding of meat chunks. The study conducted by Tsafraqidou et al. (2023) found that the product yield from vacuum and aerobically tumbled meats was similar. However, Muller (1991) reported a higher product yield for restructured chicken meat products. Muller also found that using 10% water chestnut flour (hydrated at a 1:1 ratio) was optimal for these products. Furthermore, the products remained acceptable for up to 10 days when stored in refrigerated conditions. It was confirmed that restructured cooked ham formulation could incorporate up to 38% of fresh liquid whey, yielding comparable outcomes to products cured using a traditional formulation (Dutra et al., 2012). Another use of RM was incorporated to increase the chicken meat yield by treating some parts of 5.0%, 7.5%, and 10.0% Hydrated Colocasia Flour (HCF). The most effective amount of HCF to use was determined to be 7.5% based on sensory evaluations, physicochemical qualities, and microbiological quality (Talukder et al., 2013). The appropriateness of tetra-potassium pyrophosphate (0.4%), tetra-sodium pyrophosphate (0.4%), and their combination (0.2% each) was assessed for the production of a low-salt reformed goat meat product. Using phosphates and including tetrapotassium pyrophosphate, which has a soapy smell, resulted in a substantial enhancement in the product output. Combining tetra-sodium and tetra-potassium pyrophosphate can create a low-sodium product (Gadekar et al., 2014). Reducing the duration of tumbling before injecting can enhance the yield and tenderness of roast beef (Boles and Shand, 2002). Table 1 provides an overview of the key factors influencing the quality of RM products for better understanding on future work to improve processing and RM product development.

Meat Particle Size and Product Quality

The particle size can significantly affect the texture of RM products. Several studies have been done to check the effect of meat particle size on meat quality. Restructured pork steaks made from smaller flakes exhibited reduced shear force values and were preferable to consumers (Patel et al., 2023). Pork chops and pork shoulder products that were restructured and manufactured with smaller flakes exhibited more excellent softness (Bhaskar Reddy et al., 2015).

It was observed that decreasing the size of meat particles can enhance the quality of RM products. Sen and Karim (2003) found that reducing the mutton particle size from 0.7–1.2 mm performed better than 20 mm cut while producing restructured mutton chops. The restructured pork blocks with a chunk size of 2–3 cm exhibited a more significant product yield (89.31%)

Table 1. Factors affecting the quality of restructured meat

Factors	Observations	References
Selection of ingredient	The selection of different ingredients affects the overall restructured meat quality positively or negatively. Using NaCl as a binding agent in meat can affect the flavor of meat.	Zhou et al., 2020
Processing techniques	Processing techniques can affect the overall quality of meat. It was observed that particle size in restructured meat affects the texture.	Xia et al., 2023
Temperature and time	A range from -2°C to -10°C was found as the ideal temperature for good binding when kept for five hours. Temperature fluctuations have been seen to affect the binding ability and quality of restructured meat.	Sheard, 2002
Additives and binders	Walnuts can be used as additives because it was observed that walnuts were able to improve the color of restructured meat.	Florowski et al., 2019
	MTgase can bind meat particles together, and in this way, it plays a crucial role in improving the texture.	Xu and Xu, 2021
	Acacia gum can also help in the preservation of restructured meat.	Sharma et al., 2014
Packaging and storage	In a study, packaging and storage were found to be very important variables, and deviation from standards could be vulnerable to meat microbial contaminations.	Sofos, 2014
	Preservatives that can be used to improve the shelf quality of meat. Alginate is a plant-based alternative and can be used as a preservative of restructured meat without affecting its taste.	Sha and Xiong, 2020

than those with a chunk size of 4–5 cm (85.12%). The pork block rebuilt using meat chunks measuring 2–3 cm exhibited a considerably lower shear force value and a higher tenderness level than those produced with 4–5 cm meat chunks (Gurikar et al., 2014).

Binding Agents in Processing of Restructured Meat

It is necessary to add binding agents to bind meat particles together to form products properly. Besides muscle proteins, additional non-meat components significantly bind meat chunks or bits. Meat chunks containing 0.1% microbial transglutaminase (MTGase) and 3% salt exhibited superior binding. The enzyme MTGase (0.05%–0.1%) and sodium caseinate (0.5%–0.1%) effectively made RM with sufficient binding in its raw, refrigerated form without adding salt at 5°C for 2 hours (Kuraishi et al., 1997). In another study, the use of beef rolls cooked with 1% salt and 0.5% sodium salt of phytate was shown to be more efficient than using 1% salt+0.5% sodium pyrophosphate, 0.5% STP in enhancing binding strength and cooked yield (Lee et al., 1998). To increase the strength of binding sites, adding 0.5% calcium lactate, 0.5% algin, 0.5 phosphate, and 1.5% salt gave significant results (Shao et al., 1999). They further noted that binding capacity was better than the previous treatment (5% fibrinogen and 0.25% thrombin).

An effective combination is crucial to developing an effective RM product, and researchers are conducting studies on this issue. A study by Devatkal and Mendiratta (2001) found that the most practical combination of binding agents for producing restructured pork rolls under refrigeration was 0.7% sodium alginate, 0.125% calcium carbonate, and 0.3% calcium lactate. The utilization of tri-calcium phosphate at a concentration of 0.3% resulted in notable enhancements in both the tenderness of flesh and the binding properties of restructured buffalo meat rolls, as compared to products containing 0.3% STP (Mendiratta et al., 2002). A formulation with 1% soybean protein isolates, 0.3% carrageenan, and 3% potato starch can make cheap RM because these binders are more affordable than others (Silva et al., 2021). Fiber-based binders also have potential in RM

Table 2. Commonly used materials in the processing of restructured meat

Source	Material	Observations	References
Chemical	Calcium alginate	Examined that it helped in restructuring and improving the quality of meat.	Boles, 2011
	Sodium triphosphate	Have the potential to improve the structure of meat	Hu et al., 2021
	MT Gase, caseinate	Efficient to increase the binding ability of meat	Chen et al., 2024
	Calcium lactate	Can be used as a salt substitute and improved the taste	Wang, 2023
	Sodium chloride aqueous solution	Flavor and texture quality were enhanced	Zhang et al., 2023
Plant	Alginate	Used to enhance the shelf life of meat	Montone et al., 2023
	Carrageenan	Improved the microstructure of meat and may improve RM product quality.	Feng et al., 2024
	Soya bean protein isolates	Helped in improving the crosslinking between meat particles	Wei et al., 2023
	Walnut	Improved color and texture	Afshar and Seyyed Qavami, 2023
	Acacia gum	Useful as an emulsifier in RM	Inguglia et al., 2023
Animal	Blood plasma	Aided the binding capacity of meat	Zou et al., 2019

RM, restructured meat.

processing and show good binding strength without affecting the quality of RM (Shafit et al., 2007). Summarized information on the commonly used in the processing of RM is shown in Table 2, which could bring effective physical and sensory characteristics.

Factors Mediate the Binding Strength

Along with including a binding agent, it is essential to have a sufficient capacity for binding multiple materials to form the final product successfully. A study conducted by Serrano et al. (2007) found that microwaved restructured beef steaks exhibited considerably higher Kramer shear force and binding strength compared to steaks that were pan-fried or normally oven-cooked ($p < 0.05$). According to Sharma et al. (2014), vacuum tumbling buffalo meat at a pressure of 0.4 bar for 3 hours at a speed of 11 rpm resulted in improved extraction of salt soluble proteins and enhanced binding and cohesiveness of the product compared to aerobic tumbling for the same duration.

To provide the best possible binding and food safety, mechanical measures like performing tumbler and massager processes in cold rooms at 0 degrees Celsius are recommended (Sikorski, 2004). The protein exudates generated during the procedure act as a binding agent, adhering the meat chunks together while cooking. The use of a vacuum during the operation aids in the elimination of air bubbles from the exudates and facilitates the extraction of proteins (Barbut, 2005).

Techniques to Improve Shelf Life

As meat products are perishable and require storage at a minimum temperature of -18°C , enhancing their shelf life during processing is necessary. Thus, for this reason, different approaches and designs were used to enhance the shelf life of RM products. Using grape seed extract at a concentration of 0.1% improved the duration for which reconstructed mutton slices may be stored without spoiling and can be refrigerated for up to 28 days (Bhaskar Reddy et al., 2013). The inclusion of

additives can improve the overall quality of meat products. A study conducted by Gadekar et al. (2014) showed that the addition of sodium ascorbate (500 ppm) and alpha-tocopherol acetate (10 ppm) to restructured goat meat products enhanced the stability of lipids during refrigeration and frozen storage, therefore improving the overall storage stability of the meat. Results from Reverte et al. (2003) showed a positive effect of adding flavoring compounds and antioxidants as they improved beef's shelf life and flavor, which can be incorporated in RM processing. To evaluate the storage condition in a study, Malav et al. (2012) assessed the impact of water chestnut flour on the quality and durability of storage. They had positive results and concluded that water chestnut flour enhances the storage time.

Recent Advancements in Restructured Meat Research and Development

The RM market is growing gradually. Presently, dedicated research is being conducted on RM product development. Gurikar et al. (2014) was successful in producing restructured pork blocks by using some processing conditions. In another study, the quality of RM was improved to meet consumer demands (Anna Anandh and Annal Villi, 2018). This study revealed that spent Hen can effectively produce nuggets using salt, STP, and sodium nitrite. Some of the latest techniques are also being used in the field of RM, as Zhu et al. (2019) utilized pressure-transform rolling techniques to make restructured pork chops. Some genes from plant or fungi sources are also used for RM; Yang and Zhang (2019) used a recombinant transglutaminase gene from *Pichia pastoris* (methylotrophic yeast) to restructure the meat. Citric acid (0.2%) was used to restructure fish (*Oreochromis mossambicus*) by Gu et al. (2021) to improve some of the physical qualities of the meat.

Nowadays, researchers are focusing on functional attributes and trying to improve the health properties of RM products. In such an effort, Ahmad et al. (2021) manufactured restructured buffalo meat pieces that were low in sodium, high in fiber, and full of antioxidants. Gorbunova (2022) used the shockwave method to restructure fresh meat and make it more tender. Saengsuk et al. (2022) used alginate/calcium and κ -carrageenan to make restructured pork steak. In contrast, the samples supplemented with κ -carrageenan retained their red color better and can be easily chewed while maintaining the unique taste by adjusting hardness following the USDA guidelines. Lemma et al. (2022) used raisin paste as a natural preservative in jerky dressing to make low-fat restructured jerky products. Extrusion-based 3D food printing was used by Park et al. (2023) to make restructured beef steak. Gupta and Sharma (2023) conducted a study to examine the quality of spent hen meat after adding some binder and extender to meat. They restructured spent hen meat slices by adding soy protein to make the chunks more tender.

Market Scenario

Countries with high incomes also have higher meat demand, which puts significant pressure on livestock meat production (Parlasca and Qaim, 2022). Reducing meat consumption may facilitate the reduction of the environmental impact and minimize the pressure on livestock meat production (Sijpestijn et al., 2022). The increase in meat consumption may aid the growth of RM products. According to Hankyoreh (2023), meat consumption is increasing in Korea by an average of 2.8% per annum, and consumers are becoming health conscious. This could be an opportunity for increased demand for RM products. North America has the biggest market for RM products due to the high population (DI, 2023). Pork is a significant part of RM product development (Saengsuk et al., 2021). Meat scientists are working to improve the quality of RM by improving its color, texture, and taste (Shevchenko et al., 2021). Furthermore, the number of RM customers is also increasing rapidly (DI, 2023). Lifestyle, eating habits, and people's priorities have changed rapidly over the past few decades, which has increased

the demand for meat (Whitton et al., 2021). Advancements in technology (freezing and restructuring technology) and awareness of the benefits of RM can potentially enhance the RM market in the coming years (Ritchie et al., 2021).

According to future market insight (FMI, 2023), efficient marketing is essential for any RM industry's profitability and long-term viability. The demand for RM was predicted to reach 5 billion US\$ by the end of this year, and it is also expected to increase by 18% by 2033 (FMI, 2023). RM in processed food is a new advancement in the meat sector. Thus, meat scientists are assuming that the market value of this type of meat will increase rapidly because RM can be modified according to consumers' demand.

The RM production and selling system requires modernizing. Appropriate processing facilities are necessary to export RM products to the international market. As meat consumption will double in 2050 (FAO, 2014), there is an opportunity to grow the RM market, and that requires potential strategies to meet consumer demand and increase sales by analyzing customer needs, adjusting processing and dispensing methods, developing a solid brand name, implementing rigorous quality control measures, and creating healthier meat products. The majority of this growth is anticipated to occur in developing countries.

The processed meat sector offers significant potential for entrepreneurial development and employment creation, and a gradual increase in RM products (DI, 2023) will open further opportunities. Rigorous quality control measures and producing healthier meat products in response to consumer demand can also be effective for RM product growth.

Conclusion

This review concludes that meat restructuring technology can convert meat trimmings, lesser-value meat cuts, and plant-based protein/fibers into value-added RM products, improving palatability and customer acceptance. Additional studies are required to enhance the processing techniques, product ranges, and functional quality of RM. RM will open opportunities to add different kinds of meat, by-products, plant fibers, and plant proteins to produce HBM, which is healthier for consumers. In addition to its economic impact, the restructuring process can also increase the overall yields and reduce the cost of meat products. The most crucial feature of RM could be its healthy products with reduced fat content, which aligns with the preferences of health-conscious consumers who seek low-fat, low-salt, high-fiber options with minimal synthetic additives. Several reconstructed types of meat, specially HBM with diverse substances, have been created by different existing meat-producing companies, e.g., Tyson Foods, JBS S.A., Cargill, Smithfield Foods, Marfrig Global Foods, Hormel Foods, Maple Leaf Foods, BRF S.A., etc. However, customer acceptance of these items remains challenging due to clarity on the product formation, a specific declaration of health benefits, and the need for a marketing campaign. Categorizing products to particular consumers, ages, and health groups, along with simultaneous engagement in effective marketing campaigns to educate consumers on the RM products, are crucial to increasing the market share and popularization.

Conflicts of Interest

The authors declare no potential conflicts of interest.

Acknowledgements

This research was supported by a National Research Foundation of Korea (NRF) grant funded by the Korean government (MSIT) (No. 2020R1I1A2069379 and 2023R1A2C1004867), and the Korean Institute of Planning and Evaluation for

Technology in Food, Agriculture, Forestry and Fisheries (IPET) through the Agri-Bioindustry Technology Development Program, funded by the Ministry of Agriculture, Food, and Rural Affairs (MAFRA) (Project No. 321028-5), Korea.

Author Contributions

Conceptualization: Samad A, Joo ST. Writing - original draft: Samad A, Alam AMMN. Writing - review & editing: Samad A, Alam AMMN, Kumari S, Hossain MJ, Lee EY, Hwang YH, Joo ST.

Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

References

- Afshar N, Seyyed Qavami SH. 2023. Evaluation of edible coating based on chicken foot gelatin/green walnut skin extract on physical characteristics, color, and texture properties of rainbow trout fillet during storage in the refrigerator. *J Food Sci Technol* 20:1-14.
- Ahmad SR, Sharma BD, Irshad A, Kumar RR, Malav OP, Talukder S. 2021. Effect of aerobic storage conditions on the quality of functional restructured buffalo meat fillets enriched with natural sources of dietary fibers and antioxidant components. *J Food Process Preserv* 45:e15072.
- Alam AMM, Kim CJ, Kim SH, Kumari S, Lee SY, Hwang YH, Joo ST. 2024. Trends in hybrid cultured meat manufacturing technology to improve sensory characteristics. *Food Sci Anim Resour* 44:39-50
- Andrade BF, Guimarães AS, do Carmo LR, Tanaka MS, Fontes PR, Ramos ALS, Ramos EM. 2023. S-nitrosothiols as nitrite alternatives: Effects on residual nitrite, lipid oxidation, volatile profile, and cured color of restructured cooked ham. *Meat Sci* 209:109397.
- Anna Anandh M, Annal Villi R. 2018. Effect of spent hen meat emulsion and ground meat on quality and acceptability of chicken meat cutlets. *Int J Livest Res* 8:33-40.
- Barbut S. 2005. Effect of enhanced fluorescent light on acceptability of meat cuts. *J Muscle Foods* 16:77-86.
- Baugreet S, Kerry JP, Allen P, Gallagher E, Hamill RM. 2018. Physicochemical characteristics of protein-enriched restructured beef steaks with phosphates, transglutaminase, and elasticised package forming. *J Food Qual* 2018:4737602.
- Baune MC, Broucke K, Ebert S, Gibis M, Weiss J, Enneking U, Profeta A, Terjung N, Heinz V. 2023. Meat hybrids: An assessment of sensorial aspects, consumer acceptance, and nutritional properties. *Front Nutr* 10:1101479.
- Bhaskar Reddy GV, Mandal PK, Sen AR, Reddy KS. 2015. Developments in science, technology, quality and constraints of restructured meat products: A review. *Int J Meat Sci* 5:14-48.
- Bhaskar Reddy GV, Sen AR, Nair PN, Sudhakar Reddy K, Kondal Reddy K, Kondaiah N. 2013. Effects of grape seed extract on the oxidative and microbial stability of restructured mutton slices. *Meat Sci* 95:288-294.
- Blaustein-Rejto D, Smith A. 2021. We're on track to set a new record for global meat consumption. Available from: <https://www.technologyreview.com/2021/04/26/1023636/sustainable-meat-livestock-production-climate-change/>. Accessed at Dec 29, 2023.
- Boles JA. 2011. Use of cold-set binders in meat systems. In *Processed meats*. Kerry JP, Kerry JF (ed). Woodhead, Sawston,

- UK. pp 270-298.
- Boles JA, Shand PJ. 2002. Tumbling regime effects on the processing characteristics and tenderness of cooked roast beef. *J Muscle Foods* 13:25-35.
- Carpentieri S, Larrea-Wachtendorff D, Donsi F, Ferrari G. 2022. Functionalization of pasta through the incorporation of bioactive compounds from agri-food by-products: Fundamentals, opportunities, and drawbacks. *Trends Food Sci Technol* 122:49-65.
- Chen G, Cai Y, Su Y, Wang D, Pan X, Zhi X. 2021. Study of meat quality and flavour in different cuts of Duroc-Bamei binary hybrid pigs. *Vet Med Sci* 7:724-734.
- Chen Z, Dai Z, Liu C, Wang S, Li J, Mao X. 2024. Microbial transglutaminase promotes cross-linking for enhancing gelation of myofibrillar protein in frozen *Litopenaeus vannamei* through deamination reaction. *Food Hydrocoll* 147:109332.
- Cifuentes-Galindres DF, Fuenmayor CA, López-Vargas JH. 2024. Restructured beef steaks with inclusion of digestion-resistant maltodextrin: Techno-functional characteristics, nutritional value, sensory quality, and storage stability. *Bioact Carbohydr Diet Fibre* 31:100391.
- Cornet SHV, Snel SJE, Lesschen J, van der Goot AJ, van der Sman RGM. 2021. Enhancing the water holding capacity of model meat analogues through marinade composition. *J Food Eng* 290:110283.
- Data Intelligence [DI]. 2023. Reconstituted meat market by type (pork, beef, lamb/mutton, chicken, turkey, others), by application (food industry, pet food, HoReCa), by region (North America, Latin America, Europe, Asia Pacific, Middle East, and Africa): Share, size, outlook, and opportunity analysis, 2023–2030. Available from: <https://www.datamintelligence.com/research-report/reconstituted-meat-market>. Accessed at Dec 31, 2023.
- Devatkal S, Mendiratta SK. 2001. Use of calcium lactate with salt-phosphate and alginate-calcium gels in restructured pork rolls. *Meat Sci* 58:371-379.
- Dinani ST, Broekema NL, Boom R, van der Goot AJ. 2023. Investigation potential of hydrocolloids in meat analogue preparation. *Food Hydrocoll* 135:108199.
- Dutra MP, Cardoso GP, Ramos EM, Ramos ADLS, Pinheiro ACM, Fontes PR. 2012. Technological and sensory quality of restructured low-fat cooked ham containing liquid whey. *Ciênc Agrotec* 36:86-92.
- Farouk MM. 2010. Restructured whole-tissue meats. In *Handbook of meat processing*. Toldrá F (ed). Blackwell, Hoboken, NJ, USA. pp 399-421.
- Feng Y, Liang X, Zhang J, Kong B, Shi P, Cao C, Zhang H, Liu Q, Zhang Y. 2024. Effects of transglutaminase coupled with κ -carrageenan on the rheological behaviours, gel properties and microstructures of meat batters. *Food Hydrocoll* 146:109265.
- Florowski T, Florowska A, Chmiel M, Dasiewicz K, Adamczak L, Pietrzak D. 2019. The effect of nuts and oilseeds enriching on the quality of restructured beef steaks. *LWT-Food Sci Technol* 104:128-133.
- Food and Agriculture Organization [FAO]. 2014. Meat and meat products. Available from: https://www.fao.org/fileadmin/templates/est/COMM_MARKETS_MONITORING/Meat/Documents/May_2014_FO_MEAT.pdf Accessed at Feb 21, 2024.
- Freire M, Bou R, Cofrades S, Solas MT, Jiménez-Colmenero F. 2016. Double emulsions to improve frankfurter lipid content: Impact of perilla oil and pork backfat. *J Sci Food Agric* 96:900-908.
- Future Market Insight [FMI]. 2023. Reconstituted meat market outlook (2023 to 2033). Available from: <https://www.futuremarketinsights.com/reports/reconstituted-meat-market>. Accessed at Dec 22, 2023.

- Gadekar YP, Sharma BD, Shinde AK, Mendiratta SK. 2014. Effect of different phosphates on quality of goat meat and restructured goat meat product. *Agric Res* 3:370-376.
- Gadekar YP, Sharma BD, Shinde AK, Mendiratta SK. 2015. Restructured meat products - Production, processing and marketing: A review. *Indian J Small Rumin* 21:1-12.
- Gómez I, Janardhanan R, Ibañez FC, Beriain MJ. 2020. The effects of processing and preservation technologies on meat quality: Sensory and nutritional aspects. *Foods* 9:1416.
- Gorbunova NA. 2022. Shockwave effects in the technology of meat raw material processing. *Theory Pract Meat Process* 7:22-29.
- Gu Z, Liu S, Duan Z, Kang R, Zhao M, Xia G, Shen X. 2021. Effect of citric acid on physicochemical properties and protein structure of low-salt restructured tilapia (*Oreochromis mossambicus*) meat products. *J Sci Food Agric* 101:1636-1645.
- Gupta S, Sharma BD. 2023. Effect of texturized soy protein on quality of restructured meat slices from spent hen. *Agric Res* 12:319-324.
- Gurikar AM, Lakshmanan V, Gadekar YP, Sharma BD, Anjaneyulu ASR. 2014. Effect of meat chunk size, massaging time and cooking time on quality of restructured pork blocks. *J Food Sci Technol* 51:1363-1369.
- Hankyoreh. 2023. Meat consumption overtakes rice in Korean diet for first time, report shows. Available from: https://english.hani.co.kr/arti/english_edition/e_national/1078688. Accessed at Jan 2, 2024.
- Hu Y, Zhang L, Yi Y, Solangi I, Zan L, Zhu J. 2021. Effects of sodium hexametaphosphate, sodium tripolyphosphate and sodium pyrophosphate on the ultrastructure of beef myofibrillar proteins investigated with atomic force microscopy. *Food Chem* 338:128146.
- Inguglia ES, Song Z, Kerry JP, O'Sullivan MG, Hamill RM. 2023. Addressing clean label trends in commercial meat processing: Strategies, challenges and insights from consumer perspectives. *Foods* 12:2062.
- Kumar A, Hanjabam MD, Kishore P, Uchoi D, Panda SK, Mohan CO, Chatterjee SN, Zynudheen AA, Ravishankar CN. 2023. Exploitation of seaweed functionality for the development of food products. *Food Bioprocess Technol* 16:1873-1903.
- Kumari S, Alam AN, Hossain MJ, Lee EY, Hwang YH, Joo ST. 2024. Sensory evaluation of plant-based meat: Bridging the gap with animal meat, challenges and future prospects. *Foods* 13:108.
- Kuraishi C, Sakamoto J, Yamazaki K, Susa Y, Kuhara C, Soeda T. 1997. Production of restructured meat using microbial transglutaminase without salt or cooking. *J Food Sci* 62:488-490.
- Lee BJ, Hendricks DG, Cornforth DP. 1998. Effect of sodium phytate, sodium pyrophosphate and sodium tripolyphosphate on physico-chemical characteristics of restructured beef. *Meat Sci* 50:273-283.
- Lemma BB, Lee JH, Kannan G, Kouakou B. 2022. Natural preservative properties of raisins in restructured goat meat (chevon) jerky. *Int J Food Prop* 25:1736-1752.
- Malav OP, Sharma BD, Gokulakrishnan P, Talukder S, Kumar RR. 2012. Effect of water chestnut flour on quality characteristics and storage stability. *Fleischwirtschaft Int* 5:58-62.
- Mandigo RW, Osburn WN. 2019. Cured and processed meats. In *Freezing effects on food quality*. Jeremiah (ed). CRC Press, Boca Raton, FL, USA. pp 135-182.
- Mendiratta SK, Anjaneyulu ASR, Devatkal S, Chauhan G, Lakshmanan V. 2002. Preparation of restructured buffalo meat rolls using calcium phosphate. *J Food Sci Technol* 39:534-536.
- Mireles-Arriaga AI, Ruiz-Nieto JE, Juárez-Abraham MR, Mendoza-Carrillo JM, Martínez-Lopereana R. 2017. Functional

- restructured meat: Applications of ingredients derived from plants. *Vitae* 24:196-204.
- Montone AMI, Malvano F, Taiano R, Capparelli R, Capuano F, Albanese D. 2023. Alginate coating charged by hydroxyapatite complexes with lactoferrin and quercetin enhances the pork meat shelf life. *Foods* 12:553.
- Muller WB. 1991. Effect of method of manufacture. *Fleischwirtschaft* 71:10-18.
- Organisation for Economic Co-operation and Development [OECD], Food and Agriculture Organization [FAO]. 2021. OECD-FAO agricultural outlook 2021-2030. OECD, Paris, France.
- Park JW, Lee SH, Kim HW, Park HJ. 2023. Application of extrusion-based 3D food printing to regulate marbling patterns of restructured beef steak. *Meat Sci* 202:109203.
- Parlasca MC, Qaim M. 2022. Meat consumption and sustainability. *Annu Rev Resour Econ* 14:17-41.
- Patel D, Nayak NK, Chauhan P. 2023. Recent developments in restructured meat products. *Pharma Innov* 12:1124-1129.
- Polásek Z, Salek RN, Vašina M, Lyčková A, Gál R, Pachlová V, Buňka F. 2021. The effect of furcellaran or κ -carrageenan addition on the textural, rheological and mechanical vibration damping properties of restructured chicken breast ham. *LWT-Food Sci Technol* 138:110623.
- Post MJ. 2012. Cultured meat from stem cells: Challenges and prospects. *Meat Sci* 92:297-301.
- Post ML, Levenberg S, Kaplan DL, Genovese N, Fu J, Bryant CJ, Negowetti N, Verzijden K, Moutsatsou P. 2020. Scientific, sustainability and regulatory challenges of cultured meat. *Nat Food* 1:403-415.
- Reverte D, Xiong YL, Moody WG. 2003. Properties of restructured beef steaks from forage- and grain-fed cattle as affected by antioxidant and flavoring agents. *Meat Sci* 65:539-546.
- Ribeiro WO, Ozaki MM, dos Santos M, Rodriguez AP, de Castro RJS, Sato HH, Campagnol PCB, Pollonio MAR. 2023. Improving the textural and nutritional properties in restructured meat loaf by adding fibers and papain designed for elderly. *Food Res Int* 165:112539.
- Ritchie H, Rosado P, Roser M. 2021. Meat and dairy production. Available from: <https://ourworldindata.org/meat-production>. Accessed at Jan 02, 2024.
- Saengsuk N, Laohakunjit N, Sanporkha P, Kaisangsri N, Selamassakul O, Ratanakhanokchai K, Uthairatanakij A. 2021. Physicochemical characteristics and textural parameters of restructured pork steaks hydrolysed with bromelain. *Food Chem* 361:130079.
- Saengsuk N, Laohakunjit N, Sanporkha P, Kaisangsri N, Selamassakul O, Ratanakhanokchai K, Uthairatanakij A, Waeonukul R. 2022. Comparative physicochemical characteristics and *in vitro* protein digestibility of alginate/calcium salt restructured pork steak hydrolyzed with bromelain and addition of various hydrocolloids (low acyl gellan, low methoxy pectin and κ -carrageenan). *Food Chem* 393:133315.
- Schönfeldt HC, Strydom PE. 2011. Effect of age and cut on cooking loss, juiciness and flavour of South African beef. *Meat Sci* 87:180-190.
- Sen AR, Karim SA. 2003. Effect of meat particle size on quality attributes of restructured mutton steaks. *J Food Sci Technol* 40:423-425.
- Serrano A, Cofrades S, Jiménez-Colmenero F. 2006. Characteristics of restructured beef steak with different proportions of walnut during frozen storage. *Meat Sci* 72:108-115.
- Serrano A, Librelotto J, Cofrades S, Sánchez-Muniz FJ, Jiménez-Colmenero F. 2007. Composition and physicochemical characteristics of restructured beef steaks containing walnuts as affected by cooking method. *Meat Sci* 77:304-313.
- Sha L, Xiong YL. 2020. Plant protein-based alternatives of reconstructed meat: Science, technology, and challenges. *Trends*

Food Sci Technol 102:51-61.

- Shafit HM, West RL, Johnson DD, Lindsay JA, Bacus JN. 2007. Effects of sodium lactate and/or potassium lactate on the microbial and sensory quality of beef frankfurter. *J Trop Agric Food Sci* 35:253-259.
- Shao CH, Avens JS, Schmidt GR, Maga JA. 1999. Functional, sensory, and microbiological properties of restructured beef and emu steaks. *J Food Sci* 64:1052-1054.
- Sharma BD, Kumar S, Nanda PK. 2014. Optimization of short term tumbling schedule for the processing of cured and restructured buffalo meat blocks. *Indian J Anim Sci* 72:684-688.
- Sheard P. 2002. Processing and quality control of restructured meat. In *Meat processing: Improving quality*. Kerry J, Kerry J, Ledward D (ed). Woodhead, Sawston, UK. pp 332-358.
- Shevchenko I, Polishchuk G, Topchii O, Kotliar Y, Osmak T. 2021. Improving the technology of restructured ham-type products from turkey meat and pse pork. *Food Sci Technol* 15:106-115.
- Sijpestijn GF, Wezel A, Chriki S. 2022. Can agroecology help in meeting our 2050 protein requirements? *Livest Sci* 256:104822.
- Sikorski ZE. 2004. SMOKING | Traditional. In *Encyclopedia of meat sciences*. 1st ed. Jensen WK, Devine C, Dikeman K (ed). Elsevier, Amsterdam, The Netherlands. pp 1265-1272.
- Silva DRG, Moura APR, Haddad GBS, Souza PM, Ramos ALS, Hopkins DL, Ramos EM. 2021. Reducing the safety risk of low nitrite restructured sliced cooked ham by gamma radiation. *J Food Process Preserv* 45:e15071.
- Sofos JN. 2014. Meat and meat products. In *Food safety management: A practical guide for the food industry*. Academic Press, Cambridge, MA, USA. pp 119-162.
- Talukder S, Sharma BD, Mendiratta SK, Malav OP, Sharma H, Gokulakrishnan P. 2013. Development and evaluation of extended restructured chicken meat block incorporated with colocasia (*Colocasia esculenta*) flour. *J Food Process Technol* 4:207.
- Tsafirakidou P, Sameli N, Kakouri A, Bosnea L, Samelis J. 2023. Assessment of the spoilage microbiota and the growth potential of *Listeria monocytogenes* in minced free-range chicken meat stored at 4 °C in vacuum: Comparison with the spoilage community of resultant retail modified atmosphere packaged products. *Appl Microbiol* 3:1277-1301.
- Wang S. 2023. An investigation of saltiness perception of lysine and calcium lactate and their application in developing reduced salt meat products. Ph.D. dissertation, University of Reading, Reading, UK.
- Wang Y, Domínguez R, Lorenzo JM, Bohrer BM. 2021. The relationship between lipid content in ground beef patties with rate of discoloration and lipid oxidation during simulated retail display. *Foods* 10:1982.
- Wei Z, Dai S, Huang J, Hu X, Ge C, Zhang X, Yang K, Shao P, Sun P, Xiang N. 2023. Soy protein amyloid fibril scaffold for cultivated meat application. *ACS Appl Mater Interfaces* 15:15108-15119.
- Whitton C, Bogueva D, Marinova D, Phillips CJC. 2021. Are we approaching peak meat consumption? Analysis of meat consumption from 2000 to 2019 in 35 countries and its relationship to gross domestic product. *Animals* 11:3466.
- Xia Y, Qian J, Zhao Y, Zheng B, Wei K, Peng B, Yuan J, Xing C, Yan W. 2023. Effects of food components and processing parameters on plant-based meat texture formation and evaluation methods. *J Texture Stud* 54:394-409.
- Xu Y, Xu X. 2021. Modification of myofibrillar protein functional properties prepared by various strategies: A comprehensive review. *Compr Rev Food Sci Food Saf* 20:458-500.
- Yang X, Zhang Y. 2019. Expression of recombinant transglutaminase gene in *Pichia pastoris* and its uses in restructured meat products. *Food Chem* 291:245-252.

- Zhang D, Ge X, Jiao Y, Liu Y. 2024. Quality analysis of steamed beef with black tea and the mechanism of action of main active ingredients of black tea on myofibrillar protein. *Food Chem* 441:137997.
- Zhang M, Fu C, Chen M, Jin C. 2023. The effect of sodium chloride on the physicochemical and textural properties and flavor characteristics of sous vide cooked duck meat. *Foods* 12:3452.
- Zhou Y, Wang Y, Pan Q, Wang X, Li P, Cai K, Chen C. 2020. Effect of salt mixture on flavor of reduced-sodium restructured bacon with ultrasound treatment. *Food Sci Nutr* 8:3857-3871.
- Zhu C, Yin F, Tian W, Zhu Y, Zhao L, Zhao G. 2019. Application of a pressure-transform tumbling assisted curing technique for improving the tenderness of restructured pork chops. *LWT-Food Sci Technol* 111:125-132.
- Zou Y, Yang H, Li PP, Zhang MH, Zhang XX, Xu WM, Wang DY. 2019. Effect of different time of ultrasound treatment on physicochemical, thermal, and antioxidant properties of chicken plasma protein. *Poult Sci* 98:1925-1933.