

**REVIEW**

# Potential Prebiotic Properties of Whey Protein and Glycomacropeptide in Gut Microbiome

 Bryna Rackerby<sup>1</sup>, Hoang Ngoc M. Le<sup>1</sup>, Avery Haymowicz<sup>1</sup>, David C. Dallas<sup>1,2</sup>, and Si Hong Park<sup>1,3,\*</sup>
<sup>1</sup>Department of Food Science and Technology, Oregon State University, Corvallis, OR 97331, USA

<sup>2</sup>School of Biological and Population Health Sciences, Nutrition, Oregon State University, Corvallis, OR 97331, USA

<sup>3</sup>Department of Food Science and Technology, Chung-Ang University, Anseong 17546, Korea

 OPEN ACCESS

**Received** October 18, 2023

**Revised** January 22, 2024

**Accepted** January 29, 2024

**\*Corresponding author** : Si Hong Park  
 Department of Food Science and Technology,  
 Oregon State University, Corvallis, OR  
 97331, USA  
 Tel: +1-541-737-1684  
 Fax: +1-541-737-1877  
 E-mail: [sihong.park@oregonstate.edu](mailto:sihong.park@oregonstate.edu)

**\*ORCID**  
 Bryna Rackerby  
<https://orcid.org/0000-0003-0582-5928>  
 Hoang Ngoc M. Le  
<https://orcid.org/0009-0005-5890-5245>  
 Avery Haymowicz  
<https://orcid.org/0009-0003-5131-7788>  
 David C. Dallas  
<https://orcid.org/0000-0002-9696-0967>  
 Si Hong Park  
<https://orcid.org/0000-0001-6587-7020>


**Abstract** Proteins in whey have prebiotic and antimicrobial properties. Whey protein comprises numerous bioactive proteins and peptides, including glycomacropeptide (GMP), a hydrophilic casein peptide that separates with the whey fraction during cheese making. GMP has traditionally been used as a protein source for individuals with phenylketonuria and also has prebiotic (supporting the growth of *Bifidobacterium* and lactic acid bacteria) and antimicrobial activities. GMP supplementation may help positively modulate the gut microbiome, help treat dysbiosis-related gastrointestinal disorders and improve overall health in consumers.

**Keywords** whey protein, glycomacropeptide (GMP), prebiotic effect, gut microbiome

## Introduction

Whey is a byproduct of the cheese manufacturing process. It is the liquid portion that is drained away after curd formation. Bovine milk is the most common source of whey due to its predominance in the dairy industry. The extracted product referred to as whey protein is a mixture of numerous proteins including  $\beta$ -lactoglobulin (~65%),  $\alpha$ -lactalbumin (~25%), bovine serum albumin (~8%), lactoferrin (~1%), lactoperoxidase (0.25%–0.5%) and immunoglobulins (<1%) (Boscaini et al., 2020).

Bovine whey protein is used in a large array of products, including supplements to promote muscle synthesis and infant formula. Whey protein helps increase muscle synthesis which can help with muscle and exercise recovery in athletes (Moore, 2019; West et al., 2017), and decrease the risk of age-related sarcopenia (Liao et al., 2019; Yang et al., 2012; Fig. 1). Bovine whey protein is added to most infant formulas to mimic human milk. Beyond serving as a source of amino acids, the proteins in whey



**Fig. 1.** Overall relations between whey and GMP with their primary functions.

can exert additional functions, including prebiotic, antimicrobial (Kareb and Aider, 2019), and enhancing gastrointestinal health (Li et al., 2018). For example, many whey proteins, including lactoferrin, lactoperoxidase and immunoglobulins have direct or indirect antimicrobial activity (Bielecka et al., 2022). Perhaps due to these bioactivities, whey protein supplementation can alter the gut microbiome (at least in animal models; Boscaini et al., 2020; Nilaweera et al., 2017) and thereby impact metabolism (Boscaini et al., 2020; Fig. 1).

Glycomacropeptide (GMP) is a hydrophilic peptide cleaved from  $\kappa$ -casein during cheesemaking and makes up about 20% by mass of commercial whey protein (Neelima et al., 2013). About half of the GMP in whey is the unglycosylated form known as caseinomacropeptide, whereas the other 50% consists of 14 different glycovariants (Robitaille, 2012). GMP has several functions, including prebiotic which promotes the the growth of *Bifidobacterium* and lactic acid bacteria (Córdova-Dávalos et al., 2019; Fig. 1), antimicrobial and immunomodulatory (Córdova-Dávalos et al., 2019).

## Impact of Whey on the Gut Microbiome

Whey protein supplementation can impact the gut microbiome. In an infant fecal culture system with 3 feeding conditions; 1) bovine milk, 2) formula with  $\alpha$ -lactalbumin, and 3) formula with GMP, both formulas supplemented with  $\alpha$ -lactalbumin and GMP both supported *Bifidobacterium* as the predominant organism and decreased *Bacteroides*, *Clostridium* and *Escherichia coli* (Brück et al., 2003). In the same study, all groups reduced enteropathogenic *E. coli* and *Salmonella* Typhimurium after initial inoculation of these pathogens into the fecal culture system (Brück et al., 2003).

Whey protein feeding has also been shown to alter the microbiome in animal models. Nilaweera et al. (2017) found that whey protein isolate supplementation reduced the susceptibility of mice to sucrose-induced microbial changes to a larger degree than those supplemented with casein. The study indicated that whey protein reduced the abundance of Firmicutes and Actinobacteria, which is associated with enhanced metabolic health and reduced inflammation, and increased Bacteroidetes, a phylum that can break down complex carbohydrates and produce beneficial short-chain fatty acids (SCFAs). In the same

study, whey protein feeding increased the families *Unclassified\_Sutterellaceae*, *Sutterellaceae*, *Anaeroplasmanaceae*, *Unclassified\_Porphyrimonadaceae* and *Porphyrimonadaceae*, and decreased *Streptococcaceae* and *Enterobacteriaceae*, resulting in the increase of potential healthy gut microbiomes (Nilaweera et al., 2017). Similarly, mice with autoimmune prostatitis fed glycated whey had decreased Firmicutes and an increased *Porphyrimonadaceae*, among other families and genera (Chen et al., 2020). In mice fed a high-fat diet (HFD), whey protein increased *Lactobacillaceae* and *Clostridiaceae* at the family level and *Desulfovibrio* and *Mucisprillum* at the genus level (McAllan et al., 2014). Boscaini et al. (2020) found that 5-week-old mice fed whey (compared with casein) on a HFD had increased *Streptococcaceae* and *Lactococcus* at the family and genus levels, increased *Lactococcus lactis* and *Bacteroides vulgatus* at the species level and increased abundance of *Lactobacillus murinis*. Sprong et al. (2010) found that casein supplemented with either whey protein or threonine and cysteine both increased *Lactobacillus* and *Bifidobacteria*, decreased markers of inflammation and increased mucin secretion in rats with dextran sulfate sodium-induced colitis. Chen et al. (2020) found that non-obese diabetic mice fed whey protein-derived early glycation products for 6 months had increased *Allobaculum*, *Anaerostipes*, *Bacteroides*, *Parabacteroides* and *Prevotella* and decreased *Adlercreutzia* and *Roseburia* at the genus level. In this study, some of the changes in the microbiome correlated with immune markers measured, which suggested that the microbial changes may have contributed to the observed anti-inflammatory effects (Chen et al., 2020). Specifically, total splenocytes were negatively correlated with *Bacteroides (uniformis and acidifaciens)*, *Parabacteroides*, *Prevotella*, and *Anaerostipes*; splenic M1 macrophages were negatively correlated with *Bacteroides (uniformis and acidifaciens)* and *Parabacteroides*; and splenic CD4+ T-cells were negatively correlated with *Bacteroides acidifaciens* (Chen et al., 2020).

Nielsen et al. (2020) found that piglets given whey protein concentrate (WPC) with high  $\alpha$ -lactalbumin tended to have higher alpha-diversity and a higher abundance of *Clostridiaceae*, *Enterobacteriaceae*, *Streptococcus* and *Streptomyces* than those supplemented with WPC with low  $\alpha$ -lactalbumin. However, the microbial composition resulting from dietary interventions did not show evidence of correlation with physiological changes or functional, or performance aspects of preterm pigs (Nielsen et al., 2020).

Though dietary whey protein modulates the gut microbiome in fecal cultures, murine models and piglet models, the few human studies have not been able to replicate these results. This lack of findings in humans may be due to the large degree of variation in human gut microbiomes, behaviors, environmental factors and genetics. Two studies on primarily overweight or obese adult humans found that whey protein did not significantly alter the gut microbiome (Cronin et al., 2018; Reimer et al., 2017). However, one study on endurance athletes fed between whey isolate and beef hydrolysate for 10 weeks indicated an altered gut microbiome with increased *Bacteroidetes* and decreased health-related taxa, suggesting a potential negative impact of long-term protein supplementation that requires further research (Moreno-Pérez et al., 2018). A summary of recent studies related to whey protein diets is listed in Table 1.

## Impact of Glycomacropeptide on the Gut Microbiome

GMP can promote the growth of beneficial organisms and inhibit the adhesion of pathogens to intestinal cells (Córdova-Dávalos et al., 2019). Córdova-Dávalos et al. (2019) provided an extensive review of GMP's ability to prevent the adhesion of pathogens (e.g., *S. Typhimurium* and enterohemorrhagic *E. coli* 0157) and toxins (e.g., cholera toxin, *E. coli* enterotoxin) to intestinal cells. GMP also demonstrates an enhancement in the growth of some specific probiotic organisms. For example, bovine and caprine GMP both improved the growth of *Lactobacillus rhamnosus* RW-9595-M and *Bifidobacterium*

**Table 1. A summary of the recent studies examining the effect of different whey protein diets on gut microbiome diversity**

Area	Criteria of study	Outcome: Changes in gut microbiome (whey only)		Reference
		Increase	Decrease	
Animal studies	Mice fed whey protein isolate (WPI) for 17 weeks	<i>Bacteroidetes</i>	Firmicutes and actinobacteria	Nilaweera et al., 2017
	Non-obese diabetic mice fed glycosylated whey proteins for 6 months	<i>Porphyromonadaceae</i>	Firmicutes	Chen et al., 2020
	Mice on a high-fat diet (HFD) fed for 21 weeks	<i>Lactobacillaceae</i>	<i>Clostridiaceae/Clostridium</i>	McAllan et al., 2014
	Mice on a HFD fed WPI versus casein for 5 weeks	<i>Lactobacillus murinus</i>	<i>HsL</i> and <i>Lpl</i> expression	Boscaini et al., 2020
	Rats with 3% dextran sulfate sodium-induced colitis fed cheese whey protein for 14 days	<i>Lactobacilli</i> and <i>Bifidobacteria</i>	N/A	Sprong et al., 2010
	Preterm piglets fed $\alpha$ -lactalbumin-enriched whey protein concentrate (WPC) versus regular WPC for 19 days	<i>Clostridiaceae</i> , <i>Enterobacteriaceae</i> , and <i>Lachnospiraceae</i>	None noted	Nielsen et al., 2020
	Pre-pubertal male rats under 60 days of food restriction and refeeding with casein- or whey-based diet	<i>Burkholderiales</i> (phylum Proteobacteria), <i>Bacilli</i> (phylum Firmicutes), and <i>Lactibacillaceae</i>	<i>Erysipelotrichales</i> , <i>Cytophagales</i> , and <i>Flavobacteriales</i>	Masarwi et al., 2018
	Mice fed high-fat-whey protein concentrate and high-fat whey-protein hydrolysate for 9 weeks	Bacteroidetes phylum ( <i>B. rodentium</i> , <i>B. acidifaciens</i> and <i>B. stercoris</i> )	<i>Firmicutes</i>	Monteiro et al., 2016
	Rats with and without tumors fed a whey-based diet containing medium-chain triglycerides	<i>Muribaculaceae</i> and <i>Peptostreptococcaeae</i>	<i>Ruminococcaceae</i>	Wardill et al., 2023
Human study	Endurance athletes fed whey isolate or beef hydrolysate for 10 weeks	Bacteroidetes phylum	<i>Citrobacter</i> , <i>Klebsiella</i> , <i>Coprococcus</i> , <i>Roseburia</i> , and <i>Blautia</i> genera	Moreno-Pérez et al., 2018
	With or without whey protein supplement on adult engaged in aerobic and resistance training for 8 weeks	<i>Lactococcus</i> phage, $\beta$ -diversity of gut virome	None noted	Cronin et al., 2018
	Adults with overweight/obesity assigned snack bar with whey protein for 12 weeks	None	None	Reimer et al., 2017
<i>In vitro</i> study	<i>In vitro</i> infant fecal culture exposed to whey protein hydrolysate enriched in essential amino acids	<i>Lactobacillus acidophilus</i> , <i>Proteobacteria</i> , <i>Streptococcus</i> , and <i>Bacteroides</i>	Actinobacteriota:Bacteroidota (ratio)	Feng et al., 2022

N/A, not applicable.

*thermophilum* RBL67 in a dose-dependent manner and to a greater degree than bovine  $\beta$ -lactoglobulin (Robitaille, 2012).

The mechanism by which GMP exhibits prebiotic activity is unclear. Though studies on the antimicrobial impact of GMP tend to implicate the glycosylation structures in its bioactivity (Feeney et al., 2017), studies examining growth-promoting ability are less concordant. Some studies suggest the prebiotic effect stems from the glycan moieties of GMP. For example, periodate oxidation to remove the glycans from GMP significantly reduced its bifidogenic effect, which was interpreted as evidence that its prebiotic activity is linked to glycosylation (O’Riordan et al., 2018). This study also found that GMP induces the expression of some glycogenes, but repressed others, perhaps representing a glycan-structure specific response (O’Riordan et al., 2018).

Further, sialyl glycopeptide concentrate created through digestion and ultrafiltration of GMP-containing WPC (G-WPC) outperformed G-WPC as the sole carbon source for the growth of certain *Bifidobacteria* (Fukudome et al., 2021). Although the glycan component of GMP is often hypothesized to be the basis for prebiotic activity, the peptide portion may be involved as well, as periodate-treated GMP did possess a small growth-promoting effect (O’Riordan et al., 2018). Similarly, Tian et al. (2014) found that GMP’s *Bifidobacteria* growth-promoting effect did not directly depend on sialic acid content and may be related to its high glutamine, leucine, and alanine contents, despite the poor proteolytic activity of most *Bifidobacteria*. GMP hydrolysate produced with papain (GHP) had a stronger growth-promoting effect on *Bifidobacterium animalis* subsp. *lactis* (Bb12) than intact GMP or GMP hydrolyzed by trypsin, although GHP had the lowest sialic acid content of the three (Tian et al., 2014). Robitaille (2012) found that glycosylated, unglycosylated, and mixed GMP treatments equally promoted the growth of lactic acid bacteria, indicating that the glycosylation state is not necessarily a factor in prebiotic activity, even though neither *Bifidobacteria* strain studied was proteolytic. Robitaille (2012) suggested that GMP allows improved growth of *Bifidobacteria* and *Lactobacillus* in acidic media during fermentation by triggering metabolic adaptations. Regardless of the mechanism, GMP is effective in encouraging the growth of probiotic organisms *in vitro* and could be used in the production of probiotics or as a functional ingredient to promote the growth of probiotic cultures in fermented dairy products and influence beneficial organisms in the gut (O’Riordan et al., 2018; Robitaille, 2012; Tian et al., 2014).

Like whey protein, GMP can impact the gut microbiome. In an *in vitro* human fecal culture system, GMP addition supported stable *Bifidobacterium* presence and decreased *Bacteroides*, *Clostridium* and *E. coli* (Brück et al., 2003). Chen et al. (2012) demonstrated that feeding GMP to mice promoted the growth of *Lactobacillus* and *Bifidobacteria* while decreasing *Enterobacteriaceae* and coliforms.

In piglets, compare with control diet, diet supplemented with 1.5% casein GMP resulted in positive changes to the gut microbiome (increased *Lactobacillus* and decreased *Enterobacteria*; Hermes et al., 2012). Supplementation of sows with a combination of galacto-oligosaccharide (GOS) and casein GMP (GOS+GMP) during late gestation through farrowing induced changes to the intestinal microbiome in both the sow and their offspring, when comparing to control treatment based on nutrient requirements from National Research Council (Wu et al., 2020). At the phylum level, Fusobacteria became more prevalent in GOS+GMP-fed sows, whereas their offspring had an increase in Synergistetes and a decrease in Patescibacteria. At the genus level, GOS+GMP-fed sows had higher *Prevotella*, *Fusobacterium*, and *unclassified\_f\_Prevotellaceae*, and their offspring had higher *norank\_f\_Ruminococcaceae*, *Christensenellaceae\_R-7\_group*, *Ruminococcaceae\_UCG-005*, and *Ruminococcaceae\_UCG-010* (Wu et al., 2020). Beyond these changes in the microbiome, this supplementation improved the number of live and healthy piglets, total litter weight, and average birth weight of live piglets (Wu et al., 2020). These findings align with previous work demonstrating that maternal diet impacts the health and microbial composition of the intestinal tract of offspring (Kashtanova et al., 2016). The study design did not allow identification of any effect due to GMP alone.

## Effects of Glycomacropeptide on the Microbiome in Humans

Though GMP has growth-promoting effects on probiotic organisms *in vitro* and in animal models, these results are not consistently replicable in humans. Wernlund et al. (2020) found that GMP supplementation in healthy human subjects had no effects on Shannon or observed diversity and no changes in microbiome composition or fecal SCFAs. Moreover, that study showed no effect of GMP on gastrointestinal symptoms. Likewise, the study found no effect of GMP on high-sensitivity C-reactive protein, fecal calprotectin, indicating little systemic immunomodulatory impacts (Wernlund et al., 2020). A possible

explanation for the observed lack of change in humans is the increased variability in human genetics, environments, behaviors and microbiomes compared to laboratory mice.

A recent crossover study (Hansen et al., 2023) in which obese women consumed twice or thrice daily GMP supplements (15 g GMP+10 g whey protein/dose) found that compared to baselines, the high dose GMP resulted in lower overall alpha-diversity and the low dose resulted in lower relative abundance of the genus *Streptococcus*. The observed changes in the gut microbiome were unlikely to be the cause of the observed increases in satiety and higher area under the curves of the glucoregulatory/satiety hormone amylin and the C-peptide of insulin and lower glucagon in a blood test after a GMP meal tolerance test compared with a baseline soy meal tolerance test. In this study, GMP supplementation did not affect weight, markers of systemic inflammation or plasma SCFAs (Hansen et al., 2023).

## Glycomacropeptide Effects on Dysbiosis

GMP may be able to improve dysbiosis induced by old age or metabolic syndromes. In the elderly, a loss of microbial diversity may be a contributing factor in suboptimal health (Ntemiri et al., 2017). In an artificial colon model of elderly gut microbiota, the addition of GMP increased microbial diversity and increased the growth of the beneficial gut bacteria *Coprococcus* and *Clostridium* cluster XIVb (Ntemiri et al., 2017). In fecal cultures from free-living subjects GMP supplementation increased *Roseburia* and tended to increase *Dorea*, whereas in fecal culture from subjects residing in long-stay facilities *Pseudoflavonifactor* increased (Ntemiri et al., 2017). Type 2 diabetes is also associated with a loss of microbial diversity and changes to the microbial composition (Yuan et al., 2020). Mice with type 2 diabetes had increased *Helicobacteraceae* and *Lachnospiraceae* and decreased *Bacteroidales\_S24-7\_group* (Yuan et al., 2020). Feeding GHP to these diabetic mice recovered lost microbial diversity, reduced *Helicobacteraceae*, and increased *Ruminococcaceae* and *Bacteroidales\_S24-7\_group*, the ratio of Bacteroidetes:Firmicutes, and *Ruminiclostridium*, *Blautia*, and *Allobaculum* (Yuan et al., 2020). Similarly, feeding non-hydrolyzed GMP increased the abundance of *Allobaculum* in wild-type mice and the abundance of *Bacteroidales\_f\_S24-7;g\_* in mice with phenylketonuria and reduced *Desulfovibrio* in both wild-type and phenylketonuria mice, which is associated with inflammatory bowel disease (Sawin et al., 2015).

## Impacts of Glycomacropeptide-Induced Microbial Shifts on Physiology

The changes in the gut microbiome induced by GMP can have physiological impacts (Table 2). For example, in a study by Yuan et al. (2020), supplementing GHP to mice with high fat diet and streptozotocin-induced type 2 diabetes induced antidiabetic effects that correlated with changes in the microbiome. Similarly, microbial changes in piglets whose mothers were supplemented with a combination of GOS and GMP were positively correlated with mRNA transcript levels for claudin-1, claudin-2, occludin, mucin-4, and mucin-13 (Wu et al., 2020).

## Future Perspective

Though many studies have examined the effects of GMP in animal models [e.g., neonatal piglets (Wu et al., 2020), rhesus monkeys (Kelleher et al., 2003), and mice (Nilaweera et al., 2017)], studies in humans are scarce. Moreover, we know little about how changes in the microbiome induced by GMP affect physiology. Future research is needed to examine the effects of whey protein and GMP on the microbiome in humans and determine their clinical impacts. Moreover, more studies

**Table 2. A summary of glycomacropeptide (GMP) effects on gut microbiome diversity**

Area/criteria of study	Specific effect	Reference
General population health	Decrease <i>Streptococcus</i> abundance in the gut	Hansen et al., 2023
	Enhance indicators of satiety and glycemic control	
	Positive relationship of GMP+galacto-oligosaccharide (GOS) feeding with increased mRNA transcript levels for claudin-1, claudin-2, occludin, mucin-4, and mucin-13	Wu et al., 2020
	Changes in the microbiome were positively correlated with antidiabetic effects after feeding GMP hydrolysates	Yuan et al., 2020
	Maternal GMP+GOS supplementation improved litter characteristics in piglet (number of live and healthy piglets, total litter weight, and average birth weight of live piglets)	Wu et al., 2020
	Maternal GMP+GOS supplementation increased immunoglobulins (IgA, IGF-1, IgG, IgM) in neonatal piglets	
Microbiome – Dysbiosis	Increased <i>Lactobacillus</i> , stable <i>Bifidobacterium</i> , decreased <i>Bacteroides</i> , <i>Clostridium</i> , and <i>Escherichia coli</i> in human fecal cultures	Brück et al., 2003
	Improved microbial diversity in an artificial colon model for elderly people: Increased growth of <i>Coprococcus</i> , <i>Clostridium cluster XIVb</i> , <i>Roseburia</i> ; decreased <i>Dorea</i>	Ntemiri et al., 2017
	Reduced <i>Helicobacteraceae</i> ; increased <i>Ruminococcaceae</i> , <i>Bacteroidales_S24-7_group</i> in mice with type-2 diabetes	Yuan et al., 2020

examining the long-term effect of whey protein and GMP supplementation on gut health and immune function are needed (Pena et al., 2018). Further research can guide the use of GMP and whey protein as therapeutics.

## Conclusion

Both whey protein and GMP have a range of bioactivities, including prebiotic action and antimicrobial actions, and may enhance human health beyond provision of their amino acids. In animal models, whey protein supplementation frequently results in increased gut microbiota diversity, increased growth of beneficial microbial species and decreased markers of inflammation. GMP's capacity to promote the growth of beneficial organisms makes it a potential prebiotic dietary supplement. More research is needed to determine the extent to which whey protein and GMP affect the microbiome in humans and the extent to which any such changes affect overall physiology.

## Conflicts of Interest

The authors declare no potential conflicts of interest.

## Acknowledgements

This review manuscript was supported by the BUILD (Building University-Industry linkages through Learning and Discovery) Dairy program and Glanbia.

## Author Contributions

Conceptualization: Rackerby B, Dallas DC, Park SH. Data curation: Le HNM. Validation: Le HNM, Haymowicz A, Park

SH. Investigation: Park SH. Writing - original draft: Rackerby B. Writing - review & editing: Rackerby B, Le HNM, Haymowicz A, Dallas DC, Park SH.

## Ethics Approval

This article does not require IRB/IACUC approval because there are no human and animal participants.

## References

- Bielecka M, Cichosz G, Czczot H. 2022. Antioxidant, antimicrobial and anticarcinogenic activities of bovine milk proteins and their hydrolysates: A review. *Int Dairy J* 127:105208.
- Boscaini S, Cabrera-Rubio R, Nychyk O, Speakman JR, Cryan JF, Cotter PD, Nilaweera KN. 2020. Age- and duration-dependent effects of whey protein on high-fat diet-induced changes in body weight, lipid metabolism, and gut microbiota in mice. *Physiol Rep* 8:e14523.
- Brück WM, Graverholt G, Gibson GR. 2003. A two-stage continuous culture system to study the effect of supplemental  $\alpha$ -lactalbumin and glycomacropeptide on mixed cultures of human gut bacteria challenged with enteropathogenic *Escherichia coli* and *Salmonella* serotype Typhimurium. *J Appl Microbiol* 95:44-53.
- Chen Q, Cao J, Jia Y, Liu X, Yan Y, Pang G. 2012. Modulation of mice fecal microbiota by administration of casein glycomacropeptide. *Microbiol Res* 3:e3.
- Chen Y, Guo KM, Nagy T, Guo TL. 2020. Chronic oral exposure to glycated whey proteins increases survival of aged male NOD mice with autoimmune prostatitis by regulating the gut microbiome and anti-inflammatory responses. *Food Funct* 11:153-162.
- Córdova-Dávalos LE, Jiménez M, Salinas E. 2019. Glycomacropeptide bioactivity and health: A review highlighting action mechanisms and signaling pathways. *Nutrients* 11:598.
- Cronin O, Barton W, Skuse P, Penney NC, Garcia-Perez I, Murphy EF, Woods T, Nugent H, Fanning A, Melgar S, Falvey EC, Holmes E, Cotter PD, O'Sullivan O, Molloy MG, Shanahan F. 2018. A prospective metagenomic and metabolomic analysis of the impact of exercise and/or whey protein supplementation on the gut microbiome of sedentary adults. *mSystems* 3:e00044-18.
- Feeney S, Ryan JT, Kilcoyne M, Joshi L, Hickey R. 2017. Glycomacropeptide reduces intestinal epithelial cell barrier dysfunction and adhesion of entero-hemorrhagic and entero-pathogenic *Escherichia coli* *in vitro*. *Foods* 6:93.
- Feng C, Tian L, Hong H, Wang Q, Zhan X, Luo Y, Tan Y. 2022. *In vitro* gut fermentation of whey protein hydrolysate: An evaluation of its potential modulation on infant gut microbiome. *Nutrients* 14:1374.
- Fukudome H, Yamaguchi T, Higuchi J, Ogawa A, Taguchi Y, Li J, Kabuki T, Ito K, Sakai F. 2021. Large-scale preparation and glycan characterization of sialylglycopeptide from bovine milk glycomacropeptide and its bifidogenic properties. *J Dairy Sci* 104:1433-1444.
- Hansen KE, Murali S, Chaves IZ, Suen G, Ney DM. 2023. Glycomacropeptide impacts amylin-mediated satiety, postprandial markers of glucose homeostasis, and the fecal microbiome in obese postmenopausal women. *J Nutr* 153:1915-1929.
- Hermes RG, Molist F, Francisco Pérez J, de Segura AG, Ywazaki M, Davin R, Nofrarias M, Korhonen TK, Virkola R, Martín-Orúe SM. 2012. Casein glycomacropeptide in the diet may reduce *Escherichia coli* attachment to the intestinal mucosa and increase the intestinal lactobacilli of early weaned piglets after an enterotoxigenic *E. coli* K88 challenge. *Br J*



Nutr 109:1001-1012.

- Kareb O, Aider M. 2019. Whey and its derivatives for probiotics, prebiotics, synbiotics, and functional foods: A critical review. *Probiotics Antimicrob Proteins* 11:348-369.
- Kashtanova DA, Popenko AS, Tkacheva ON, Tyakht AB, Alexeev DG, Boytsov SA. 2016. Association between the gut microbiota and diet: Fetal life, early childhood, and further life. *Nutrition* 32:620-627.
- Kelleher SL, Chatterton D, Nielsen K, Lönnerdal B. 2003. Glycomacropeptide and  $\alpha$ -lactalbumin supplementation of infant formula affects growth and nutritional status in infant rhesus monkeys. *Am J Clin Nutr* 77:1261-1268.
- Li Y, Nguyen DN, Obelitz-Ryom K, Andersen AD, Thymann T, Chatterton DEW, Purup S, Heckmann AB, Bering SB, Sangild PT. 2018. Bioactive whey protein concentrate and lactose stimulate gut function in formula-fed preterm pigs. *J Pediatr Gastroenterol Nutr* 66:128-134.
- Liao Y, Peng Z, Chen L, Zhang Y, Cheng Q, Nüssler AK, Bao W, Liu L, Yang W. 2019. Prospective views for whey protein and/or resistance training against age-related sarcopenia. *Aging Dis* 10:157-173.
- Masarwi M, Solnik HI, Phillip M, Yaron S, Shamir R, Pasmanic-Chor M, Gat-Yablonski G. 2018. Food restriction followed by refeeding with a casein- or whey-based diet differentially affects the gut microbiota of pre-pubertal male rats. *J Nutr Biochem* 51:27-39.
- McAllan L, Skuse P, Cotter PD, O'Connor P, Cryan JF, Ross RP, Fitzgerald G, Roche HM, Nilaweera KN. 2014. Protein quality and the protein to carbohydrate ratio within a high fat diet influences energy balance and the gut microbiota in C57BL/6J mice. *PLOS ONE* 9:e88904.
- Monteiro NES, Roquette AR, de Pace F, Moura CS, Santos A, Yamada AT, Saad MJA, Amaya-Farfan J. 2016. Dietary whey proteins shield murine cecal microbiota from extensive disarray caused by a high-fat diet. *Food Res Int* 85:121-130.
- Moore DR. 2019. Maximizing post-exercise anabolism: The case for relative protein intakes. *Front Nutr* 6:147.
- Moreno-Pérez D, Bressa C, Bailén M, Hamed-Bousdar S, Naclerio F, Carmona M, Pérez M, González-Soltero R, Montalvo-Lominchar MG, Carabaña C, Larrosa M. 2018. Effect of a protein supplement on the gut microbiota of endurance athletes: A randomized, controlled, double-blind pilot study. *Nutrients* 10:337.
- Neelima, Sharma R, Rajput YS, Mann B. 2013. Chemical and functional properties of glycomacropeptide (GMP) and its role in the detection of cheese whey adulteration in milk: A review. *Dairy Sci Technol* 93:21-43.
- Nielsen CH, Hui Y, Nguyen DN, Ahnfeldt AM, Burrin DG, Hartmann B, Heckmann AB, Sangild PT, Thymann T, Bering SB. 2020. Alpha-lactalbumin enriched whey protein concentrate to improve gut, immunity and brain development in preterm pigs. *Nutrients* 12:245.
- Nilaweera KN, Cabrera-Rubio R, Speakman JR, O'Connor PM, McAuliffe A, Guinane CM, Lawton EM, Crispie F, Aguilera M, Stanley M, Boscaini S, Joyce S, Melgar S, Cryan JF, Cotter PD. 2017. Whey protein effects on energy balance link the intestinal mechanisms of energy absorption with adiposity and hypothalamic neuropeptide gene expression. *Am J Physiol Endocrinol Metab* 313:E1-E11.
- Ntemiri A, Chonchúir FN, O'Callaghan TF, Stanton C, Ross RP, O'Toole PW. 2017. Glycomacropeptide sustains microbiota diversity and promotes specific taxa in an artificial colon model of elderly gut microbiota. *J Agric Food Chem* 65:1836-1846.
- O'Riordan N, O'Callaghan J, Buttò LF, Kilcoyne M, Joshi L, Hickey RM. 2018. Bovine glycomacropeptide promotes the growth of *Bifidobacterium longum* ssp. *infantis* and modulates its gene expression. *J Dairy Sci* 101:6730-6741.
- Pena MJ, Pinto A, Daly A, MacDonald A, Azevedo L, Rocha JC, Borges N. 2018. The use of glycomacropeptide in patients

- with phenylketonuria: A systematic review and meta-analysis. *Nutrients* 10:1794.
- Reimer RA, Willis HJ, Tunnicliffe JM, Park H, Madsen KL, Soto-Vaca A. 2017. Inulin-type fructans and whey protein both modulate appetite but only fructans alter gut microbiota in adults with overweight/obesity: A randomized controlled trial. *Mol Nutr Food Res* 61:1700484.
- Robitaille G. 2012. Growth-promoting effects of caseinomacropeptide from cow and goat milk on probiotics. *J Dairy Res* 80:58-63.
- Sawin EA, Wolfe TJD, Aktas B, Stroup BM, Murali SG, Steele JL, Ney DM. 2015. Glycomacropeptide is a prebiotic that reduces *Desulfovibrio* bacteria, increases cecal short-chain fatty acids, and is anti-inflammatory in mice. *Am J Physiol Gastrointest Liver Physiol* 309:G590-G601.
- Sprong RC, Schonewille AJ, van der Meer R. 2010. Dietary cheese whey protein protects rats against mild dextran sulfate sodium-induced colitis: Role of mucin and microbiota. *J Dairy Sci* 93:1364-1371.
- Tian Q, Wang TT, Tang X, Han MZ, Leng XJ, Mao XY. 2014. Developing a potential prebiotic of yogurt: Growth of *Bifidobacterium* and yogurt cultures with addition of glycomacropeptide hydrolysate. *Int J Food Sci Technol* 50:120-127.
- Wardill HR, Ferreira ARDS, Kumar H, Bateman EH, Cross CB, Bowen JM, Havinga R, Harmsen HJM, Knol J, Dorresteyn B, van Dijk M, van Bergenhenegouwen J, Tissing WJE. 2023. Whey-based diet containing medium chain triglycerides modulates the gut microbiota and protects the intestinal mucosa from chemotherapy while maintaining therapy efficacy. *Cell Death Dis* 14:338.
- Wernlund PG, Hvas CL, Dahlerup JF, Bahl MI, Licht TR, Knudsen KEB, Agnholt JS. 2020. Casein glycomacropeptide is well tolerated in healthy adults and changes neither high-sensitive C-reactive protein, gut microbiota nor faecal butyrate: A restricted randomised trial. *Br J Nutr* 125:1374-1385.
- West DWD, Sawan SA, Mazzulla M, Williamson E, Moore DR. 2017. Whey protein supplementation enhances whole body protein metabolism and performance recovery after resistance exercise: A double-blind crossover study. *Nutrients* 9:735.
- Wu Y, Zhang X, Tao S, Pi Y, Han D, Ye H, Feng C, Zhao J, Chen L, Wang J. 2020. Maternal supplementation with combined galactooligosaccharides and casein glycomacropeptides modulated microbial colonization and intestinal development of neonatal piglets. *J Funct Foods* 74:104170.
- Yang Y, Breen L, Burd NA, Hector AJ, Churchward-Venne TA, Josse AR, Tarnopolsky MA, Phillips SM. 2012. Resistance exercise enhances myofibrillar protein synthesis with graded intakes of whey protein in older men. *Br J Nutr* 108:1780-1788.
- Yuan Q, Zhan B, Chang R, Du M, Mao X. 2020. Antidiabetic effect of casein glycomacropeptide hydrolysates on high-fat diet and STZ-induced diabetic mice via regulating insulin signaling in skeletal muscle and modulating gut microbiota. *Nutrients* 12:220.