



## A Systematic Review of Kinesiology Taping in Patients With Lymphedema

Jaehee Yang, RN, CLT, MSc, Eun Jin Lim, MPH, PhD

Department of Integrative Medicine, The Graduate School of Integrative Medicine, CHA University, Pochun, Korea

<https://doi.org/10.12674/ptk.2023.30.4.288>

Phys Ther Korea 2023;30(4):288-305

The authors wish to notify readers of the following change: In page 298 - sentence in the “1) Changes in volume - (1) KT vs. control group (MLB & CG)”:

### <Before correction>

#### 1) Changes in volume

##### (1) KT vs. control group (MLB & CG)

The effect size was statistically significant ( $p < 0.05$ ), with an effect size mean of  $-7.18$  (95% confidence interval [CI] = 12.64 to  $-1.72$ ) across the six literatures, but there was heterogeneity between the literature ( $I^2 = 90\%$ ).

### <After correction>

#### 1) Changes in volume

##### (1) KT vs. control group (MLB & CG)

The effect size was statistically significant ( $p < 0.05$ ), with an effect size mean of  $-7.18$  (95% confidence interval [CI] =  $-12.64$  to  $-1.72$ ) across the six literatures, but there was heterogeneity between the literature ( $I^2 = 90\%$ ).

We apologize for any inconvenience our mistake may have caused.

