

A NEW EXTENSION OF THE MITTAG-LEFFLER FUNCTION

MUHAMMAD ARSHAD, JUNESANG CHOI, SHAHID MUBEEN,
KOTTAKKARAN SOOPPY NISAR, AND GAUHAR RAHMAN

ABSTRACT. Since Mittag-Leffler introduced the so-called Mittag-Leffler function in 1903, due to its usefulness and diverse applications, a variety and large number of its extensions (and generalizations) and variants have been presented and investigated. In this sequel, we aim to introduce a new extension of the Mittag-Leffler function by using a known extended beta function. Then we investigate certain useful properties and formulas associated with the extended Mittag-Leffler function such as integral representation, Mellin transform, recurrence relation, and derivative formulas. We also introduce an extended Riemann-Liouville fractional derivative to present a fractional derivative formula for a known extended Mittag-Leffler function, the result of which is expressed in terms of the new extended Mittag-Leffler functions.

1. Introduction and preliminaries

Gösta Mittag-Leffler (1846-1927) [13] presented an entire function

$$(1) \quad E_{\rho}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\rho n + 1)} \quad (z \in \mathbb{C}; \Re(\rho) > 0),$$

which is called Mittag-Leffler function. Wiman [28, 29] gave and investigated a generalization of the Mittag-Leffler function (1)

$$(2) \quad E_{\rho, \sigma}(z) = \sum_{n=0}^{\infty} \frac{z^n}{\Gamma(\rho n + \sigma)} \quad (z, \sigma \in \mathbb{C}; \Re(\rho) > 0).$$

Here and in the following, let \mathbb{C} , \mathbb{R}^+ , \mathbb{N} , and \mathbb{Z}_0^- be the sets of complex numbers, positive real numbers, positive integers, and non-positive integers, respectively, and let $\mathbb{R}_0^+ := \mathbb{R}^+ \cup \{0\}$. Since then, a variety of generalizations and

Received May 22, 2017; Accepted January 10, 2018.

2010 *Mathematics Subject Classification.* 33B20, 33C20, 33C45, 33C60, 33B15, 33C05, 26A33.

Key words and phrases. gamma function, beta function, extended beta function, Mittag-Leffler function, extended Mittag-Leffler functions, Fox-Wright function, generalized hypergeometric function, Mellin transform, Riemann-Liouville fractional derivative, extended Riemann-Liouville fractional derivative.

extensions of the Mittag-Leffler function (1) have been investigated (see, e.g., [2, 3, 5, 8, 9, 11, 12, 15–17, 19, 20, 27] and the references cited therein). The Mittag-Leffler function and its diverse generalizations are important, in particular, in connection with the theories of fractional calculus and special functions (see, e.g., [5], [21, 22, 24, 25]). Certain interesting applications of the Mittag-Leffler function and its generalizations have been carried out in a wide range of research subjects including physics and engineering such as kinetic equations, Lorenz system, random walk, Levy flights, complex system, fluid flow, electric network, probability, and statistical distribution theory.

Prabhakar [16] introduced and investigated the following generalization of the function $E_{\rho,\sigma}(z)$ (2)

$$(3) \quad E_{\rho,\sigma}^{\gamma}(z) = \sum_{n=0}^{\infty} \frac{(\gamma)_n}{\Gamma(\rho n + \sigma)} \frac{z^n}{n!} \quad (z, \sigma, \gamma \in \mathbb{C}; \Re(\rho) > 0),$$

where $(\lambda)_n$ is the Pochhammer symbol defined (for $\lambda \in \mathbb{C}$) by (see, e.g., [23, p. 2 and pp. 4–6]):

$$(4) \quad (\lambda)_n := \begin{cases} 1 & (n = 0) \\ \lambda(\lambda + 1) \cdots (\lambda + n - 1) & (n \in \mathbb{N}) \end{cases} \\ = \frac{\Gamma(\lambda + n)}{\Gamma(\lambda)} \quad (\lambda \in \mathbb{C} \setminus \mathbb{Z}_0^-),$$

and Γ being the familiar gamma function (see, e.g., [23, Section 1.1]).

Shukla and Prajapati [20] (see also [27]) introduced a generalized Mittag-Leffler function $E_{\rho,\sigma}^{\delta,q}(z)$ defined by

$$(5) \quad E_{\rho,\sigma}^{\delta,q}(z) = \sum_{n=0}^{\infty} \frac{(\delta)_{nq}}{\Gamma(\rho n + \sigma)} \frac{z^n}{n!}$$

$$(z, \rho, \sigma, \delta \in \mathbb{C}; \min\{\Re(\rho), \Re(\sigma), \Re(\delta)\} > 0; q \in (0, 1) \cup \mathbb{N})$$

and investigated to present many interesting properties and formulas involving the generalized Mittag-Leffler function such as Laplace transforms and Whittaker transforms, in a systematic way (see also [6, 7, 10, 18]).

Özarslan and Yilmaz [15] introduced to investigate the following extended Mittag-Leffler function $E_{\rho,\sigma}^{\delta;c}(z; p)$ (see also [14]):

$$(6) \quad E_{\rho,\sigma}^{\delta;c}(z; p) = \sum_{n=0}^{\infty} \frac{B_p(\delta + n, c - \delta)}{B(\delta, c - \delta)} \frac{(c)_n}{\Gamma(\rho n + \sigma)} \frac{z^n}{n!} \\ (p \in \mathbb{R}_0^+; \Re(c) > \Re(\delta) > 0; \Re(\rho) > 0),$$

where $B_p(x, y)$ is extended beta function defined by (see, e.g., [1, 12])

$$(7) \quad B_p(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} e^{-\frac{p}{t(1-t)}} dt \\ (\min\{\Re(p), \Re(x), \Re(y)\} > 0)$$

and $B_0(x, y) = B(x, y)$ is the familiar beta function (see, [23, Section 1.1]).

By giving a generalization of the extended beta function in (7)

$$(8) \quad B_p^{\lambda, \rho}(x, y) = \int_0^1 t^{x-1} (1-t)^{y-1} {}_1F_1 \left[\lambda; \rho; -\frac{p}{t(1-t)} \right] dt$$

$$(p \in \mathbb{R}_0^+; \min\{\Re(x), \Re(y)\} > 0),$$

we define a further extension of the extended Mittag-Leffler function in (6)

$$(9) \quad E_{\alpha, \beta}^{\gamma, c; \lambda, \rho}(z; p) = \sum_{n=0}^{\infty} \frac{B_p^{\lambda, \rho}(\gamma + n, c - \gamma)}{B(\gamma, c - \gamma)} \frac{(c)_n}{\Gamma(\alpha n + \beta)} \frac{z^n}{n!}$$

$$(p \in \mathbb{R}_0^+; \Re(c) > \Re(\gamma) > 0; \Re(\alpha) > 0; \rho \in \mathbb{C} \setminus \mathbb{Z}_0^-).$$

Then we investigate certain interesting properties and formulas involving the extended Mittag-Leffler function (9) such as integral representations, Mellin transform, and recurrence relation. We also introduce an extended Riemann-Liouville fractional derivative to present a fractional derivative formula (40) for the extended Mittag-Leffler function (3), the result of which is expressed in terms of the new extended Mittag-Leffler functions (9). The results presented here are sure to new and potentially useful.

It is remarked in passing that the special cases of (9) when $\rho = \lambda$ and $p = 0$ reduce to the extended Mittag-Leffler functions (6) and (3), respectively.

For our purpose, we recall the Fox-Wright function ${}_p\Psi_q$ (see [4] and [30–32]) (see also [26, p. 21])

$$(10) \quad {}_p\Psi_q \left[\begin{matrix} (\alpha_1, A_1), \dots, (\alpha_p, A_p); \\ (\beta_1, B_1), \dots, (\beta_q, B_q); \end{matrix} z \right] = \sum_{k=0}^{\infty} \frac{\prod_{j=1}^p \Gamma(\alpha_j + A_j k)}{\prod_{j=1}^q \Gamma(\beta_j + B_j k)} \frac{z^k}{k!},$$

where the coefficients $A_j \in \mathbb{R}^+$ ($j = 1, \dots, p$) and $B_j \in \mathbb{R}^+$ ($j = 1, \dots, q$) such that

$$(11) \quad 1 + \sum_{j=1}^q B_j - \sum_{j=1}^p A_j \geq 0.$$

A special case of (10) reduces to the generalized hypergeometric function ${}_pF_q$

$$(12) \quad {}_p\Psi_q \left[\begin{matrix} (\alpha_1, 1), \dots, (\alpha_p, 1); \\ (\beta_1, 1), \dots, (\beta_q, 1); \end{matrix} z \right] = \frac{\prod_{j=1}^p \Gamma(\alpha_j)}{\prod_{j=1}^q \Gamma(\beta_j)} {}_pF_q \left[\begin{matrix} \alpha_1, \dots, \alpha_p; \\ \beta_1, \dots, \beta_q; \end{matrix} z \right].$$

We also recall the generalized gamma function $\Gamma_p^{(\alpha, \beta)}$ (see [14, Eq. (3)])

$$(13) \quad \Gamma_p^{(\alpha, \beta)}(x) = \int_0^{\infty} t^{x-1} {}_1F_1 \left(\alpha; \beta; -t - \frac{p}{t} \right) dt$$

$$(\min\{\Re(\alpha), \Re(\beta), \Re(x)\} > 0; p \in \mathbb{R}_0^+)$$

and (see [14, Eq. (5)])

$$(14) \quad \Gamma_0^{(\alpha, \beta)}(x) =: \Gamma^{(\alpha, \beta)}(x) = \frac{\Gamma(\beta) \Gamma(\alpha - x) \Gamma(x)}{\Gamma(\alpha) \Gamma(\beta - x)}.$$

The special case of (13) when $\alpha = \beta$ and $p = 0$ is seen to yield the familiar gamma function Γ .

2. Properties of the extended Mittag-Leffler function (9)

Here, we give an integral representation and the Mellin transform of the extended Mittag-Leffler function (9).

Theorem 2.1. *Let $c, \alpha, \beta, \gamma, \lambda \in \mathbb{C}$ with $\Re(c) > \Re(\gamma) > 0$ and $\Re(\alpha) > 0$. Also, let $p \in \mathbb{R}_0^+$ and $\rho \in \mathbb{C} \setminus \mathbb{Z}_0^-$. Then*

$$(15) \quad E_{\alpha, \beta}^{\gamma, c; \lambda, \rho}(z; p) = \frac{1}{B(\gamma, c - \gamma)} \times \int_0^1 t^{\gamma-1} (1-t)^{c-\gamma-1} {}_1F_1\left[\lambda; \rho; -\frac{p}{t(1-t)}\right] E_{\alpha, \beta}^c(tz) dt.$$

Proof. Using (8) in (9), we have

$$E_{\alpha, \beta}^{\gamma, c; \lambda, \rho}(z; p) = \sum_{n=0}^{\infty} \left\{ \int_0^1 t^{\gamma+n-1} (1-t)^{c-\gamma-1} \times {}_1F_1\left[\lambda; \rho; -\frac{p}{t(1-t)}\right] dt \frac{(c)_n}{B(\gamma, c - \gamma)} \frac{z^n}{\Gamma(\alpha n + \beta) n!} \right\}.$$

Interchanging the order of summation and integration in the above equation, which is verified under the given conditions here, we get

$$E_{\alpha, \beta}^{\gamma, c; \lambda, \rho}(z; p) = \int_0^1 t^{\gamma-1} (1-t)^{c-\gamma-1} \times {}_1F_1\left[\lambda; \rho; -\frac{p}{t(1-t)}\right] \sum_{n=0}^{\infty} \frac{(c)_n}{B(\gamma, c - \gamma)} \frac{(tz)^n}{\Gamma(\alpha n + \beta) n!} dt.$$

Using (3) in the above equation, we get the desired result. \square

Setting $t = \frac{u}{1+u}$ and $t = \sin^2 \theta$ in the result in Theorem 2.1, we get two interesting integral formulas for the extended Mittag-Leffler function (9), which are given in Corollaries 2.2 and 2.3, respectively.

Corollary 2.2. *Let $c, \alpha, \beta, \gamma, \lambda \in \mathbb{C}$ with $\Re(c) > \Re(\gamma) > 0$ and $\Re(\alpha) > 0$. Also, let $p \in \mathbb{R}_0^+$ and $\rho \in \mathbb{C} \setminus \mathbb{Z}_0^-$. Then*

$$(16) \quad E_{\alpha, \beta}^{\gamma, c; \lambda, \rho}(z; p) = \frac{1}{B(\gamma, c - \gamma)} \int_0^{\infty} \frac{u^{\gamma-1}}{(u+1)^c} \times {}_1F_1\left[\lambda; \rho; -\frac{p(1+u)^2}{u}\right] E_{\alpha, \beta}^c\left(\frac{uz}{1+u}\right) du.$$

Corollary 2.3. *Let $c, \alpha, \beta, \gamma, \lambda \in \mathbb{C}$ with $\Re(c) > \Re(\gamma) > 0$ and $\Re(\alpha) > 0$. Also, let $p \in \mathbb{R}_0^+$ and $\rho \in \mathbb{C} \setminus \mathbb{Z}_0^-$. Then*

$$(17) \quad E_{\alpha,\beta}^{\gamma,c;\lambda,\rho}(z;p) = \frac{2}{B(\gamma,c-\gamma)} \int_0^{\frac{\pi}{2}} \sin^{2\gamma-1} \theta \cos^{2c-1} \theta \times {}_1F_1\left[\lambda; \rho; -\frac{p}{\sin^2 \theta \cos^2 \theta}\right] E_{\alpha,\beta}^c(z \sin^2 \theta) d\theta.$$

Kurulay and Bayram [11] presented the following recurrence relation for the extended Mittag-Leffler function (3)

$$(18) \quad E_{\alpha,\beta}^c(tz) = \beta E_{\alpha,\beta+1}^c(tz) + \alpha z \frac{d}{dz} E_{\alpha,\beta+1}^c(tz).$$

We also present a recurrence relation for the extended Mittag-Leffler function (9) asserted by the following corollary.

Corollary 2.4. *Let $c, \alpha, \beta, \gamma, \lambda \in \mathbb{C}$ with $\Re(c) > \Re(\gamma) > 0$ and $\Re(\alpha) > 0$. Also, let $p \in \mathbb{R}_0^+$ and $\rho \in \mathbb{C} \setminus \mathbb{Z}_0^-$. Then*

$$(19) \quad E_{\alpha,\beta}^{\gamma,c;\lambda,\rho}(z;p) = \beta E_{\alpha,\beta+1}^{\gamma,c;\lambda,\rho}(z;p) + \alpha \gamma z E_{\alpha,\alpha+\beta+1}^{\gamma+1,c+1;\lambda,\rho}(z;p).$$

Proof. Applying (18) to the integrand in (15), we obtain

$$(20) \quad E_{\alpha,\beta}^{\gamma,c;\lambda,\rho}(z;p) = \beta E_{\alpha,\beta+1}^{\gamma,c;\lambda,\rho}(z;p) + \alpha z \frac{d}{dz} E_{\alpha,\beta+1}^{\gamma,c;\lambda,\rho}(z;p).$$

Then we use (37) with $n = 1$ in the derivative in (20) to yield (19). □

Theorem 2.5. *Let $\alpha \in \mathbb{R}^+, p \in \mathbb{R}_0^+, \Re(c) > \Re(\gamma) > 0, \min\{\Re(\beta), \Re(\lambda), \Re(\rho)\} > 0$. Then Mellin transform of the extended Mittag-Leffler function (9) is given by*

$$(21) \quad \mathfrak{M}\left\{E_{\alpha,\beta}^{\gamma,c;\lambda,\rho}(z;p); s\right\} = \frac{\Gamma(\lambda;\rho)(s) \Gamma(c+s-\gamma)}{\Gamma(\gamma)\Gamma(c-\gamma)} {}_2\Psi_2\left[\begin{matrix} (c, 1), (\gamma+s, 1) \\ (\beta, \alpha), (c+2s, 1) \end{matrix}; z\right].$$

Proof. Taking Mellin transform of the extended Mittag-Leffler function (9), we have

$$(22) \quad \mathfrak{M}\left\{E_{\alpha,\beta}^{\gamma,c;\lambda,\rho}(z;p); s\right\} = \int_0^\infty p^{s-1} E_{\alpha,\beta}^{\gamma,c;\lambda,\rho}(z;p) dp.$$

Applying (15) to (22), we have

$$(23) \quad \mathfrak{M}\left\{E_{\alpha,\beta}^{\gamma,c;\lambda,\rho}(z;p); s\right\} = \frac{1}{B(\gamma,c-\gamma)} \times \int_0^\infty p^{s-1} \left\{ \int_0^1 t^{\gamma-1} (1-t)^{c-\gamma-1} {}_1F_1\left[\lambda; \rho; -\frac{p}{t(1-t)}\right] \right\} E_{\alpha,\beta}^c(tz) dt dp.$$

Interchanging the order of integrations in (23), which is guaranteed under the conditions here, we have

$$(24) \quad \mathfrak{M}\left\{E_{\alpha,\beta}^{\gamma,c;\lambda,\rho}(z;p);s\right\} = \frac{1}{B(\gamma,c-\gamma)} \int_0^1 \left[t^{\gamma-1}(1-t)^{c-\gamma-1} E_{\alpha,\beta}^c(tz) \right] \int_0^\infty p^{s-1} {}_1F_1\left[\lambda;\rho;-\frac{p}{t(1-t)}\right] dp dt.$$

Substituting $u = \frac{p}{t(1-t)}$ in the inner integral in (24), we get

$$(25) \quad \begin{aligned} \int_0^\infty p^{s-1} {}_1F_1\left[\lambda;\rho;-\frac{p}{t(1-t)}\right] dp &= \int_0^\infty u^{s-1} t^s (1-t)^s {}_1F_1[\lambda;\rho;-u] du \\ &= t^s (1-t)^s \int_0^\infty u^{s-1} {}_1F_1[\lambda;\rho;-u] du \\ &= t^s (1-t)^s \Gamma^{(\lambda,\rho)}(s), \end{aligned}$$

where $\Gamma_p^{(\lambda,\rho)}(s)$ is the extended gamma function given in (13). Using (25) and (24), we find

$$\begin{aligned} \mathfrak{M}\left\{E_{\alpha,\beta}^{\gamma,c;\lambda,\rho}(z;p);s\right\} &= \frac{\Gamma^{(\lambda,\rho)}(s)}{B(\gamma,c-\gamma)} \int_0^1 t^{\gamma+s-1} (1-t)^{c+s-\gamma-1} \\ &\quad \times \sum_{n=0}^\infty \frac{(c)_n z^n}{\Gamma(\alpha n + \beta) n!} \int_0^1 t^{\gamma+n+s-1} (1-t)^{c+s-\gamma-1} dt \\ &= \frac{\Gamma^{(\lambda,\rho)}(s)}{B(\gamma,c-\gamma)} \sum_{n=0}^\infty \frac{(c)_n z^n}{\Gamma(\alpha n + \beta) n!} \frac{\Gamma(\gamma+n+s)\Gamma(c+s-\gamma)}{\Gamma(c+n+2s)} \\ &= \frac{\Gamma^{(\lambda,\rho)}(s)\Gamma(c+s-\gamma)}{\Gamma(\gamma)\Gamma(c-\gamma)} \sum_{n=0}^\infty \frac{\Gamma(c+n)}{\Gamma(\alpha n + \beta)} \frac{\Gamma(\gamma+n+s)}{\Gamma(c+n+2s)} \frac{z^n}{n!}, \end{aligned}$$

which, in terms of (10), leads to the right side of (21). \square

Setting $s = 1$ in the result in Theorem 2.5 and using (14), we obtain an integral formula involving the extended Mittag-Leffler function (9), which is asserted by the following corollary.

Corollary 2.6. *Let $\alpha \in \mathbb{R}^+$, $p \in \mathbb{R}_0^+$, $\Re(c) > \Re(\gamma) > 0$, $\Re(\beta) > 0$, $\min\{\Re(\lambda), \Re(\rho)\} > 1$. Then*

$$(26) \quad \begin{aligned} &\int_0^\infty E_{\alpha,\beta}^{\gamma,c;\lambda,\rho}(z;p) dp \\ &= \frac{(\rho-1)\Gamma(c+1-\gamma)}{(\lambda-1)\Gamma(\gamma)\Gamma(c-\gamma)} {}_2\Psi_2 \left[\begin{matrix} (c, 1), (\gamma+1, 1); \\ (\beta, \alpha), (c+2, 1); \end{matrix} z \right]. \end{aligned}$$

Setting $\lambda = \rho$ in Corollary 2.6, we get an integral formula involving the extended Mittag-Leffler function (6), which is asserted by the following corollary.

Corollary 2.7. *Let $\alpha \in \mathbb{R}^+$, $p \in \mathbb{R}_0^+$, $\Re(c) > \Re(\gamma) > 0$, and $\Re(\beta) > 0$. Then*

$$(27) \quad \int_0^\infty E_{\alpha,\beta}^{\gamma,c}(z;p) dp = \frac{c-\gamma}{\Gamma(\gamma)} {}_2\Psi_2 \left[\begin{matrix} (c,1), (\gamma+1,1); \\ (\beta,\alpha), (c+2,1); \end{matrix} z \right].$$

3. Derivative formulas of the extended Mittag-Leffler function (9)

Here, we define a further extension of a known extended Riemann-Liouville fractional derivative and present a derivative formula for the extended Mittag-Leffler function (3) involving this extended fractional derivative, which is expressed in terms of the extended Mittag-Leffler functions (9). To do this, we recall the well-known Riemann-Liouville fractional derivative of order $\mu \in \mathbb{C}$ ($\Re(\mu) \geq 0$) defined by

$$(28) \quad \mathfrak{D}_x^\mu \{f(x)\} := \frac{1}{\Gamma(m-\mu)} \frac{d^m}{dx^m} \int_0^x f(t)(x-t)^{-\mu+m-1} dt$$

$$(m \in \mathbb{N}; m-1 < \Re(\mu) < m; x > 0).$$

In particular,

$$(29) \quad \mathfrak{D}_x^\mu \{f(x)\} = \frac{1}{\Gamma(1-\mu)} \frac{d}{dx} \int_0^x f(t)(x-t)^{-\mu} dt$$

$$(0 < \Re(\mu) < 1; x > 0).$$

Özarslan and Yilmaz [15] extended the Riemann-Liouville fractional derivative (28) as follows:

$$(30) \quad \mathfrak{D}_x^{\mu,p} \{f(x)\}$$

$$:= \frac{1}{\Gamma(m-\mu)} \frac{d^m}{dx^m} \int_0^x f(t)(x-t)^{-\mu+m-1} \exp\left(-\frac{px^2}{t(x-t)}\right) dt$$

$$(p \in \mathbb{R}_0^+; m \in \mathbb{N}; m-1 < \Re(\mu) < m; x > 0).$$

In particular,

$$(31) \quad \mathfrak{D}_x^{\mu,p} \{f(x)\} := \frac{1}{\Gamma(1-\mu)} \frac{d}{dx} \int_0^x f(t)(x-t)^{-\mu} \exp\left(-\frac{px^2}{t(x-t)}\right) dt$$

$$(p \in \mathbb{R}_0^+; 0 < \Re(\mu) < 1; x > 0).$$

Here, we define a further extension of the extended Riemann-Liouville fractional derivative (30) by

$$(32) \quad \mathfrak{D}_{x;\lambda,\rho}^{\mu,p} \{f(x)\} := \frac{1}{\Gamma(m-\mu)} \frac{d^m}{dx^m} \int_0^x f(t)(x-t)^{-\mu+m-1} {}_1F_1 \left[\lambda; \rho; -\frac{px^2}{t(x-t)} \right] dt$$

$$(p \in \mathbb{R}_0^+; m \in \mathbb{N}; m-1 < \Re(\mu) < m; x \in \mathbb{R}^+; \lambda \in \mathbb{C}, \rho \in \mathbb{C} \setminus \mathbb{Z}_0^-).$$

In particular,

$$(33) \quad \mathfrak{D}_{x;\lambda,\rho}^{\mu,p} \{f(x)\} := \frac{1}{\Gamma(1-\mu)} \frac{d}{dx} \int_0^x f(t)(x-t)^{-\mu} {}_1F_1 \left[\lambda; \rho; -\frac{px^2}{t(x-t)} \right] dt$$

$$(p \in \mathbb{R}_0^+; 0 < \Re(\mu) < 1; x \in \mathbb{R}^+; \lambda \in \mathbb{C}, \rho \in \mathbb{C} \setminus \mathbb{Z}_0^-).$$

It is obvious that the special cases of (32) when $\lambda = \rho$ and $p = 0$ reduce to the extended Riemann-Liouville fractional derivative (30) and the Riemann-Liouville fractional derivative (28), respectively.

We provide some formulas for the $B_p^{\lambda, \rho}(x, y)$ function in (8) and the beta function $B(x, y)$, which are easily derivable and given in the following lemma.

Lemma 3.1. *The following formulas hold.*

$$(34) \quad B(x, y + 1) = \frac{y}{x + y} B(x, y);$$

$$(35) \quad B(x + 1, y) = \frac{x}{x + y} B(x, y);$$

$$(36) \quad B_p^{\lambda, \rho}(x, y) = B_p^{\lambda, \rho}(x + 1, y) + B_p^{\lambda, \rho}(x, y + 1).$$

We give a derivative formula for the extended Mittag-Leffler function (9), which is asserted by the following theorem.

Theorem 3.2. *Let $p \in \mathbb{R}_0^+$, $\Re(c) > \Re(\gamma) > 0$, $\Re(\alpha) > 0$, and $\rho \in \mathbb{C} \setminus \mathbb{Z}_0^-$. Then*

$$(37) \quad \frac{d^n}{dz^n} E_{\alpha, \beta}^{\gamma, c; \lambda, \rho}(z; p) = (\gamma)_n E_{\alpha, n\alpha + \beta}^{\gamma + n, c + n; \lambda, \rho}(z; p) \quad (n \in \mathbb{N}_0).$$

Proof. We find from (9) that

$$\begin{aligned} \frac{d}{dz} E_{\alpha, \beta}^{\gamma, c; \lambda, \rho}(z; p) &= \sum_{n=1}^{\infty} \frac{B_p^{\lambda, \rho}(\gamma + n, c - \gamma)}{B(\gamma, c - \gamma)} \frac{(c)_n}{\Gamma(\alpha n + \beta)} \frac{z^{n-1}}{(n-1)!} \\ &= \sum_{n=0}^{\infty} \frac{B_p^{\lambda, \rho}(\gamma + 1 + n, c - \gamma)}{B(\gamma, c - \gamma)} \frac{(c)_{n+1}}{\Gamma(\alpha n + \alpha + \beta)} \frac{z^n}{n!} \\ &= c \sum_{n=0}^{\infty} \frac{B_p^{\lambda, \rho}(\gamma + 1 + n, c - \gamma)}{B(\gamma, c - \gamma)} \frac{(c+1)_n}{\Gamma(\alpha n + \alpha + \beta)} \frac{z^n}{n!}, \end{aligned}$$

which, upon using (35), leads to

$$\frac{d}{dz} E_{\alpha, \beta}^{\gamma, c; \lambda, \rho}(z; p) = \gamma \sum_{n=0}^{\infty} \frac{B_p^{\lambda, \rho}(\gamma + 1 + n, c - \gamma)}{B(\gamma + 1, c - \gamma)} \frac{(c+1)_n}{\Gamma(\alpha n + \alpha + \beta)} \frac{z^n}{n!}.$$

In terms of (9), we obtain

$$(38) \quad \frac{d}{dz} E_{\alpha, \beta}^{\gamma, c; \lambda, \rho}(z; p) = \gamma E_{\alpha, \alpha + \beta}^{\gamma + 1, c + 1; \lambda, \rho}(z; p).$$

Similarly, we have

$$\frac{d^2}{dz^2} E_{\alpha, \beta}^{\gamma, c; \lambda, \rho}(z; p) = \gamma(\gamma + 1) E_{\alpha, 2\alpha + \beta}^{\gamma + 2, c + 2; \lambda, \rho}(z; p).$$

Then, continuing this process $n - 2$ times on the last formula is seen to yield the desired result (37). \square

Theorem 3.3. Let $p \in \mathbb{R}_0^+$, $\Re(c) > \Re(\gamma) > 0$, $\Re(\alpha) > 0$, $\rho \in \mathbb{C} \setminus \mathbb{Z}_0^-$, and $\beta, \mu \in \mathbb{C}$. Then

$$(39) \quad \frac{d^n}{dz^n} \left\{ z^{\beta-1} E_{\alpha,\beta}^{\gamma,c;\lambda,\rho}(\mu z^\alpha; p) \right\} = z^{\beta-n-1} E_{\alpha,\beta-n}^{\gamma,c;\lambda,\rho}(\mu z^\alpha; p) \quad (n \in \mathbb{N}_0).$$

Proof. By using (9), it is easy to get the desired result. The detailed account of the proof is omitted. \square

Theorem 3.4. Let $p \in \mathbb{R}_0^+$, $x \in \mathbb{R}^+$, and $\rho \in \mathbb{C} \setminus \mathbb{Z}_0^-$. Also, let $m - 2 < \Re(\delta - c) < m - 1$ for $m \in \mathbb{N}$. Then

$$(40) \quad \begin{aligned} & \mathfrak{D}_{x;\lambda,\rho}^{\delta+1-c,p} \left\{ x^{\delta-1} E_{\alpha,\beta}^{c+m-1}(x) \right\} \\ &= \sum_{\ell=0}^m \binom{m}{\ell} \frac{\Gamma(\delta + \ell)}{\Gamma(c + \ell - 1)} x^{c+\ell-2} E_{\alpha,\ell\alpha+\beta}^{\delta+\ell,c+\ell+m-1;\lambda,\rho}(x; p) \quad (m \in \mathbb{N}). \end{aligned}$$

Proof. Let \mathcal{L}_1 be the left side of (40). By using (32), we have

$$\begin{aligned} \mathcal{L}_1 &= \frac{1}{\Gamma(m + c - \delta - 1)} \frac{d^m}{dx^m} \int_0^x t^{\delta-1} (x-t)^{c+m-\delta-2} \\ &\quad \times E_{\alpha,\beta}^{c+m-1}(t) {}_1F_1 \left[\lambda; \rho; -\frac{px^2}{t(x-t)} \right] dt, \end{aligned}$$

which, upon taking $t = ux$, becomes

$$(41) \quad \begin{aligned} \mathcal{L}_1 &= \frac{1}{\Gamma(m + c - \delta - 1)} \frac{d^m}{dx^m} \left\{ x^{c+m-2} \int_0^1 u^{\delta-1} (1-u)^{c+m-\delta-2} \right. \\ &\quad \left. \times E_{\alpha,\beta}^{c+m-1}(xu) {}_1F_1 \left[\lambda; \rho; -\frac{p}{u(1-u)} \right] du \right\}. \end{aligned}$$

Applying (15) to the integral in (41), we obtain

$$\mathcal{L}_1 = \frac{\Gamma(\delta)}{\Gamma(c + m - 1)} \frac{d^m}{dx^m} \left\{ x^{c+m-2} E_{\alpha,\beta}^{\delta,c+m-1;\lambda,\rho}(x; p) \right\}.$$

By using the Leibniz's generalized rule for differentiation of product of two functions, we have

$$(42) \quad \mathcal{L}_1 = \frac{\Gamma(\delta)}{\Gamma(c + m - 1)} \sum_{\ell=0}^m \binom{m}{\ell} \left\{ \frac{d^{m-\ell}}{dx^{m-\ell}} x^{c+m-2} \right\} \left\{ \frac{d^\ell}{dx^\ell} E_{\alpha,\beta}^{\delta,c+m-1;\lambda,\rho}(x; p) \right\}.$$

Using (37) and the following easily derivable formula

$$(43) \quad \frac{d^\ell}{dx^\ell} x^\alpha = \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha - \ell + 1)} x^{\alpha-\ell} \quad (\ell \in \mathbb{N}_0)$$

in (42), we get the desired result. \square

The particular case of Theorem 3.4 when $m = 1$ is given in the following corollary.

Corollary 3.5. *Let $p \in \mathbb{R}_0^+$, $x \in \mathbb{R}^+$, $\rho \in \mathbb{C} \setminus \mathbb{Z}_0^-$, $0 < \Re(\delta) < \Re(c)$ and $\Re(\delta - c) > -1$. Then*

$$(44) \quad \begin{aligned} & \mathfrak{D}_{x;\lambda,\rho}^{\delta+1-c,p} \{x^{\delta-1} E_{\alpha,\beta}^c(x)\} \\ &= \frac{\Gamma(\delta)}{\Gamma(c)} x^{c-2} \left\{ (c-1) E_{\alpha,\beta}^{\delta,c;\lambda,\rho}(x;p) + \delta x E_{\alpha,\alpha+\beta}^{\delta+1,c+1;\lambda,\rho}(x;p) \right\}. \end{aligned}$$

References

- [1] M. A. Chaudhry, A. Qadir, H. M. Srivastava, and R. B. Paris, *Extended hypergeometric and confluent hypergeometric functions*, Appl. Math. Comput. **159** (2004), no. 2, 589–602.
- [2] J. Choi and P. Agarwal, *A note on fractional integral operator associated with multiindex Mittag-Leffler functions*, Filomat **30** (2016), no. 7, 1931–1939.
- [3] J. Choi and D. Kumar, *Solutions of generalized fractional kinetic equations involving Aleph functions*, Math. Commun. **20** (2015), no. 1, 113–123.
- [4] C. Fox, *The asymptotic expansion of generalized hypergeometric functions*, Proc. London Math. Soc. (2) **27** (1928), no. 5, 389–400.
- [5] R. Gorenflo, A. A. Kilbas, and S. V. Rogosin, *On the generalized Mittag-Leffler type functions*, Integral Transform. Spec. Funct. **7** (1998), no. 3-4, 215–224.
- [6] R. Hilfer, *Applications of Fractional Calculus in Physics*, World Scientific Publishing Co., Inc., River Edge, NJ, 2000.
- [7] ———, *Fractional time evolution*, in Applications of fractional calculus in physics, 87–130, World Sci. Publ., River Edge, NJ, 2000.
- [8] R. Hilfer and H. J. Seybold, *Computation of the generalized Mittag-Leffler function and its inverse in the complex plane*, Integral Transforms Spec. Funct. **17** (2006), no. 9, 637–652.
- [9] A. A. Kilbas and M. Saigo, *On Mittag-Leffler type function, fractional calculus operators and solutions of integral equations*, Integral Transform. Spec. Funct. **4** (1996), no. 4, 355–370.
- [10] A. A. Kilbas, H. M. Srivastava, and J. J. Trujillo, *Theory and Applications of Fractional Differential Equations*, North-Holland Mathematics Studies, **204**, Elsevier Science B.V., Amsterdam, 2006.
- [11] M. Kurulay and M. Bayram, *Some properties of the Mittag-Leffler functions and their relation with the Wright functions*, Adv. Difference Equ. **2012** (2012), 181, 8 pp.
- [12] F. Mainardi, *On some properties of the Mittag-Leffler function $E_\alpha(-t^\alpha)$, completely monotone for $t > 0$ with $0 < \alpha < 1$* , Discrete Contin. Dyn. Syst. Ser. B **19** (2014), no. 7, 2267–2278.
- [13] G. M. Mittag-Leffler, *Sur la nouvelle fonction $E_\alpha(x)$* , C. R. Acad. Sci. Paris **137** (1903), 554–558.
- [14] E. Özergin, M. A. Özarslan, and A. Altin, *Extension of gamma, beta and hypergeometric functions*, J. Comput. Appl. Math. **235** (2011), no. 16, 4601–4610.
- [15] M. A. Özarslan and B. Yilmaz, *The extended Mittag-Leffler function and its properties*, J. Inequal. Appl. **2014** (2014), 85, 10 pp.
- [16] T. R. Prabhakar, *A singular integral equation with a generalized Mittag Leffler function in the kernel*, Yokohama Math. J. **19** (1971), 7–15.
- [17] M. Saigo and A. A. Kilbas, *On Mittag-Leffler type function and applications*, Integral Transform. Spec. Funct. **7** (1998), no. 1-2, 97–112.
- [18] S. G. Samko, A. A. Kilbas, and O. I. Marichev, *Fractional Integrals and Derivatives*, translated from the 1987 Russian original, Gordon and Breach Science Publishers, Yverdon, 1993.

- [19] H. J. Seybold and R. Hilfer, *Numerical results for the generalized Mittag-Leffler function*, *Fract. Calc. Appl. Anal.* **8** (2005), no. 2, 127–139.
- [20] A. K. Shukla and J. C. Prajapati, *On a generalization of Mittag-Leffler function and its properties*, *J. Math. Anal. Appl.* **336** (2007), no. 2, 797–811.
- [21] H. M. Srivastava, *A contour integral involving Fox's H-function*, *Indian J. Math.* **14** (1972), 1–6.
- [22] ———, *A note on the integral representation for the product of two generalized Rice polynomials*, *Collect. Math.* **24** (1973), 117–121.
- [23] H. M. Srivastava and J. Choi, *Zeta and q-Zeta Functions and Associated Series and Integrals*, Elsevier, Inc., Amsterdam, 2012.
- [24] H. M. Srivastava, K. C. Gupta, and S. P. Goyal, *The H-functions of one and two variables*, South Asian Publishers Pvt. Ltd., New Delhi, 1982.
- [25] H. M. Srivastava and C. M. Joshi, *Integral representation for the product of a class of generalized hypergeometric polynomials*, *Acad. Roy. Belg. Bull. Cl. Sci. (5)* **60** (1974), 919–926.
- [26] H. M. Srivastava and P. W. Karlsson, *Multiple Gaussian Hypergeometric Series*, Ellis Horwood Series: Mathematics and its Applications, Ellis Horwood Ltd., Chichester, 1985.
- [27] H. M. Srivastava and Ž. Tomovski, *Fractional calculus with an integral operator containing a generalized Mittag-Leffler function in the kernel*, *Appl. Math. Comput.* **211** (2009), no. 1, 198–210.
- [28] A. Wiman, *Über den Fundamentalsatz in der Theorie der Funktionen $E^a(x)$* , *Acta Math.* **29** (1905), no. 1, 191–201.
- [29] ———, *Über die Nullstellen der Funktionen $E^a(x)$* , *Acta Math.* **29** (1905), no. 1, 217–234.
- [30] E. M. Wright, *The asymptotic expansion of the generalized hypergeometric functions*, *J. London Math. Soc.* **10** (1935), 286–293.
- [31] ———, *The asymptotic expansion of integral functions defined by Taylor series*, *Philos. Trans. Roy. Soc. London, Ser. A.* **238** (1940), 423–451.
- [32] ———, *The asymptotic expansion of the generalized hypergeometric function*, *Proc. London Math. Soc. (2)* **46** (1940), 389–408.

MUHAMMAD ARSHAD
 DEPARTMENT OF MATHEMATICS
 INTERNATIONAL ISLAMIC UNIVERSITY
 ISLAMABAD, PAKISTAN
Email address: marshad_zia@yahoo.com

JUNESANG CHOI
 DEPARTMENT OF MATHEMATICS
 DONGGUK UNIVERSITY
 GYEONGJU 38066, KOREA
Email address: junesang@mail.dongguk.ac.kr

SHAHID MUBEN
 DEPARTMENT OF MATHEMATICS
 UNIVERSITY OF SARGODHA
 SARGODHA, PAKISTAN
Email address: smjhanda@gmail.com

KOTTAKKARAN SOOPPY NISAR
DEPARTMENT OF MATHEMATICS
COLLEGE OF ARTS AND SCIENCE AT WADI AL-DAWASER
PRINCE SATTAM BIN ABDULAZIZ UNIVERSITY
ALKHARJ, RIYADH REGION 11991, KINGDOM OF SAUDI ARABIA
Email address: knsisar1@gmail.com

GAUHAR RAHMAN
DEPARTMENT OF MATHEMATICS
INTERNATIONAL ISLAMIC UNIVERSITY
ISLAMABAD, PAKISTAN
Email address: gauhar55uom@gmail.com